

Effects of Jatropha Seed Cake (JSC) and Different Inorganic Fertilizers on Growth and Yield of Two Groundnut Cultivars under Three Harvesting Periods.

S. M. H Elseed¹, S. O.Yagoub^{2*}, I. S. Mohamed³

¹ Ministry of Agriculture and Animal Resources and Irrigation, Khartoum State, Sudan.

^{2*}Department of Agronomy, College of Agricultural Studies, Sudan University of Science and Technology Box, 73,, Sudan. *Email of corresponding author: umreelah2003@yahoo.com

³Department of Plant Protection, College of Agricultural Studies, Sudan University of Science and Technology, Sudan.

Authors' contributions

This work was carried out in collaboration between three authors. Author SMHE initiated the experiments, collected the data, performed the statistical analysis, Author SOY designed the study, managed the literature review and wrote the first drafts of the manuscript. Author ISM assisted with statistical analysis and contributed to the final draft. Three authors read and approved the final manuscript.

ABSTRACT

Aim: to investigate the effect of jatropha seed cake and different fertilizers on growth and yield of two groundnut cultivars under three harvesting periods

Study Design: The experimental design was a randomized complete block in a split-plot replicated three times. Groundnut cultivars arranged as whole plots and five fertilizers treatment as sub plot.

Locations: field experiments were conducted for two seasons 2011/12-2012/13 under semi-arid condition of Khartoum state. in demonstration farm of Sudan University of Science and Technology, Khartoum North, Shambat.

Methodology: The treatments were Jatropha seed cake (2.5 ton/ha), sulphur (119 kg/ha), single super phosphate (119 kg/ha), ammonium sulphate(119 kg/ha) and control on two groundnut cultivars (Sodri) and (Gebish) under three harvesting time(every 10 days).

Results: The results showed that yield increased with delayed in harvest periods for two seasons. The two tested cultivars revealed similar behavior during vegetative and reproductive growth. Except for plant height of third harvest of first season, primary branch of second harvest and yield of first and third harvest of two seasons. Results showed significant difference among fertilizers treatments and interaction between cultivars and fertilizers concerning the majority of growth and yield components. Super phosphate gave significant difference on leaf area index, plant height, number of primary branches, number of primary branches, number of pods/plant, germination %. On the other hand, sulphur and ammonium sulphate gave significant difference on number of pods/plant, germination % and yield. Ammonium sulphate and jatropha seed cake reviled significant effect on yield kg/ha in first and second season respectively

Conclusion: In conclusion addition of jatropha seed cake and inorganic fertilizers appeared to be more promising. Delaying harvesting periods gave positive results.

Keywords:; groundnut inorganic fertilizers, jatropha seed cake

1.INTRODUCTION

Groundnut *Arachis hypogaea* L is a major cash crop grown mainly under rain-fed conditions in the semi-arid tropics. Groundnut is the 13th most important food crop of the world. It is the world's 4th most important source of edible oil and 3rd most important source of vegetable protein {1}. Groundnut is primarily used for oil extraction in Sudan {2}. It is consumed directly because of its high food

value. Sudan is one of the major groundnut producing countries. The total area under groundnut production is approximately one million ha with an average yield of 855 kg ha⁻¹ {2}.

Substantial evidence showed that groundnut responds well to additional inorganic and organic fertilizers application {3, 4, 5, 6, 7, 8, 9}. Phosphorus is very important nutrient element for crop growth and yield. It plays an important role in physiological processes of plants. As P source, single superphosphate is the most suitable fertilizer for groundnut in Nigeria Savannah {10}. Sulphur deficiency in legume crops affects not only yield, but also the nutritional quality of the seeds, {11}.

Biodiesel is an environmental-friendly substitute for fossil fuel. Research on the production of biodiesel from vegetable oils has concentrated on *Jatropha*, soybean, palm kernel, sunflower and rapeseed oils with scarce information on groundnut oils {12}. The biodiesel from *Jatropha curcas* had received much attention in past decades. The byproduct of oil extraction from *Jatropha curcas* seeds and kernel is called seedcake. It contains highly toxic protein that is not suitable for animal feeds, although good for organic manure, it is being used as an organic fertilizer {13}. The cake is rich in nitrogen (3.2%) phosphorus (1.4%) and potassium (1.3%) and can be used as manures {14 and 15}. Limited studies have been reported on the potential of *Jatropha* seed cake to be used as organic fertilizers. Organic manures are very important, as they contain both major and minor elements necessary for plant growth, and improve the physical, chemical and biological properties of soil {16}. The productivity of groundnut depends on proper selection of variety, fertilizer management. This study was carried out to identify the effect of *Jatropha* seed cake (JSC) and different types of fertilizers on growth and yield of two groundnut cultivars under three harvesting periods.

2. MATERIALS AND METHODS.

The experiment was aimed to study the effect of different organic and inorganic fertilizers on growth of two groundnut cultivars under different harvesting periods.

2.1 Experimental site

Field experiments were conducted in Demonstration farm, College of Agricultural Studies, Sudan University of Science and Technology, Shambat, Khartoum North, Sudan for two consecutive seasons 2011/12- 2012/13. Shambat is located at longitude 32°35'E and latitude 15°31'N, within the semi-desert region. Climate of the locality is semi-desert and tropical with low relative humidity.

2.2 Layout of the Experiment and Land Preparation

The experiment was designed in split plots replicated three times. Two cultivars of groundnut (Sodri =V1) and (Gebish=V2) and four types of fertilizers were applied; (F1) control without fertilizer, (F2) inorganic Jatropha Seed Cake (JSC) of 2.5 tons/ha, and three organic fertilizers: (F3) pure sulfate of 119 kg/ha, (F4) super phosphate of 119 kg/ha and (F5) Ammonia sulfate of 119 kg/ha. After maturing(about 110 days from sowing) three harvesting periods were taken (every 10 days), early, medium and late. The field was prepared according to adopted method by Engineering Department of Agricultural College. The seeds were sown in July 12th, and July 21th respectively for two seasons, 3-4 seeds per hole then thinned to two seed per hole. During July until August (about two months) it was irrigated by rain-fed then irrigation was applied every 15 days. For first season the three harvesting time were, October, 11- 21 and first November, and for second season October, 24, and November 4-14.

2.3 The Source of Seeds:

Two cultivars of groundnut certified seed (Sodri = V1) and (Gebish= V2) were obtained from Al –Obied Research Center (North Kordfan) , and Arabian Company for Seed, Khartoum.

2. 3 Data Collection

Observation was taken from 3 plants selected randomly each subplot. Data was recorded on pre and post harvesting stage. The data recorded during pre-harvest; were percentage of germination after 3 weeks and 5 weeks, seeds that failed to germinate were counted per hole (implanting seeds), Leaf area index after 90 days. The harvest stage data were: - height of plant, number of primary and secondary branches, and number of pods per plant. The yield of threshed seeds from one meter of each plots were taken and transformed in kg/ha.

2-4 Statistical Analysis:

The collected data were subjected to standard Statistical analysis. The procedure of analysis of variance and mean separation were followed according to the description of {17}. The data was analyzed by MSTAT-C Statistical Package.

4. RESULTS AND DISCUSSION

The effect of Jatropha Seed Cake (JSC), sulfur, super phosphate and ammonium sulfate fertilizer on germination % of two groundnut cultivars (Sodri and Gebish) were presented in Table (1a) for season 2011/12 and Table (1b) season 2012/13. There were no significant difference between two cultivars for 3 and 5 weeks of sowing, and among the fertilizers for 5 weeks of sowing, but there were significant differences among fertilizers X cultivars for two seasons for 3 and 5 days of sowing. Sulfur fertilizer with V1(Sodri) and super phosphate with V2 (Gebish) gave the highest values for first and second seasons respectively. Among

fertilizers for two seasons in 3 weeks of sowing application of sulfur and (JSC) obtained the highest values. This evaluated that groundnut responded well to application of organic and inorganic fertilizers. This was in line with {9}.

Table 1a. Effect of Jatropha seed cake and different fertilizers on germination % of two groundnut cultivars season 2011/12.

	3 WS			5WS		
	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X
F1(Control)	70.0 b	69.0 b	69.5 ab	77.5 bc	81.3 b	79.4 a
F2(Jateropha)	65.0 bc	64.6 bc	64.8 ab	71.3 c	76.8 bc	74.1 a
F3(Sulfur)	82.9 a	58.7 c	70.8 a	89.2 a	69.5 c	79.4 a
F4(Super ph.)	66.7 bc	63.8 bc	65.2 ab	75.2 bc	71.7 c	73.4 a
F5(Amonium sulfate)	69.6 b	49.6 d	59.6 b	70.6 c	72.8 bc	71.7 a
X	70.8 a	61.1 a		76.8 a	74.4 a	75.6
CV%			21.5			18.9
LSD V			7.8			7.8
LSD F			10.2			10.2
LSD V*F			62.7			54.1

Table 1b. Effect of Jatropha seed cake and different fertilizers on germination % of two groundnut cultivars season 2012/13.

	3 WS			5WS		
	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X
F1(Control)	45.8 c	60.3 b	53.1 b	36.3 c	50.4 ab	43.4 a
F2(Jateropha)	54.6 b	60.4 b	59.5 a	46.9 b	50.7 ab	48.8 a
F3(Sulfur)	41.9 c	65.4 b	53.7 b	31.7 c	50.5 ab	41.1 a
F4(Super ph.)	46.1 c	68.9 a	57.5 ab	37.2 c	56.8 a	47.0 a
F5(Amonium sulfate)	57.5 b	57.5 b	57.5 ab	46.4 b	46.4 b	46.4 a

X	49.9 a	62.5 a	39.7 a	51.0 a
CV%		31%		34.6%
LSD V		4.86		11.30
LSD F		9.63		8.57
LSD V*F		26.7		15.00

The results of leaf area index presented in Table 2, there were significant difference in first season between cultivars, fertilizers and interaction among cultivars and fertilizers. V2, F4 (super phosphate) and the interaction of V2XF4 gave the highest values. While the second season the significant differences were noticed only among the interaction of cultivars and fertilizers and V1XF4 showed the highest values.

Table 2. Effect of Jatropha seed cake (JSC) and different fertilizers on leaf area index of two groundnut cultivars seasons 2011/12-2012/13.

	2011/12			2012/13		
	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X
F1(Control)	29.2 efg	30.6 def	29.9 b	34.7 b	29.3 de	31.9 a
F2(Jateropha)	32.3 cd	32.6 cd	32.5 ab	38.1 a	28.7 e	33.4 a
F3(Sulfur)	35.2 ab	30.6 def	33.4 a	36.2 ab	31.7 cd	33.9 a
F4(Super ph.)	27.8 9	36.4 a	32.1 ab	34.0 bc	28.8 e	31.4 a
F5(Amonium sulfate)	28.4 fg	33.8 bc	31.1 ab	34.4 b	30.3 de	32.3 a
X	30.6 b	32.9 a		35.5 a	29.7 a	
CV%	13.2			14.1		
LSD V			1.5			2.5
LSD F			2.3			3.3
LSD V*F			3.0			13.7

Tables 3a-3b showed the results of effect of JSC and different fertilizers on plant height of two groundnut cultivars for three harvesting periods. The results indicated non significant differences between cultivars except in third harvest of second season and V1 (Sodri) gave the highest value. Among fertilizers treatments all of them revealed highly significant difference except the first harvest of second season. In general, F4 (super phosphate fertilizers) showed the highest records in first harvest of first season and second harvest of second season also with control in second and third harvest of first season, but in second season third harvest F2(JSC) showed the highest value. The interaction of cultivars with fertilizers showed significant difference for all parameters and F4 and F1 obtained the highest values except in the third harvest of second season F2 gave the higher value.

The results showed the effect of JSC and different fertilizers on number of primary branches of two groundnut cultivars seasons 2011/12-2012/13. It was presented in Table 4a-4b for two seasons respectively. The results obtained that there were no significant differences between cultivars for three different harvesting time except in season two second harvest and V1(Sodri) gave higher record than V2(Gebish). Among fertilizers in the first season the results showed that F2 (JSC) in first and third harvest and F4 (Super phosphate) gave the highest values with significant differences. Meanwhile, second season showed slight difference in first and second harvest and no significant difference in third harvest for fertilizers treatment. The results of interaction of cultivars with fertilizers showed that F2 for two cultivar in first harvest and V1F4 second harvest and V2 F2 for thirds harvest had the highest values with significant difference. The interaction in season two showed results similar to results of fertilizers with difference in V1 which gave the biggest values in first and second harvest, but thirds harvest gave significant difference in V1F4.

Numbers of pods/plants were obtained in table 5a-5b. The results showed non significant differences between cultivars for two seasons in all harvest time. Fertilizer treatment showed no significant differences in first harvest of first season and second and third harvest of second season. In first season super phosphate gave the significant difference among treatments. In second season F5 gave the highest values. {18} study the relationships between nutrient uptake parameters, biomass and pod dry weight and found that they were positive and significant in both seasons. The interaction treatment revealed highly significant difference for three harvests for two seasons. In general third harvest of two seasons had the highest number of pods/plant and V2F4 of third harvest of second season obtained the heaviest numbers of pods/plant which was 18.3 pods/plant. {19} gave the same result and showed that the genotypes obtained an increasing trend in the number of mature pods as harvesting date delayed.

The yield kg/ha of two groundnut cultivars for three harvests were presented in table 6a-6b for two seasons. The result showed that yield increased with advanced in harvest time for two seasons, and the yield of third harvest is the best of the first and second harvest this result was in line with {19} but {20} found that peanut harvest at physiological maturing period gave better result than

that at 10 days earlier or later.. V2 (Gebish) gave slight big yield compared with V1 (Sodri) with significant difference in first and third harvest of first season and first harvest of second seasons. Among fertilizers treatment in second seasons Tables 6b the results showed that F2 for three harvests gave the highest yield and also for interaction had the same results with variation in cultivars V2F2 gave the highest values. In first season Table 6a, F5 (Amonium sulphate) gave the highest values for three harvest. The interaction of cultivars and fertilizers in this season gave significant difference. The same results among fertilizers treatments were detected{21}.

Table 3a. Effect of Jatropha seed cake and different fertilizers on plant height of two groundnut cultivars season 2011/12.

	First harvest			Second harvest			Third harvest		
	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X
F1(Control)	18.1 cd	19.8 bc	18.9 b	21.3 c	23.3 b	22.3 a	21.3 cde	26.1 a	23.7 a
F2(Jateropha)	20.4 b	17.3 d	18.8 b	19.6 def	20.4 cde	19.9 b	21.9 cd	20.3 ef	21.1 b
F3(Sulfur)	18.7 bcd	20.2 b	19.4 b	19.2 ef	22.8 b	21.0 ab	20.1 ef	19.6 f	19.9 b
F4(Super ph.)	22.3 a	23.3 a	22.8 a	19.5 ef	24.7 a	22.1 a	22.7 bc	23.6 b	23.2 a
F5(Amonium sulfate)	19.5 bc	19.8 bc	19.7 b	18.9 f	21.0 cd	19.9 b	20.7 def	19.8 ef	20.2 b
X	19.8 a	20.1 a		19.7 a	22.4 a		21.3 a	21.9 a	
CV%			16.4			11.7			12.2
LSD V			1.79			1.35			1.45
LSD F			2.3			1.77			1.89
LSD V*F			3.6			9.4			2.7

Table 3b. Effect of Jatropha seed cake and different fertilizers on plant height of two groundnut cultivars season 2011/12.

	First harvest			Second harvest			Third harvest		
	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X
F1(Control)	19.0 d	22.5 bc	20.8 a	18.1 d	21.6 c	19.9 cd	18.8 e	24.7 bc	21.8 ab
F2(Jateropha)	20.5 cd	25.9 a	23.2 a	21.5 c	26.1 b	23.8 ab	21.6 d	27.4 a	24.5 a

F3(Sulfur)	23.1 b	22.5 bc	22.8 a	18.6 d	25.9 b	22.3 bc	18.6 e	25.6 ab	22.1 ab
F4(Super ph.)	20.4 cd	24.1 ab	22.2 a	23.0 c	29.6 a	26.3 a	23.2 cd	24.1 bc	23.7 ab
F5(Amonium sulfate)	21.8 bc	21.7 bc	21.7 a	17.9 d	18.8 d	18.4 d	21.7 d	21.7 d	21.7 b
X	20.9 a	23.3 a		19.9 a	24.4 a		20.8 b	24.7 a	
CV%			19.2			21.3			16.3
LSD V			0.78			2.52			2.66
LSD F			2.33			3.31			2.03
LSD V*F			3.05			7.60			13.7

Different concentration of JSC gave the various results on the yield of each vegetable crop. Treatment with the half rate of chemical fertilizer mixed with high rate (10 t/ha) of JSC had proven to provide the best marketable yield of Chinese kale, in which such yield was comparable to that of full rate of chemical fertilizer. In addition, the half rate of chemical fertilizer combined with any rates of JSC (low, medium, and high) gave the most tomato fruit yield. When applied with low rate (2.5 t/ha) of JSC to the sweet potato plot, the outcome showed the highest tuber yield [22]. As mentioned above, fertilizers in form of jatropha seed cake, super phosphate, sulphur and ammonium sulphate on different yield and yield gave positive results with significant difference. {6} concluded that phosphorus rate increased the number of filled pods and seed yield. Also {23} found that application of FYM and S fertilization increased the yield and yield attributed of groundnut. {24} confirmed that sulphur application increased pod yield of groundnut. On the other hand, {25} found that sulphur application significantly influenced the growth and yield attributed characters, yield and oil content over control regardless of the source and levels of sulphur. In contrast {3} found that sulphur supply only increase the sulphur concentration of the plants without enhancing the yield.

Table 4a. Effect of Jatropha seed cake and different fertilizers on number of primary branches of two groundnut cultivars season 2011/12.

	First harvest			Second harvest			Third harvest		
	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X
F1(Control)	4.5 bc	4.1 c	4.5 bc	4.1 d	4.5 cd	4.3 c	5.0 bc	4.8 bcd	4.9 ab

F2(Jateropha)	5.6 a	5.3 a	5.6 a	4.7 c	5.2 b	4.9 ab	5.1 b	5.7 a	5.4 a
F3(Sulfur)	5.1 ab	3.9 c	5.1 ab	4.7 d	5.2 b	4.9 ab	4.3 de	4.9 bc	4.6 b
F4(Super ph.)	4.5 bc	4.2 c	4.5 bc	5.8 a	4.6 cd	5.1 a	4.9 bc	4.5 cde	4.7 b
F5(Amonium sulfate)	4.5 bc	4.5	4.5 bc	4.4 cd	4.6 cd	4.5 bc	4.8 bcd	4.2 c	4.5 b
X	4.8 a	4.4 a		4.7 a	4.8 a		4.8 a	4.8 a	
CV%			24			17			16.5
LSD V			0.61			0.44			0.43
LSD F			0.79			0.58			0.57
LSD V*F			2.5			2.6			1.5

Table 4b. Effect of Jatropha seed cake and different fertilizers on number of primary branches of two groundnut cultivars season 2012/13.

	First harvest			Second harvest			Third harvest		
	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X
F1(Control)	6.1 a	4.7 cd	5.4 a	6.2 a	4.3 de	5.3 a	5.0 bc	4.6 cd	4.8 a
F2(Jateropha)	5.3 b	3.8 e	4.6 b	4.2 e	3.9 e	4.0 b	5.4 abc	5.2 bc	5.3 a
F3(Sulfur)	4.1 de	5.4 b	4.8 ab	5.3 bc	3.8 e	4.6 ab	5.8 ab	5.0 bc	5.4 a
F4(Super ph.)	4.5 cd	5.0 bc	4.7 ab	4.9 cd	5.2 bc	5.1 a	6.3 a	4.0 d	5.2 a
F5(Amonium sulfate)	6.2 a	4.1 de	5.2 ab	5.7 ab	5.0 bcd	3.3 a	5.8 ab	5.1 bc	5.5 a
X	5.2 a	4.6a		5.3 a	4.4 b		5.7 a	5.7 a	
CV%			20.2			24.0			32.5
LSD V			0.71			0.83			0.93
LSD F			0.54			0.63			1.21
LSD V*F			2.20			0.66			2.31

Taable5a. Effect of Jatropha seed cake and different fertilizers on number of pods/plant of two groundnut cultivars season 2011/12.

	First harvest			Second harvest			Third harvest		
	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X
F1(Control)	5.4 cd	8.1 a	6.7 a	8.0 b	5.0 d	6.5 b	13.1 bc	10.9 cd	12.0 ab
F2(Jateropha)	7.0 abcd	7.5 ab	7.3 a	5.5 cd	8.1 b	6.8 b	10.6 cde	12.0 cd	11.3 b
F3(Sulfur)	5.5 cd	5.3 cd	5.4 a	7.6 bc	8.0 b	7.8 ab	9.6 de	15.8 b	12.7 ab
F4(Super ph.)	5.6 bcd	5.0 d	5.3 a	9.0 ab	11.0 a	10.0	22.8 a	7.7 e	15.2 a
F5(Amonium sulfate)	7.1 abc	5.3 cd	6.2 a	8.7 ab	7.2 bcd	7.9 ab	13.4 bc	9.3 de	11.5 ab
X	6.1 a	6.2 a		7.8 a	7.8 a		13.9 a	11.2 a	
CV%			53.3			51.5			40.7
LSD V			1.80			2.10			2.80
LSD F			1.55			2.20			3.66
LSD V*F			2.37			4.10			7.80

Table 5b. Effect of Jatropha seed cake and different fertilizers on number of pods/plant of two groundnut cultivars season 2012/13.

	First harvest			Second harvest			Third harvest		
	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X
F1(Control)	5.9 d	9.1 b	7.5 bc	12.5 bc	15.8 a	14.2 a	12.4 cd	17.9 a	15.1 a
F2(Jateropha)	7.4 bcd	6.8 cd	7.1 c	11.5 c	14.6 ab	13.1 a	14.4 bc	14.4 bc	14.4 a
F3(Sulfur)	8.4 bc	10.9 b	9.7 ab	11.2 c	15.0 a	13.1 a	11.9 cd	14.9 b	13.4 a
F4(Super ph.)	10.9 a	7.8 bc	9.3 ab	11.9 c	15.7 a	1.3.8 a	11.9 d	18.3 a	15.1
F5(Amonium sulfate)	10.9 a	10.9 a	10.9 a	11.7 c	14.3 ab	12.9	15.4 b	10.3 d	12.9 a
X	8.7 a	9.1 a		11.8 a	15.1 a		13.2 a	15.2 a	
CV%			33.5			26			29.3
LSD V			6.60			2.00			2.27

LSD F	8.89	2.63	2.97
LSD V*F	14.53	7.10	23.9
LSD V*F	14.53	7.10	23.9

Table 6a. Effect of Jatropha seed cake and different fertilizers on yield (kg/ha) of two groundnut cultivars season 2011/12.

	First harvest			Second harvest			Third harvest		
	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X
F1(Control)	250.2 f	445.7 de	347.9 c	309.3 f	695.5 cd	502.4 d	760.5 e	1224.0 cde	992.3 e
F2(Jateropha)	791.2 b	360.3 ef	575.8 ab	942.3 cd	450.8 ef	69.60 e	2675.3 ab	936.5 de	1805.9 c
F3(Sulfur)	967.3 a	455.7 cde	711.5 a	1331.5 a	705.0 de	1018.3 b	2710.5 ab	1621.8 c	2166.2 b
F4(Super ph.)	390.7 ef	595.0 cd	492.8 bc	430.0 f	895.2 cd	662.6 c	1331.0 cd	1569.3 c	1450.2 d
F5(Amonium sulfate)	831.0 ab	609.5 c	720.3 a	1119.5bc	1424.7 a	1272.1 a	2761.1 a	2254.5 b	2507.8 a
X	646.1 a	493.2 b		826.5 a	834.2 a		2047.7 a	1521.2 b	
CV%			50.7%			57.3%			49.5%
LSD V			132.0			2216			584.7
LSD F			340.9			572.2			150.0
LSD V*F			157.9			260.3			483.1

Table 6b. Effect of Jatropha seed cake and different fertilizers on yield (kg/ha) of two groundnut cultivars season 2011/12.

	First harvest			Second harvest			Third harvest		
	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X	V1(Sodri)	V2(Gebish)	X
F1(Control)	124.7 e	126.5 e	193.1 d	327.3 d	242.8 de	285.1 c	792.3 c	444.3 d	618.3 cd

F2(Jateropha)	62.7 f	677.5 a	370.1 a	89.5 f	1607.7 a	848.6 a	191.2 e	1721.2 a	956.2 a
F3(Sulfur)	90.0 g	377.2 b	193.1 d	85.7 f	713.7 b	399.7 b	142.7 e	1340.5 b	741.6 bc
F4(Super ph.)	242.7 a	119.5 e	186.8 c	162.7 ef	274.0 d	218.3 c	556.5 d	558.5 d	557.5 d
F5(Amonium sulfate)	325.0 c	196.7 d	260.8 b	512.7 c	432.5 c	472.6 b	719.7 c	794.8 c	757.3 b
X	152,7 a	299.5 b		235.6 a	654.19 a		480.5 a	971.9 a	
CV%			38.9%			35.5%			32.6%
LSD V			71.8			666.4			884.2
LSD F			18.5			172.1			228.3
LSD V*F			48.4			86.4			129.4

4. SUMMARY AND CONCLUSION

Groundnut is very important oil seed in the Sudan. The productivity of groundnut depends on proper selection of variety, fertilizer management and other management practices. Currently application of jatropha seed byproduct as oil diesel remain seed cake waste. This jatropha seed cake can be use as good green manure fertilizers. In this study using jatropha seed cake compared with different inorganic fertilizers; sulphur, superphosphate and ammonium sulphate gave significant difference in growth, yield and yield components of groundnut. On the other hand, delayed harvesting periods gave good results in growth and yield of groundnut. The two cultivars of groundnut obtained similar results without clear variation in their growth. Thus additional studies under more conductive condition are needed to better understand the role of jatropha seed cake plays as fertilizer to increase growth and yield of crops.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Taru VB, Kyagya IZ Mshelia SI. Profitability of Groundnut Production in Michika Local Government Area of Adamawa State, Nigeria. J.Agric. Sci. 2010; 1(1):25-29.
2. ARC(Agricultural Research Corporation) of Sudan. Annual report (2003-2010) Wad-Madani, Sudan.
3. Gupta, U.C., Mc Leads J.A Effects of various sources of sulphur on yield and sulphur concentrations of cereals and forages. Can. J. of Soil Sci. 1984; 64:169-1174.
4. Ashraf, M., Athar, H. R .Harris, P. J. C. Kwon, T.R. Some prospective strategies for improving crop salt tolerance. Adv. Agro. 2008;97:45-110
5. DAFF. Groundnut production guide line. Department of Agriculture, Forestry and Fisheries. Pretoria, South Africa 2010.
6. Shiyam, J.O.. Growth and yield response of groundnut (*Arachis hypogea*, L.) to plant densities and phosphorus on an ultisol in southeastern Nigeria. Libyan Agriculture Research Center Journal 2012; 1(4):211-214.
7. Moraditochae, M.. Effects of humic acid, foliar spraying and nitrogen fertilizer management on yield of peanut(*Arachis hypogea*, L.) in Iran. ARPN journal of Agriculture and Biological Science, 2012; 7(4):289-293.
8. Habbasha, S.F., M.H.Taha and N.A.Jafar. Effect of nitrogen fertilizer levels and zinc foliar application on yield, yield attributes and some chemical traits of groundnut. Research Journal of Agriculture and Biological Sciences, 2013; 9(1): 1-7.
9. Rumbidzai, D.K. and C.Mabwe (2014). Response of groundnut to inorganic fertilizer use in smallholder farming of Makonde, District, Zimbabwe. Journal of Biology Agriculture and healthcare, 2013; 4(7):78-82.
10. Lombin, G., L. Single, J.K. Yayock,. A decade of fertilizer research on groundnuts (*Arachis hypogea*, L.) in the Savanna zone of Nigeria. Fertilizer research, 1985; 6(2):157-170.
11. Arshad, J., Yong Sun Moon, Malik, Z.A. (.Enzyme activity assessment of peanut (*Arachis hypogea*, L.) under slow- release sulphur fertilization. Australian J. of crop science, 2010; 4(3):169-174.

12. Oluwatoyin O. O., A. I. Bamgboye. Production of biodiesel from groundnut (*Arachis hypogea*, L.) oil. Agricultural Engineering International: CIGR Journal, 2014; 16(1).
13. Srinophakun P. Prospect of Deoiled *Jatropha curcas* Seedcake as Fertilizer for Vegetables Crops – A Case Study Journal of Agricultural Science, 2012; 4 (3): 211-226
14. B. G. Keremane, G. V. Hegde, and Sheshachar, “Jatropha curcas: Production systems and uses”. BAIF, Tiptur, 2003.
15. Openshaw K., “A review of Jatropha curcas: an oil plant of unfulfilled Promise”, Biomass and Bioenergy. 2000; 19, 1-15.
16. Ganapathy G. Selvam, K. Sivakumar. Influence of seaweed extract as an organic fertilizer on the growth and yield of *Arachis hypogea* L. and their elemental composition using SEM–Energy Dispersive Spectroscopic analysis. Asian Pacific Journal of Reproduction 2014; 3(1): 18-22*

17. Gomez KA, Gomez A. A Statistical Procedures For Agricultural Research, 2nd edition, A Wiley Inter. Sci publication. (1984) 6 John Wiley and Son, New York

18. Junjittakarn, S. Pmratch, S. Jogoloy, W. Htoon, N. Singkhan, N. Vorasoot, B. Toomsan, C. C. Hdbrook, A. Patanothai. Nutrient uptake of peanut genotypes under different water regimes. International Journal of plant Production, 2013; 7(4):677-692.

19. Kaba, J.S., K. Ofori, and F.K. Kumaga. Inter- relationships of yield and yield components of yield at different stages of maturity in three groundnuts (*Arachis hypogaea* L) varieties. International J. of life sci. research 2014; 2(1):43-48.

20. Rahmianna, A.A., A. Taurfig and E. Yusnawan.. Pod yield and kernel quality of peanut grown under two different irrigation and two harvest times. Indonesian Journal of Agriculture, 2009; 2(2):103-109.

21. Migawer, E.A., and M.A.M. Soliman. Performance of two peanut cultivars and their response to NPK fertilization in newly reclaimed loamy sand soil. J. Agric. Sci. Mansoura Uni., 2001; 26(11):6653-6667.

22. Srinophakun, P. A Saimaneerat, I. Sooksathan, N Visarathanon, S. Malaipan, K Charernsom, W Chongrattanameteekul. World Renewable Energy Congress 2011-Sweden 8-13 May 2011. Linköping Sweden 232-238

23. Jat R.A. and Alhawat ,I.P.S. . Effect of organic manure and sulphur fertilization in pigeon pea(*Cajanus cajan*) and groundnut(*Arachis hypogea, L.*) intercropping system, 2010 55(4): 276-281.
24. Dash, A.K., Nayak, B.P.,Panigrany, N.,Mohapatra, S.Samant P.K. Performance of groundnut(*Arachis hypogea, L.*) under different levels of sulphur and irrigation. Indian J.of Agronomy, 2014; 58(4):578-582.
25. Tajeswara, K.R., A.Upendra Rao and D.Seuha. Effect of source and levels of sulphur on groundnut .Journal of academic and Industrial Research, 2013; 2(5):268-274.