

Original Research Article

Differentiated Contribution of Minerals through Soil and Foliar Fertilization to the Winter Wheat Yield

ABSTRACT

Research has evaluated the contribution of minerals through soil and foliar fertilization on winter wheat yield, the purpose being to identify the interdependence relations for the optimization of the fertilization system. Two types of mineral and foliar fertilizers were tested. The complex mineral fertilizer of the type NPK (1:1:1) was used for achieving three differentiated nutrition levels: NPK₀, NPK₃₀ (30 kg active substance ha⁻¹) and NPK₆₀ (60 kg active substance ha⁻¹). There are five types of foliar fertilizers applied and 18 experimental variants were studied. The research was conducted in Soil Science and Plant Nutrition (SSPN) and Mathematics and Statistics (MS), Banat University of Agricultural Sciences and Veterinary Medicine „Regele Mihai I al României„ from Timisoara, Romania (USAMVBT), between 2011 and 2012. The influence of interaction between the two types of fertilizer compared to their individual influence on wheat production was analyzed. Based on regression analysis were obtained mathematical model of interdependence, reliability of the results and models have been certified by LSD, p and R² (maximum value is better). The complex fertilizers generated yield increase between 925*** and 1425*** kg ha⁻¹. The foliar fertilizers determined yield increase between 355 and 810*** kg ha⁻¹ with natural soil fertility (NPK₀), between 550** and 900*** kg ha⁻¹ with NPK₃₀, and between 500** and 795*** kg ha⁻¹ with NPK₆₀. The yield increase determined by the joint effect of the two types of fertilization was between 1475*** and 2220*** kg ha⁻¹. The experimental results have high statistical confidence (p<.01; Fcrit<F for Alfa = .01). The study recorded a decrease in the potentiation effect of the foliar fertilizers and at the same time in their contribution to the winter wheat yield, when the doses of complex fertilizers applied on the soil were increased.

Keywords: fertilizers interaction, growth model, optimization, winter wheat, yield

1. INTRODUCTION

Wheat is one of the three main cereal crops that cover large areas of land all over the world, with annual yields of over 600 million tons [1]. The major wheat production worldwide is highly concentrated in a few countries, and the European Union is by far the most important producer, accounting for 21% of total world harvest. In the next decades, an increase is expected in the global food demand, and especially in the wheat demand. At European level, 55% of wheat production is concentrated in three countries: France (41 Mio. tons), Germany (24 Mio. tons) and the UK (15 Mio. tons), Romania also having a significant production of wheat (almost 5 Mio. Tons) [2]. The largest agricultural area in Romania country, 747,000 ha, is found in Timis county, bounding area of research, out of which 531 373 ha is arable land. In this region, there are different systems of agriculture practiced, with different levels of productivity, winter wheat and corn crops occupying the largest share. Together with the genetic performance of the biological material [3,4], perfecting the crop

technologies is the main means of ensuring high wheat yields [5,6]. The tendency towards climatic changes, characterized by increased average values of annual temperatures and variations in the distribution of rainfall, brings new challenges regarding the safety of the wheat yield [7,8]. These conditions make it necessary to improve wheat crop technologies, where fertilization is one of the primary elements [9,10].

The basic nutrition of plants takes place through the root system, based on the nutritive elements in the soil [11] supplementing the intake of nutrients through the leaves contributes to getting the nutriment balance right in what the wheat crop is concerned, in view of high and stable yields. The overall cost of the crop technology for the cultivation of wheat has determined the orientation towards cheaper fertilization systems, frequently based on unilateral nitrogen fertilization or foliar fertilization. Fertilization system based on complex fertilizers applied on soil and foliar fertilizers in vegetation is also commonly practiced in wheat crop for economic reasons. But at the same time, they has had serious implications on crop productivity and on soil fertility [12,13]. Many studies have focused on the influence of foliar fertilization on the vegetation status and on wheat yield quality and quantity, in various soil and climatic conditions [14-16].

In the context of these general and specific aspects, the present research aims at assessing the differentiated contribution of mineral fertilization and foliar fertilization to the wheat yield, for improving the fertilization system for winter wheat.

2. MATERIAL AND METHODS

The main objective of the present research was to assess and compare the part that mineral fertilization and foliar fertilization play in the winter wheat yield.

2.1.Experimental field area. The reference area for the study is in the West Plain of Romania, more exactly in Timiș County, Figure 1. Field experiments were carried out at Didactic Station of BUASVM Timișoara, plot A 363, 45° 28' N, 21° 7' E, during 2011-2012 growing seasons.



Figure 1. Timiș County area, Romania – the reference area for study and research (pe-harta.ro)

2.2.Study area and soil condition. The area of the present experiment is included in the great Pannonian Depression, the eastern part to be more exact; this was formed in Pleistocene-Quaternary by gradual clogging of the lake. The natural setting of the

experiment presents slightly gleized cambic chernosem with neutral reaction ($\text{pH} = 6.97\text{--}7.10$), good humus supply ($\text{H} = 3.2\%$), nitrogen index $\text{IN} = 2.97$, high degree of base saturation (over 87%), poor supply of phosphorus ($\text{P} = 23.6 \text{ ppm}$) and medium supply of potassium ($\text{K} = 132 \text{ ppm}$).

2.3.Climatic condition. The climate conditions in the trial area are typical to the continental climate with Mediterranean influences. During the trial period, mean daily temperatures were higher compared to the multi-annual means particularly between May and September (Figure 2). As for the precipitations, there were two periods of rainfall deficit compared to the multi-annual mean: the first interval of rainfall deficit was in May-June and the first decade of July and the second interval of rainfall deficit was in the third decade of July and in August-September (Figure 3).




Figure 2. Mean temperature during the trial period; T M Ma – multiannual mean temperature; T 2011 – the temperature in 2011; T 2012 – the temperature in 2012.




Figure 3. Distribution of rainfalls during the trial period; PM Ma – multiannual mean rainfall; P 2011 – rainfall in 2011; P2012 – rainfall in 2012.

2.4.Biological material. The biological material was represented by Alex wheat cultivar, species *Triticum aestivum* (L.) ssp vulgare.

2.5.Treatments and experimental design. Complex fertilizers of the type NPK (1:1:1), containing 15% active substance for each element, were used in three fertilization levels: NPK_0 (0 kg active substance ha^{-1}), NPK_{30} (30 kg active substance ha^{-1} for each element) and NPK_{60} (60 kg active substance ha^{-1} for each element). In addition to the three basic levels, foliar fertilizers were also applied (300 l water ha^{-1}): Ft – Fertitel 0.5%, containing N (6.7%), P_2O_5 (6%), K_2O (4.2%), Cu (0.025%), Fe (0.1%), Mn (0.01%), Zn (0.008%), B (0.02%) and Mo (0.001%); NZn - Nutrion Zn 0.5%, containing N (8.5%), P_2O_5 (8.5%), K_2O (4.61%), MgO (0.34%), Zn, Fe, Mn, Cu, B, Mo and essential amino acids; Bc - Biocomplex 900 0.5%, brown seaweed liquid extract; Mf – Megafol 0.5%, containing aminoacids (35%), N_i (5.6%), C_{org} 18.7%), K_2O (3.8%); Nt – Nutrilene 0.5% containing N (12%), P_2O_5 (7%), K_2O (8%), B

(0.1%), Fe (0.2%) and Zn (0.05%) and thus there were 6 variants for each NPK base (five foliar fertilized variants, and each control variant). Foliar fertilizers were applied in spring, in two treatments with an interval of 20 days (20 - 30 and 35 - 37 BBCH stage code). The combination of mineral soil fertilization and foliar fertilization resulted in 18 randomized experimental variants, arranged in three repetitions. The area of each plot was 30 m² (10 m length, 3 m width). The crop of each plot unit was harvested at maturity stage by a combine harvester (Hege, Germany).

2.6. Statistical analysis. The experimental data of wheat production, on each variant, were processed through analysis of variance (ANOVA) using the mathematical module on EXCEL 2007. Correlations coefficient, regressions and multivariate analysis were determined using the software PAST [17]. The symbol *, ** and *** used in the paper represent statistically significance at 99.9% (LSD_{0.01%}), 99% (LSD_{0.1%}) and 95% (LSD_{0.5%}) probability level.

3. RESULTS AND DISCUSSION

Under the experimental conditions presented above, the fertilizers applied in each fertilization variant were differently used up by the winter wheat crop, resulting in different levels of yields. The yield varied from 2925±72.17 kg ha⁻¹ in variant Mt₀ and 5145±106.81 kg ha⁻¹ in the variant fertilized with Megafol on a NPK₆₀ base, as shown in Table 1. The overall yield increase ranged from 355 kg ha⁻¹ in the variant fertilized with Biocomplex 900 with NPK₀ fertilization and 2220 kg ha⁻¹ in the variant fertilized with Megafol, with NPK₆₀.

ANOVA test reveals statistical assurance for the experimental results, both for complex mineral fertilizers (rows sources of variance; $p < .01$; $F > F_{crit}$) and for foliar fertilizers (column sources of variance; $p < .01$; $F > F_{crit}$), Table 2.

The comparative analysis of the results obtained in the experimental variants reveals the different contribution of the two types of fertilizers in the formation of the yield. The complex fertilizers – as a singular effect – generated yield increases between 925*** and 1425*** kg ha⁻¹, the comparative analysis being made between variants Mt₁ (NPK₃₀) and Mt₂ (NPK₆₀) respectively, with the control Mt₀ (NPK₀).

In the cases in which foliar fertilizers were applied together with soil fertilizers, the effect of the complex fertilizers was potentiated by 0.9 – 1.24 production units with NPK₃₀ and by 0.78 – 1.14 production units with NPK₆₀. The study revealed that the potentiation effect decreased if the doses of complex fertilizers applied on the soil were increased, because in that case plant roots were supplied with higher nutriment content. Generally, plants prefer to feed through the root system if the soil, which is their main medium for nutrition, ensures enough nutrients. The large share of root nutrient absorption as compared to the foliar nutrient absorption has been revealed by other studies as well [18,19].

With foliar fertilization on the background of NPK₀, the yield obtained ranged from 3280±75.06 kg ha⁻¹ in the variant fertilized with Biocomplex 900, to 3735±152.98 kg ha⁻¹ in the variant fertilized with Nutrilene. The control variant (Mt₀), given by the natural fertility of the soil, resulted in a yield of 2925±72.17 kg ha⁻¹. The yield increase generated by foliar fertilization combined with the natural fertility of the soil (NPK₀) was between 355 kg ha⁻¹ in the variant fertilized with Biocomplex 900 and 810*** kg ha⁻¹ in the variant fertilized with Nutrilene. In the other variants with foliar fertilization, the yield increase was 550** kg ha⁻¹ in the variant fertilized with Nutrien Zn, 675** kg ha⁻¹ in the variant fertilized with Fertitel and 700*** kg ha⁻¹ in the variant fertilized with Megafol.

The particular influence of foliar fertilization has been assessed in other studies as well, with differentiated contribution on yield quantity and quality, depending on the type of fertilizer, the type and association of nutriments, climate and soil conditions, biological material and the time of fertilization [20,21].

Complex fertilizers administered in quantities of 30 kg a.s. ha⁻¹ ensured the second medium for nutrition (NPK₃₀) for the winter wheat crop where foliar fertilizers were applied. Under

these conditions, the yield level ranged from $3850 \pm 86.60 \text{ kg ha}^{-1}$ in the control variant Mt_1 , represented by the conditions created by mineral fertilization (NPK_{30}) and $4750 \pm 144.33 \text{ kg ha}^{-1}$ in the variant fertilized with Fertitel, Table 1.

Table 1. The yield of winter wheat, Alex cultivar, obtained under the influence of foliar fertilization, combined with different NPK doses of soil fertilizers (average values over the experimental period 2011 – 2012)

Fertilization variant	Variant number	Yield (mean values 2011-2012)	General yield increase as compared to the control variant	Yield increase generated by foliar fertilizers	Yield increase generated by complex fertilizers
NPK_0	Mt_0	V_1 2925 $\pm$ 72.17	-	-	-
	Fertitel	V_2 3600 $\pm$ 144.33	675**	675**	-
	Nutrion Zn	V_3 3475 $\pm$ 129.90	550**	550**	-
	Biocomplex 900	V_4 3280 $\pm$ 75.06	355	355	-
	Megafof	V_5 3625 $\pm$ 187.64	700***	700***	-
	Nutrilene	V_6 3735 $\pm$ 152.98	810***	810***	-
The yield increase generated by foliar fertilizers was calculated by comparison with the control variant, Mt_0					
NPK_{30}	Mt_1	V_7 3850 $\pm$ 86.60	925***	-	-
	Fertitel	V_8 4750 $\pm$ 144.33	1825***	900***	1150***
	Nutrion Zn	V_9 4590 $\pm$ 132.79	1665***	740***	1115***
	Biocomplex 900	V_{10} 4400 $\pm$ 173.21	1475***	550**	1120***
	Megafof	V_{11} 4740 $\pm$ 127.02	1815***	890***	1115***
	Nutrilene	V_{12} 4595 $\pm$ 101.04	1670***	745***	860***
The yield increase generated by foliar fertilizers was calculated by comparison with the Mt_1					
The yield increase generated by complex fertilizers was calculated by comparison with the similar variants of foliar fertilizers with NPK_0 (Mt_0)					
NPK_{60}	Mt_2	V_{13} 4350 $\pm$ 109.69	1425***	-	-
	Fertitel	V_{14} 4905 $\pm$ 152.99	1980***	555**	1305***
	Nutrion Zn	V_{15} 5085 $\pm$ 147.22	2160***	735***	1610***
	Biocomplex 900	V_{16} 4900 $\pm$ 115.47	1975***	550**	1620***
	Megafof	V_{17} 5145 $\pm$ 106.81	2220***	795***	1520***
	Nutrilene	V_{18} 4850 $\pm$ 150.11	1925***	500*	1115***
The yield increase generated by foliar fertilizers was calculated by comparison with the Mt_2 variant					
The yield increase generated by complex fertilizers was calculated by comparison with the similar variants of foliar fertilizers with NPK_0 (Mt_0)					
Limits the significance of differences $LSD_{5\%} = 382.65$; $LSD_{1\%} = 512.71$; $LSD_{0.1\%} = 676.71$					

159

160

Table 2. ANOVA: Two-Factor without Replication

Source of Variation	SS	Df	MS	F	P-value	F crit
Rows (mineral fertilizer)	20229842	8	2528730	55.76621	2.81E-19	4.207037
Columns (foliar fertilizer)	3832050	5	766410	16.90168	5.84E-09	5.128263
Error	1813808	40	45345.21			
Total	25875700	53				

161 Alfa = .01

162

163 The general yield increase, as compared to the control variant Mt_0 , was from 925*** kg ha^{-1}
 164 in variant Mt_1 (NPK_{30} fertilization level) and 1825*** kg ha^{-1} in the variant fertilized with

Fertitel on the background of NPK₃₀. Complex fertilizers ensured yield increase between 860*** and 1150*** kg ha⁻¹ in comparison with similar variants with NPK₀. The highest yield, 4750±144.33 kg ha⁻¹, was obtained by the associating complex soil fertilization with Fertitel. The foliar fertilizers potentiated differently the effect of the complex fertilizers, with values of 0.93 units (Nutrilene), 1.21 units (Biocomplex 900), 1.20 units (Nutrion Zn, Megafol) and 1.24 units, respectively (Fertitel). The yield increase given by foliar fertilizers on NPK₃₀ ranged from 550** kg ha⁻¹ in the case of Biocomplex900 to 900*** kg ha⁻¹ in the case of Fertitel, with statistical assurance.

Base fertilization with complex fertilizers at the level of 60 kg active substance ha⁻¹ (NPK₆₀) constituted the third type of base fertilization that was combined with foliar fertilizers. The yield level was between 4350±109.69 kg ha⁻¹ in the variant NPK₆₀ (Mt₂) and 5145±106.81 kg ha⁻¹ in the variant where Megafol was applied, Table 1. The general yield increase, compared to Mt₀, ranged from 1425*** kg ha⁻¹ in the case of variant Mt₂ (NPK₆₀) and 2220*** kg ha⁻¹ in the case of the variant fertilized with Megafol on the background of NPK₆₀, where the yield differences are statistically assured.

Complex fertilizers determined yield increases ranging from 1115*** kg ha⁻¹ and 1620*** kg ha⁻¹, as compared to Mt₀, with statistically assured differences. The highest yield was obtained when complex mineral fertilization was combined with the foliar fertilizer Megafol (5145±106.81 kg ha⁻¹). The effect of complex fertilizers applied on the soil was potentiated in a different way by foliar fertilizers on the background of this NPK base fertilization, but generally the values were lower than in the previous NPK dosage (NPK₃₀); 0.78 (Nutrilene), 0.92 (Fertitel), 1.07 (Megafol), 1.13 (Nutrion Zn) and 1.14 (Biocomplex 900). Foliar fertilization on this NPK dosage (NPK₆₀) determined yield increases ranging from 500* kg/ha in the variant fertilized with Nutrilene and 795*** kg/ha in the variant fertilized with Megafol, the yield differences being statistically assured. Although in most cases there were differences from the control variant, the yield increase generated by foliar fertilization was relatively small, of about 300 to 800 kg ha⁻¹, in the conditions of natural soil fertility (NPK₀).

Under the soil and climate conditions of the present research, application of only foliar fertilization on the wheat crop determined relatively low yields and small increases as compared to the yield given by the natural fertility of the soil. In some cases, the yield differences were not statistically assured, which means that unilateral fertilization with foliar fertilizers presents a certain degree of uncertainty. Similar results were communicated also by [22,23], according to which, foliar fertilizers had the main influence of the quality of the wheat crop. The nutrient content in foliar fertilizers and the dilutions in which they are applied, as well as other bioactive products applied on the leaves, cause them to have an influence on the metabolic and enzymatic processes in plants that stimulate photosynthesis [24], synthesis and accumulation of carbohydrates [25,26], the increase of the protein content in beans [27,28]. When associated with complex soil fertilization (NPK₃₀, NPK₆₀), foliar fertilizers had a significantly higher contribution to the wheat yield, with variations from one type of fertilizer to another. Nevertheless, soil nutrients had the major contribution to the yield; foliar fertilizers only potentiated their effect and contributed to significant yield increases with statistical assurance.


Regression analysis of the potentiation capacity of foliar fertilization on the effect of complex fertilizers generated models that describe, with high statistical certainty, the contribution of foliar fertilization on yield increase after mineral fertilization, $R^2 = 0.844$, equation (1) with NPK₃₀ and $R^2 = 0.989$, equation (2) with NPK₆₀, Figure 4.

$$PF_{NPK_{30}} = -0.0347x^2 + 0.2153x + 0.9232 \quad (1)$$

$$PF_{NPK_{60}} = -0.0767x^2 + 0.5806x + 0.0646 \quad (2)$$

where: $PF_{NPK_{30}}$ - potentiation factor on NPK₃₀;

216 $PF_{NPK_{60}}$ - potentiation factor on NPK_{60} ;
 217 x – foliar fertilizer.



218
 219
 220 **Figure 4. The particular distribution of the potentiation values given by foliar fertilizers**
 221 **in relation to the experimental results obtained (Mt – control variant, Ft – Fertitel, NZn**
 222 **– Nutrion Zn, Bc – Biocomplex900, Mf – Megafol, Nt – Nutrilene)**
 223

224 Multivariate analysis of the experimental results facilitated the orientation and grouping of
 225 variants based on similarity. The experimental variants were grouped in two major clusters,
 226 with two large subgroups each. The first cluster contains mainly the variants with foliar
 227 fertilization on the background of NPK_0 ($V_1 - V_6$) and variant V_7 , which gave yields in the
 228 same value group, Figure 5. The second major cluster comprises the variants with foliar
 229 fertilization, on the background of NPK_{30} ($V_8 - V_{13}$) and NPK_{60} ($V_{14} - V_{18}$), which in their turn
 230 have two subclusters according to their values. The cophenetic coefficient had the value of
 231 0.859, which gives high degree of confidence to the grouping of the experimental results
 232 based on Euclidean distances.




Figure 5. Variants grouping based on similarity, according to the Euclidean distances.

4. CONCLUSION

Soil mineral fertilization and foliar fertilization had different contributions to the **winter** wheat yield, the contribution of soil fertilization being significantly higher. Foliar fertilization potentiated the effect of mineral fertilization applied on the soil, with values ranging from 0.93 to 1.24 yield units where mineral fertilization was NPK₃₀ and from 0.78 to 1.14 yield units where soil fertilization was NPK₆₀. The study revealed a reduction in the effect of foliar fertilizers on the **winter** wheat yield as the doses of soil fertilizers were increased. The prediction model for the potentiation factor (PF) is described by polynomial equations of the second degree that allow high certainty prediction of the contribution of foliar fertilizers to the yield increase. Multivariate analysis based on Euclidean distance facilitated the orientation and grouping of the variants depending on similarities in result generation, the value of the cophenetic coefficient 0.963 providing statistical assurance to the results. From the distribution of the data, convenient variants can be chosen based on similarity in result generation.

REFERENCES

1. Shewry PR. Wheat, Darwin review. J. Exp. Bot. 2009;60(6):1537-1553, doi:10.1093/jxb/erp058
2. Vigani M, Dillen K, Cerezo ER. Proceedings of a workshop on "Wheat productivity in the EU: determinants and challenges for food security and for climate change". European Commission. 2013. JRC 80645, EUR 25934, doi:10.2791/93338.
3. Fufa H, Baenziger PS, Beecher BS, Graybosch RA, Eskridge KM, Nelson LA. Genetic improvement trends in agronomic performances and end-use quality characteristics among hard red winter wheat cultivars in Nebraska. Euphytica. 2005;144:187-198, doi 10.1007/s10681-005-5811-x
4. Xiong YC, Li FM, Zhang T. Performance of wheat crops with different chromosome ploidy: root-sourced signals, drought tolerance, and yield performance. Planta.

- 2006;224(3):710-718, doi:10.1007/s00425-006-0252-x
5. Snapp SS, Swinton SM, Labarta R, Mutch D, Black JR, Leep R et al. Evaluating cover crops for benefits, costs and performance within cropping system niches. *Agron J*. 2005;97(1):322-332, doi:10.2134/agronj2005.0322
6. Tester M, Langridge P. Breeding technologies to increase crop production in a changing world. *Science*. 2010;327(5967):818-822, doi: 10.1126/science.1183700
7. Ortiz R, Sayre KD, Govaerts B, Gupta R, Subbarao GV, Ban T et al. Climate change: Can wheat beat the heat?. *Agr Ecosyst Environ*. 2008;126(1-2):46-58, <http://dx.doi.org/10.1016/j.agee.2008.01.019>
8. Olensen JE, Trnka M, Kersebaum KC, Skjelvåg AO, Seguin B, Peltonen-Sainio P et al. Impacts and adaptation of European crop production systems to climate change. *Eur J Agron*. 2011;34(2):96-112, <http://dx.doi.org/10.1016/j.eja.2010.11.003>
9. Grzebisz W, Gaj R, Sassenrath GF, Halloran JM. Fertilizer use and wheat yield in Central and Eastern European Countries from 1986 to 2005 and its implication for developing sustainable fertilizer management practices. *Commun Soil Sci Plan*. 2012;43(18):2358-2375, doi:10.1080/00103624.2012.708080
10. Grassini P, Eskridge KM, Cassman KG. Distinguishing between yield advances and yield plateaus in historical crop production trends. *Nat Commun*. 2013;4:1-11, doi:10.1038/ncomms3918
11. Mengel K, Kirkby EA. Principles of Plant Nutrition. Kluwer Academic Publisher, Dordrecht/Boston/London; 2001.
12. Hera C. CIEC – 77 years dedicated to enhancing soil fertility for the benefit of sustainable and performing agriculture. in 15th World Fertilizer Congress of the International Scientific Centre for Fertilizers (CIEC) Proceedings, 2010;17-29, Ed. Academiei Române, București.
13. Pîslea D, Sala F. The influence of mineral fertilization on the distribution of nutrients in the soil. *Bulletin USAMV Agriculture*. 2012;69(1):225-231.
14. Ahmed W, Yaseen M, Arshad M, Ali Q. Response of wheat (*Triticum aestivum* L.) to foliar feeding of micronutrients. *International Journal for Agro Veterinary and Medical Sciences*. 2011;5(2):209-220, doi: 10.5455/ijavms.20110601090456
15. Rawashdeh HM, Sala F. The effect of foliar application of iron and boron on early growth parameters of wheat (*Triticum aestivum* L.). *Research Journal of Agricultural Sciences*. 2013; 45(1):21-26.
16. Rezaeei M, Daneshvar M, Shirani AH. Effect of iron nano chelated fertilizers foliar application on three wheat cultivars in Khorramabad climatic conditions. *Scientific Journal of Crop Science*. 2014;3(2):9-16.
17. Hammer Ø, Harper DAT, Ryan PD. PAST: Paleontological statistics software package for education and data analysis. *Palaeontol Electron*. 2001;4(1):9pp. http://palaeo-electronica.org/2001_1/past/issue1_01.htm
18. Marschner H. Mineral Nutrition of Higher Plants. Second Edition, Academic Press, Elsevier; 1995.
19. Epstein E, Bloom JA. Mineral Nutrition of Plant: Principles and Perspectives, Second edition, Sinauer Associates, Inc. Publishers Sunderland, Massachusetts; 2005.
20. Woolfolk CW, Raun WR, Johnson GV, Thomason WE, Mullen RW, Wynn KJ et al. Influence of late-season foliar nitrogen applications on yield and grain nitrogen in winter wheat. *Agron J*. 2002;94(3):429–434, doi:10.2134/agronj2002.4290
21. Nadim MA, Awan IU, Baloch MS, Khan EA, Naveed K, Khan MA. Response of wheat (*Triticum aestivum* L.) to different micronutrients and their application methods. *J Anim Plant Sci*. 2012;22(1):113-119.
22. Luo C, Branlard G, Griffen WB, McNeil DL. The effect of nitrogen and sulfur fertilization and their interaction with genotype on wheat glutenins and quality parameters. *J Cereal Sci*. 2000;31(2):185- 194. doi:10.1006/jcrs.1999.0298
23. Saeed B, Gul H, Ali F, Khan AZ, Anwar S, Nasrullah et al. Contribution of soil and foliar

- 318 fertilization of nitrogen and sulfur on physiological and quality assessment of wheat
319 (*Triticum aestivum* L.). Natural Science. 2013;5(9):1012-1018,
320 doi:10.4236/ns.2013.59125
- 321 24. Dromantienė R, Pranckietienė I, Šidlauskas G, Pranckietis V. Changes in technological
322 properties of common wheat (*Triticum aestivum* L.) grain as influenced by amino acid
323 fertilizers. Zemdirbyste-Agriculture, 2013;100(1):57–62, UDK 633.11:631.8:631.55 / doi
324 10.13080/z-a.2013.100.008
- 325 25. Ebrahim MKH, Aly MM. Physiological response of wheat to foliar application of zinc and
326 inoculation with some bacterial fertilizers. J. Plant Nutr. 2005;27(10):1859-1874,
327 doi:10.1081/PLN-200026442
- 328 26. Sadak SM, Elhamid EMA, Mostafa HM. Alleviation of adverse effects of salt stress in
329 wheat cultivars by foliar treatment with antioxidants I. changes in growth, some
330 biochemical aspects and yield quantity and quality. Am Eurasian J Agric Environ Sci.
331 2013;13(11):1476-1487, doi: 10.5829/idosi.ajeaes.2013.13.11.11264
- 332 27. Abad A, Lloveras J, Michelena A. Nitrogen fertilization and foliar urea effects on durum
333 wheat yield and quality and on residual soil nitrate in irrigated Mediterranean conditions.
334 Field Crop Res. 2004;87(2-3):257-269, <http://dx.doi.org/10.1016/j.fcr.2003.11.007>
- 335 28. Zeidan MS, Mohamed MF, Hamouda HA. Effect of foliar fertilization of Fe, Mn and Zn on
336 wheat yield and quality in low sandy soils fertility. WJAS. 2010;6(6):696-699.