

COMPARATIVE EVALUATION OF ORGANIC AND INORGANIC MANURE ON SWEET PEPPER PERFORMANCE IN TWO ECOLOGICAL ZONES OF NIGERIA

ABSTRACT

Organic (poultry manure) and inorganic (Urea) fertilizers were comparatively studied with the aim of improving the performance of sweet pepper (*Capsicum annum*) in two ecological zones of Nigeria (Ado Ekiti, Ekiti state and Omu-Aran, Kwara State). Poultry manure applied at the rate of 8 tha^{-1} gave higher fruit yield than urea at the rate of 200 kg ha^{-1} twice equally at two weeks after transplanting (WAT) and at fruiting. Higher leaf number and branches were produced in the plots that received urea without a corresponding superior yield. It is indicative from this study that pepper can be successfully produced organically using poultry manure in the two ecological zones under study. It also suggests that our soils can be protected from the level of acidity associated with synthetic fertilizer use. Pepper fruits produced from such will be purely organic and safer for human consumption.

Key words: sweet pepper, organic manure, inorganic manure, yield

1. INTRODUCTION

The provision of quality food starts with planting, production and preparation of food. Organic farming is one of the fastest growing sectors of agriculture worldwide. Its main objective is to create a balance between the interconnected systems of soil organisms, plants, animals and humans [1]. Plants need nutrients in specific proportions to be present in the soil [2] and fertilizer use is a key factor in soil fertility management and yield increase in crop production [3].

Pepper is one of the most important vegetable crops grown in Nigeria [4], where it is commonly used as condiments [5]. Pepper had even been reported to prevent and treat certain ailments like cold and fever, and also improve other food intake [6, 7]. Among the various reported benefits resulting from the use of organic fertilizer in crop production includes; enhancement of soil productivity, increase in the soil organic carbon content, soil micro-organism activities, improvement of soil structure, increase in balance soil nutrient status of the soil and enhancement of crop yield [6,8,9].

Although the applications of organic and inorganic fertilizers have been reported to improve the growth and development of pepper [10], but better performance of pepper and other plants under poultry manure had been reported than under the inorganic fertilizer application [11, 12, 4].

The trend in the world today is to produce food in the organic way for quality food that will promote good health of people. The present study was, therefore, carried out to evaluate the comparative effect of poultry manure and inorganic fertilizer on the growth and yield of sweet pepper in two ecological zones of Nigeria.

2. MATERIALS AND METHODS

The experiments were carried out at the experimental site of the of the Department of Plant Science, Ekiti State University, Ado Ekiti, Ekiti state, Nigeria (7°40' N, 5°15' E) and in Landmark University

Comment [Dr1]: Add Activities

Comment [Dr2]: Add balance

Comment [Dr3]: Add and development

Teaching and Research Farm, Omu-Aran, Kwara State, Nigeria (8°25' N, 4°40' E). Both locations have a bimodal rainfall pattern with an annual mean of 1,400mm and 1232mm respectively. The land was cleared manually with cutlass and ridges made with Nigerian hoe. The soil at the experimental site in Ado Ekiti was analysed to be a sandy clay loam with an organic matter content of 4.28; 0.48%N; 5.95mg/kg P; 0.08Cmol K; and a pH of 6.65. Each ridge was 4m long with 1m between ridges. Four ridges constituted a plot and the experiment was laid out in a randomized complete block. Each treatment was replicated four times. The treatments were: Urea; poultry manure; and control. Sweet pepper (*Capsicum annum*.) seedlings were transplanted at four weeks after sowing [13] at the spacing of 0.6 m x1 m. The poultry manure was applied on the ridge three weeks before transplanting at the rate of 8 tha^{-1} [14]. Urea was applied at the rate of at the rate of 200kg/ha-1 twice equally at two weeks after transplanting (WAT) and at fruiting. Weeds were manually removed by hand pulling.

The growth and yield parameters determined were yield and yield attributing factors which include: The growth and yield parameters determined include: plant height; stem girth; number of green branches at 8 WAT; number of leaves at 8 WAT; number of fruits per plant; fruit girth; and fruit weight (tha^{-1}). Collected data were statistically analysed and means compared using the fisher's least significant difference (LSD) [1].

Comment [Dr4]: Unclear; please rephrase

Formatted: Font color: Orange

Comment [Dr5]: Add Reference??

3. RESULTS

Effects of organic and inorganic manure on pepper height and stem girth are presented in Table # 1. Pepper heights observed in poultry manure and Urea were not significantly different but higher than the one obtained in the control experiment. Stem girth obtained in the poultry manure applied plots were highest in both years except in 2014 when it was identical with those of urea applied plots. The control plots gave the lowest stem girth.

Table # 2 shows the effects of organic and inorganic manure fertilizer on number of pepper branches and leaves. The highest number of branches and leaves per plant were obtained in the urea applied plots but not significantly different from the poultry manure applied plots. The control experiment consistently resulted to the lowest number of branches and number of leaves per plant.

Comment [Dr6]: Manure or fertilizer???
Please use one word in the manuscript, sometime, you are using the manure and some time, you are using the fertilizer..... I think that the Inorganic fertilizer or synthetic fertilizer are good words

Comment [Dr7]: Please use the word maximum rather than highest

Comment [Dr8]: Replace highest with maximum

Comment [Dr9]: Correct these one also!!

Comment [Dr10]: Maximum

The highest number of fruits per plant was obtained in the poultry manure applied plots in both trials while the least was observed in the control plots. Urea produced higher number of fruits than the control but lower than the poultry manure plots (Table # 3). Fruit girth obtained from poultry manure plots and those of urea were identical but higher than the ones obtained from the control plots.

Table # 4 shows the effects of organic and inorganic manure organic and inorganic fertilizers on pepper yield. The highest fruit yield of pepper in both years was recorded in the plots that received poultry manure while the lowest was observed in the control plots. Poultry manure was significantly higher than urea in terms of fruit yield production.

4. DISCUSSION

The results obtained from this study clearly showed that the organic manure (poultry manure) used in the two locations of the experiment substantially increased the performance of pepper in terms of growth and yield.

The highest growth parameters and yield obtained from the plots that received poultry manure application probably reflected high release of essential nutrients especially nitrogen and phosphorous from this organic fertilizer (poultry manure). Reports from other researchers have shown that organic-amended soils have twice the level of N as conventional soils [15,16,17, 16, 17]. Poultry manure had been reported to possess essential nutrient elements which are associated with high photosynthetic activity capable of promoting roots and vegetative growth of plants [18, 4]. Higher but insignificant

Comment [Dr11]: Replace highest with maximum

Formatted: Font color: Red

number of leaves and branches obtained from Urea applied plots than poultry manure in this work may suggest over supply of nitrogen which increased vegetative growth without a corresponding increase in yield. Similar vegetative growth in pepper had been reported when NPK fertilizers increased the number of leaves but lower yield than Tithonia diversifolia leaf biomass [19]. Similar report had shown that hairy vetch/rye mineralization allowed for a slower release of N and more efficient absorption by sweet corn [20].

5. CONCLUSION

The present work had clearly shown that pepper can be successfully produced organically using poultry manure in the two ecological zones under study. Since, poultry manure is readily available as wastes from poultry farmers, it will be useful for farmers at very low cost and thus, reduce the high cost associated with inorganic fertilizers. Our soils will also be protected from the level of acidity associated with synthetic fertilizers use. The pepper fruits produced from such will be purely organic and safer for human consumption.

Table 1: Effects of organic and inorganic fertilizers manure on pepper height and stem girth

Comment [Dr12]: Please correct the word Inorganic manure??

Treatments	Plant height (cm)		Stem girth (mm)	
	2012	2014	2012	2014
Control	34.6b	33.4b	10.2c	12.0b
Poultry manure	45.3a	42.9a	18.3a	17.1a
Urea	43.4a	45.2a	16.1b	15.4a
LSD	5.55	6.81	1.64	2.28

Means with the same alphabet(s) within columns are not significantly different (P=.05)

Table 2: Effects of organic and inorganic fertilizers manure on number of branches and number of leaves of pepper

Treatments	Number of branches plant ⁻¹		Number of leaves plant ⁻¹	
	2012	2014	2012	2014
Control	9b	5b	109.1b	96.5b
Poultry manure	23a	21a	135a	129.5a
Urea	24a	23a	137a	129.0a
LSD	6.45	3.71	5.73	2.64

Means with the same alphabet(s) within columns are not significantly different (P=.05)

Table 3: Effects of organic and inorganic fertilizers manure on number of fruits and girth of fruits of pepper

Treatments	Number of fruits plant ⁻¹		Fruit girth (mm)	
	2012	2014	2012	2014
Control	5c	7c	34.5b	31.7a
Poultry manure	21a	17a	40.1a	39.9a
Urea	15b	13b	40.2a	36.5a
LSD	3.51	2.76	4.63	NS

Means with the same alphabet(s) within columns are not significantly different (P=.05)

113 | Table 4: Effects of organic and inorganic fertilizers manure on pepper yield (g per plant)

Treatments	Pepper Yield (t ha ⁻¹)	
	2012	2014
Control	7.40c	6.02c
Poultry manure	22.71a	20.09a
Urea	15.12b	13.17b
LSD	3.96	3.64

114 Means with the same alphabet(s) within columns are not significantly different (P=.05)

115 REFERENCE

- 116 1. Karanatsidis G, Berova M. Effect of Organic-N Fertilizer on Growth and some Physiological
117 Parameters in Pepper Plants (*Capsicum annum* L.) Biotechnol. & Biotechnol.EQ. 2009; 254-
118 257
- 119 2. Fagbayide. Growth and nutrient concentration of pepper fruits as affected by phosphorus
120 fertilizer. Proceeding, 15th annual conference, Horticultural society of Nigeria. 1997; 40-42.
- 121 3. Akinrinde EA. Issues of optimum nutrient supply for sustainable crop production in
122 Tropical Developing Countries. Pakistan Journal of Nutrition 2006; 5 (4), 377-387. Asian
123 Network for Scientific Information.
- 124 4. Aliyu L. The effect of organic and mineral fertilizer on growth yield and composition of pepper
125 (*Capsicum annum* L.) Biol Agric Hort. 2000; 18: 29-36.
- 126 5. Alabi DA. Effect of fertilizer phosphorus and poultry droppings treatments on growth and
127 nutrient components of pepper (*Capsicum annum* L.). African Journal of Biotechnology. 2006;
128 5(8): 671-677.
- 129 6. Udoh DJ, Ndon BA, Asuquo PE, and Ndaeyo NU. Crop Production Techniques for the
130 Tropics. Concept Publication ,Lagos, Nigeria. 2005; p. 446.
- 131 7. Grubben GJ. Tropical Vegetables and their Genetic Resources. IPRGR, Rome, Italy. 1977; p.
132 197.
- 133 8. FAO. Fertilizers and their uses. 2000; pp. 26-39.
- 134 9. Ademiluyi BO, Omotoso SO. Comparative evaluation of *Tithonia diversifolia* and NPK fertilizer
135 for soil improvement in maize (*Zea mays*) production in Ado Ekiti, South-western Nigeria.
136 *Research Journal of Agronomy* 2008; 2 (1): 8-11.
- 137 10. Malik AA, Chattoo MA, Sheemar G, Rashid R. Growth, yield and fruit quality of sweet pepper
138 hybrid SH-SP-5(*Capsicum annuum* L.) as affected by integration of inorganic fertilizers and
139 organic manures (FYM), Journal of Agricultural Technology. 2011; 7 (4):1037-1048.
- 140 11. Hussein TO. Effect of poultry manure on the growth of tomato. Proceeding of 15th Annual
141 Conference of Horticultural Society of Nigeria (HORTSON), April, 8-11 soil analysis Manual
142 series Nigeria. 1997; pp. 43-45.
- 143 12. De - Lannoy G, Romain HR. Crop Production in Tropical Africa. London, Mc. Graw Hill
144 Publishers Ltd. New York. 2001; pp. 234-245.
- 145 13. Ademiluyi BO. Studies on Transplanting Age and Pepper (*Capsicum annum* L) Performance
146 in a South-western Nigeria Location. *Bulletin of pure and Applied Sciences*. 2012; 31B (2):
147 53-57.
- 148 14. Ikeh AO, Ndaeyo NU, Uduak IG, Iwo GA, Ugbe LA, Udoh EI, Effiong GS. Growth and yield
149 Responses of Pepper (*Capsicum frutescens* L.) to Varied Poultry Manure Rates in Uyo,
150 Southeastern Nigeria. ARPN Journal of Agricultural and Biological Science. 2012; 7, (8): 735-
151 742
- 152 15. Burger M, Jackson LE. Microbial immobilization of ammonium and nitrate in relation to
153 ammonification and nitrification rates in organic and conventional cropping systems. *Soil Biol.*
154 *Biochem*. 2003; 35:29-36.
- 155 16. Delate K, Cambardella C, Mckern A. Effects of Organic Fertilization and Cover Crops on an
156 Organic Pepper System. HortTechnology 2008; 18: 215-226.

- 157 17. Dauda S, Aliyu NL, Chiezey UF. Effect of variety, seedling age and poultry manure on growth
158 and yield of garden egg (*Solanum gilo* L.) Aliju Acad Forum. 2005; 988-995.
159 18. John GC, Almazan LP, Paria J. Effects of Nitrogen fertilizer on the intrinsic rate of the rusty
160 plum aphid. Environmental Entomology. 2004; 34(4): 938-943.
161 19. Ademiluyi BO. Comparative Evaluation of NPK Fertilizer and *Tithonia diversifolia* Biomass in
162 Sweet Pepper (*Capsicum annum*) Production in Ado Ekiti, Nigeria. Journal of Life Sciences,
163 2013; 7(3): 288-291
164 20. Carrera LM, Abdul-Baki AA, Teasdale JR. Cover crop management and weed suppression in
165 no-tillage sweet corn production. *HortScience* 2004; 39:1262–1266.
166

167
168
169
170
171
UNDER PEER REVIEW