

Review Article**Grasscutter (*Thryonomys Swinderianus*) Production in West Africa: Prospects,
Challenges and Role in Disease Transmission****ABSTRACT**

This review brings together and consolidates the various researches that had been undertaken in grass-cutter with the aim of providing adequate information that will be capable of improving and sustaining the production of the animal as well as its consumption in West Africa. Given the above scenario, this paper reviewed the results obtained by different researchers on feeding and nutritional requirement of the grasscutter, housing, performance, anatomy and morphology, some environmental issues, the challenges and the role of grasscutter in disease transmission. It was clear from this study that grasscutter is widely acceptable, utilizes inexpensive feed to produce good meat of high biological value without any trace of pathogenic organisms, survives in simple housing apartments when confined, has good litter size and short generation interval, has simple anatomical dispositions which helps in breeding and they are capable of adapting to environments similar to their natural habitat, thus, shedding off their nocturnal habit to become more active in the daytime. However, some challenges of the grasscutter include irregular supply of breeding stock, environmental issues, poor processing and marketing plan, lack of balanced diet, poor producer training and education, inadequate infrastructural development, poor information dissemination, incidences of diseases and mortality. With good understanding of the prospects and positive utilization of the challenges, there will be great improvement and sustenance in the production of the animal, such that more jobs will be created, more income generated and protein intake in the sub-region will be increased. This overview therefore serves as a means to sustainable grasscutter production and consumption in West Africa as well as an indicator of the direction of further research on grasscutter production.

28 **Key words:** grass-cutter, feeding, housing, performance, diseases transmission, challenges.

29

30

31

32 INTRODUCTION

33 Wildlife has great potentials for meat production and serves as an important source of the
34 highly desired animal protein to the people of Africa, both in urban areas and rural
35 communities (Fonweban and Njwe, 1990). In Nigeria there is an abundant variety of wildlife
36 resources capable of supporting the protein intake of the populace. But in recent times, there
37 had been significant short fall between the production and supply of animal protein to feed
38 this ever increasing population (Akpan, *et al.*, 2009). To arrest this unacceptable trend, efforts
39 had been directed towards boosting the micro livestock sector. Among the micro-livestock
40 animals is the grass-cutter or cane rat (*Thryonomys swinderianus*) popularly called Oya by
41 the Yorubas, Nchi by the Igbos and Gegbi in Hausa Language in Nigeria, Akranti/Akrantie in
42 Ghana and simply grass-cutter in other West African Countries (Figure 1 and 2).

43

44 Grass-cutter is a widely hystricomorphic rodents widely distributed in the African sub-region
45 and exploited in most areas as a source of animal protein (NRC, 1991). It is a heavily built
46 animal with round muzzle, small round ears, short tail and hursh bristly fur (Taiwo, 2006).
47 Apart from being the most preferred (Clottey, 1981; Martin, 1985) , it is the most expensive
48 meat in most countries in West Africa including Nigeria, Togo, Benin, Ghana and Cote d'
49 Voire (Asibey and Addo, 2000). It contributes to both local and foreign earning in some of
50 these countries (Natiama-Baidu, 1998). Most rural population in Nigeria depends on bush
51 meat for their dietary protein supply (FAO 1982). Grass-cutter meat had been reported to
52 have higher nutritional value (Opara, 2010) and meat yield (Omole *et al.*, 2005).
53 Consequently, the demand for livestock products could be solved through the production of
54 grass-cutters meat (Adekola and Ogunsola, 2009). It is considered a delicacy in West and
55 Central Africa (NRC, 1991). It serves as a steady source of animal protein in many rural areas
56 of Nigeria and other West Africa countries like Benin, Ghana, Togo, Cot D' Voire
57 (Ogunsanmi *et al.*, 2002) and South Africa (Van-Zyl *et al.*, 1999). Most Chinese who are
58 resident in Nigeria cherish grass-cutter meat as regular meal and form a delicacy for
59 entertainment for their guest from abroad (The Thy Consulting, 2006). This preference for
60 grass-cutter is attributable to its high carcass quality and protein that is comparable to that of

poultry, especially turkey and other domesticated livestock like rabbit, cattle, sheep, goat and pig (Ajayi and Tewe, 1980, Fayenuwo *et al.*, 2003). A crude protein content of 22.7% had been reported for grass-cutter meat. This value is higher than the crude protein values of 20.7% for rabbit meat, 19% for chicken meat and 18.2% for beef and 22.2% for turkey (Anieunam, 2005). Its mature weight has been reported to be more than 9kg for males and 5-7kg for females (Fayenuwo *et al.*, 2003). It is therefore obvious that with the ever increasing human population and obvious protein shortage in Africa, there is need to explore other means of providing readily and acceptable meat on short term basis. A good understanding of the principles and techniques of production will lead to profitable grass-cutter business (Akinola, 2008). Thus with appropriate information regarding the prospects and possible challenges of grass-cutter production, more farmers will engage in the production of grass-cutter. This will go a long way in alleviating poverty, reducing hunger, create jobs, improve income and improve the immune system of Nigerians against diseases associated with low level of protein intake.

75

76 **PROSPECTS OF GRASS-CUTTER PRODUCTION**

77 Researches had been carried out on the nutritional requirement of grass-cutter (Mensah and Okeye, 2005, Henry and Njume, 2008), reproductive performance (Heath and Olusanya, 1985, Redford *et al.*, 1995, Ogunjobi, 2008), housing (Ogunjobi, 2008, Ogunjobi and Inah, 2008), environmental and reproductive biology (Williams *et al.*, 2011) disease and mortality (Kasali *et al.*, 1982; Fatokun *et al.*, 2010).

82

83 **Feeding and Nutritional requirement of grass-cutter**

84 The grass-cutter is primarily a herbivore, but in confinement, they require supplementary feed (Ayodele and Meduna, 2007). It is easy to feed and it is a good food transformer and practices coprophagy (Holzer *et al.*, 1986). The large caecum which forms part of the digestive tract of the grass-cutter is adapted predominantly to microbial digestion of feeds (Alaogninouwa *et al.*, (1996). Grass-cutter prefers mainly tick-stemmed grass species (Schrage and Yewadan 1999). The feeding habits of grass-cutter and other rodents (e.g. rabbits) are directly opposite. Whereas the grass-cutter prefers to eat stalks to leaves, the rabbit, for example on the contrary choose the leaves and waste stem (Vietmeyer 1991; Schrage and Yewadan 1999). This practice leads to waste of feed resources, especially during the dry season when there is scarcity of grass. Thus, irrespective of the kind of forage, grass-cutters first eat stalks, the bark of twigs and finally some leaves (Mensah and Okeyo, 2005).

This eating habit causes wearing of the animal's teeth, and unfortunately leads to high forage wastage (Adu, 2005).

They prefer grasses with lots of moisture and soluble carbohydrate (Onadeko, 1996; Agbelusi, 1997), preferring succulent grasses and stems like sugar cane (Adu, 1995; Awah, 2000) and Guatamala plants (Ndi, 2004). Grass-cutter can also be raised by feeding them with kitchen left-overs (Addo, 1997). Fitzinger (1995) reported that grass-cutters equally eat fallen fruits, nuts and many kinds of cultivated crops. Ebenebe (2005) identified some grasses and plants that are highly utilized by grass-cutter during the dry and wet season. These include: *Pennisetum purpureum* (elephant grass), *Saccharum officinarum* (sugar cane), *Zea mays* (maize), *Sorghum vulgare* (Guinea corn), *Oryza sativa* (rice), *Andropogon gayanus* (gamba grass) etc. Mensah and Okeyo (2005) reported that grass-cutters show high preference for grass and particularly favor *Pennisetum purpureum* (elephant grass) and *Panicum maximum*.

Good carcass quality and highest live weight was reported by Henry and Njume (2008) when grass-cutters were fed 2000KcalMEkg⁻¹ in combination with chopped elephant grass (*Pennisetum purpureum*). Feeds containing 12-20% crude protein (Meduna, 2002) have been reported to be suitable for grass-cutter, with Kusi *et al.* (2012) stating that 18% crude protein (CP) was required for optimum growth of grass-cutter from weaning to reproductive stage. According to Wogar (2011) gestating grass-cutter give optimum result in terms of litter size, birth weight of pups and feed conversion ratio when 14% CP is included in their diet. Wogar *et al.*, (2011) recommended that the optimum energy requirement of growing grass-cutter is 2200KcalME/kg in the humid tropics when the CP is 18% while Wogar (2012) stated that the preferred source of fibre for the growing grass-cutter is palm kernel cake.

Table 1 shows the nutrient requirement of grass-cutter while Table 2 gives the quantity of feed consumed by grass-cutter per day at different ages.

According to Holzer *et al.* (1986) and Mensah (2000) the water intake of the grass-cutter is reduced when the temperature is hot and more when the out-door temperature is low. They recommended this as a subject for further research since the reason for this unexpected behaviour was not yet understood.

Production System/ Housing

Production system of grass-cutter can be classified according to production objectives into commercial or subsistence, (Olomu *et al.*, 2003). Adu (2002) reported that the animal had

been bred and kept in boxes, empty drums, Poly Vinyl Chloride (PVC) pipes and enclosures among the rural communities and even in some urban areas among people with adequate space. The report by Ogunjobi (2008) showed that cane rat litters reared for six weeks after parturition in block-cement pens had comparative advantage in terms of weight gain and the rate of survival compared to those reared in iron cages. Ogunjobi and Inah (2008) therefore advocated for use of block pens (with wood shavings on the concrete floor) at the beginning of rearing grass-cutter as they recorded lesser deaths of the animal than those in iron cages. The housing of grass-cutter consist of stables and pens equipped with cages and hutches made of good materials and blocks, strong enough to stop the very sharp incisors of the animal (Fayenuwo *et al.*, 2003; Lameed, 2008).

Dentition in the grass-cutter

The dentition of the grass-cutter is typical of the rodentia with 10 pairs of teeth (Schrage and Yewaden, 1995; Fayenuwo *et al.*, 2003; Olomu *et al.*, 2003). These comprise of one incisor (I1), no canine (OC), one premolar (1P) and three molars (3M). Fayenuwo *et al.* (2001) reported that the incisors of grass-cutter are probably the most powerfully built of any African rodent. The time of eruption of the teeth helps the farmer to know the appropriate type of feeding to be offered to the animal and know the age of the animal. Table 3 shows a summary of a 51 weeks study on the eruption of grass-cutter teeth.

Reproductive performance of grass-cutter

Redford *et al.* (1995) reported that reproductive outputs are measured according to animal maturity, litter size, length of inter-birth interval and age at last reproduction. Ogunjobi (2008) identified some reproductive parameters of breeding grass-cutters to include having signs of mating on the 3rd day and 7th day, gestation interval of 155 ± 8 days and 157 ± 3 days, average litter size of 4 and 5, sex ratio of litters, 3 males : 1 female and 3 males : 2 females and mean weight gain of litter at six weeks of weaning 539 ± 12 g and 595 ± 12 g respectively for grass-cutter housed in iron cage (IC) and block-cement pen (BP) having same size of 120cm x 75cm x 30cm. Studies on the reproductive performance of female grass-cutter (does) at first parity in the humid tropical environment showed that 50% of the does studied had open vaginal status at first paring while the remaining 50% were either closed or plugged (Henry, 2010). The result also indicated that 50% of the does conceived at second exposure while 18.8% conceived at first exposure with more female off-springs. This confirmed the finding by Adjanohoun (1993) and Addo (1997) that grass-cutter and rabbits show variations in reproductive activity even though they are identified as induced ovulators.

Studies by Onadeko and Amubode (2002) revealed that the grass-cutter has a mean gestation length of 163.11 ± 1.58 days (with a range of 152-170 days), litter size of 4 (with a range of 2-7), mean birth weight of 117.70 ± 34.08 g (with male birth weight being generally heavier, 118.10 ± 27.70 g than females, 100.90 ± 27.50 g). They observed that breeding occurred in January, March, April, June, July, November and December with 67% of the parturition occurring at night. Further studies by the authors showed that litter weight decreased with increase in litter size, but did not influence the growth performance of the baby grass-cutter during the first months of life. A mean litter size of 4 had been previously reported by Amubode (1991).

Anatomy and morphology of grass-cutter

A good knowledge of the anatomical dispositions of the reproductive organs of grass-cutter is vital to the understanding of the reproductive biology of animals and provides information which would assist in the breeding of the cane rat and improve its domestication. The male reproductive organ of the grass-cutter is similar to that reported by previous authors (Massanyi *et al.*, 2003). The testes of the cane rat has ovoid shape with creamy white coloration covered with stroma (Dyce *et al.*, 2002; Olukole *et al.*, 2010), typical of rodents. The surface of the testes of the cane rat showed the presence of *Tunica vaginalis propria* with radical septa (*Septuli testes*) of pyramidal shape (Olukole *et al.*, 2009). The shape of the epididymis is sigmoid unlike that of a typical rodent and the distal part of the caudal epididymis is convoluted as in mammals (Olukole *et al.*, 2010) and provides useful information in the comparative regional anatomy of rodents. The morphology and morphometry of the grass-cutter male accessory sex gland had also been reported (Adebayo *et al.*, 2009). According to Ayodele and Meduna (2007) the best way of identify the different sexes is by studying the distance between the anus and the genital organs which is usually wider in the males and almost double of that obtainable in the female.

Environmental issues in grass-cutter production

The interaction of the grass-cutter in captivity with its immediate environment appears very important in relation to mortality, increase reproductive competence, improve health and overall performance. Williams *et al.* (2011) reported that a pit pen housing design was able to maintain the room temperature of the house and pens ($24.43 - 30.71^{\circ}\text{C}$) against the diurnal fluctuations outside the building ($25.86 - 34.71^{\circ}\text{C}$). They however stated that the relative humidity inside the building ($67.57 - 85.80\%$) showed a tendency to fluctuate with the

ambient relative humidity outside the building. They concluded that if captive grass-cutter were housed in environment similar to the night period, they are likely to shed their nocturnal habit and be more active during the daytime, with the likelihood of increased productivity.

Other benefits of the grass-cutter

Despite a lack of defined or measurable indications for its contribution to the gross domestic product (GDP), the grass-cutter subsector has been recognized as an important economic tool for rural poverty alleviation and household food and nutrition security (Yeboah and Adamu, 1995; Adu *et al.*, 2002). The grass-cutter is a considerable income earner for both the small scale peri-urban or rural livestock producer in the country. It also contributes to both local and export earning of countries like Kenya, Benin republic and Nigeria (Ntiamoa-Baidu, 1998). It is known to be economically important as an agricultural pest and its meat is widely accepted by all classes of people. The smoked grass-cutter could serve as a source of foreign earnings when it is well packaged and exported.

Grass-cutter meat is cheaper to produce than most other traditional livestock and its meat is more valuable and appreciated by local population. It has been showed that with only five mature grass-cutters (4 females and 1 male), a household is nutritionally secured for 6 months to one year (Juma and Ondwasy, 2002). In times of droughts and related calamities, grass-cutter serves as a critical source of animal protein.

During important occasions such as holidays, banquets and ceremonies, grass-cutter are heavily consumed by many households in Nigeria in both rural and urban areas, thus confirming that grass-cutter is the favorite bush meat species and accounts for the greater proportion pects, as ornamental, use in treatment of wound, preservation of ecological system, gift item (for example in most tribes in Nigeria, dried grass-cutter meat is used to serve the elders during traditional rights like marriages, excursion and Chieftaincy installations). The hair or fur is used to make decoration and the teeth are used to perform traditional card reading in place of cowries especially in South- South in most rural areas.

CHALLENGES OF GRASS-CUTTER PRODUCTION

Benjamin *et al.* (2006) reported some major problems encountered by grass-cutter farmers to include high initial capital, stock procurement, time constraint, inadequate medical attendant, disagreement with landlords over space to rear grass-cutter and inadequate follow-up by extension services. Recently, Ogunjobi and Inah (2008) ranked some constraints encountered

by the grass-cutter farmer from the most severe to the least as follows: lack of capital, insufficient feed, problem of disease, problem of housing, lack of enough land, problem of handling and lack of knowledge of rearing grass-cutter. Other challenges of grass-cutter challenges include its irregular supply, environmental issues, processing and marketing, feeding, producer's training and education, infrastructure development, poor information dissemination and disease/mortality.

Irregular supply

The production of grass-cutter is a relatively novel practice yet most breeding stock and cane rat meat is mainly obtained by hunting and trapping of the animals which does not ensure steady and regular supply of the meat (Taiwo *et al.*, 2009) as well as the breeding stock.

Environmental issues

The collection of grass-cutter from the wild is attended by the destruction of the environment through setting of bush fires by hunters (Ntiamoah-Baidu, 1980; NRC, 1991; Yeboah and Adamu, 1995). This leads to the destruction of valuable plants, animal life and tampers with the ecosystem (Taiwo *et al.*, 2009). Thus, there is need to domesticate the animals in order to avoid the problems associated with bush burning.

Poor processing and marketing plan

Most small scale and medium-scale farmers do not provide a good plan for processing and marketing of their grass-cutter at the initial stage of their business, as a result when the animal attain market weight, only a few buyers are seen. The farmer then devices a means of marketing (live or processed) grass-cutters while operating at a lost or reduced profit because of the extension in feeding time. This leads to the problem of storage facilities, waste disposal, and disease contamination, accompanied by reduction in selling price.

Lack of balanced diet

The domestication of cane rat has its own teething constrains, this include the need to provide regular supply of feeds rich and balanced in nutrients (Taiwo *et al.*, 2009). It has been observed that grass-cutter prefer grasses such as elephant grass, sugar cane, guinea grass with succulent stalk (Fayenuwo *et al.*, 2003) which may not be readily available. Furthermore, grass-cutter reared in captivity on forages and grasses alone does not do well compared to those living in the wild, this is because they normally obtain balanced nutrient from a variety of feeds such as grasses (forages tubers, grains, nuts, herb etc) in their natural habitat or in the wild.

Producer Training and Education

The education of farmers has been found to be one of the major factors affecting adoption of new technologies (Saha *et al.*, 1994). Intensified education on grass-cutter breeding and production to save the animal from extinction will reduce poverty and create employment. Fakoya *et al.*, (2008) reported that most grass-cutter farmers (90%) in Oyo State, Nigeria, had completed one form of formal education or another, implying that education is a variable which widens the mental horizon and predisposes farmers to new ideas. This results in having better access to knowledge and information that will be beneficial to the production and management of grass-cutter.

Infrastructure Development

Grass-cutter production is mainly concentrated in the rural areas which are characterized by poor infrastructural facilities such as road and telecommunication network. Installation of these facilities would open these areas for development (Kilungo and Mghenyi, 2001) and enhance access by the producers to market for purchase of inputs and sales of his products. Construction of good roads would help the extension services providers to reach as many producers as possible allowing training on new production technologies which will result in increased productivity of grass-cutter.

Poor information dissemination

The grass-cutter farmers in Nigeria as in other developing countries are faced with poor information dissemination about the challenges (such as disease out breaks, feeding, breeding, housing, marketing and lack of credit facilities) facing their production. Information is an essential ingredient in agricultural development programmes but Nigerian farmers seldom feel the impact of agricultural innovations either because they have no access to such vital information or because it is poorly disseminated (Ozowa, 1997). The extent of information needed by grass-cutter producers had been reported (Fakoya *et al.*, 2008). They identified grass-cutter diseases, grass-cutter housing pattern and equipment needed for production as the 1st, 2nd and 3rd most essential information needed by the grass-cutter farmer. Table 4 gives their findings and rating of the information needs of grass-cutter farmer. These problems can be solved through well-designed and implemented information dissemination and awareness programmes, including seminar which will endow all the stakeholders in the sector with necessary knowledge and skill (Gueye, 2009). Thus, there is need for networking amongst grass-cutter farmers to enable them acquire and share knowledge, views and experiences among themselves and with all other stakeholders along and grass-cutter value chain.

Mortality and Disease of grass-cutter

Disease is one of the most important limiting factors to profit in many livestock enterprise in the tropics (Hill, 1992). Majiyagbe and Lamorde (1997) reported that apart from inadequate and unbalanced feeding, high disease prevalence and associated high neonatal mortality constitute a major obstacle to the promotion of large scale holding of livestock. It is important to note that the severity of diseases depend on the nutritional state of the animals, especially during the dry season when feed is inadequate in quantity and quality (Opara and Fagbemi, 2009). Also, the incidence, severity and prevalence of disease had been shown to vary with the management system (Adu *et al.*, 2002).

Ogunjobi and Inah (2008) reported that more grass-cutter died when they were housed in iron cages at the beginning of farm operation than when they were housed in block-cement pens. In another research, Fatokun *et al.* (2010) observed that more grass-cutter (28) died of pneumonia among those kept in concrete cages with cemented floor while only (3) died among those kept in iron cages as a result of trauma and dystocia (difficulty in parturition). Kasali *et al.* (1982) observed the outbreak of intestinal coccidiosis in can rat while Onyeanusi and Famoyin (2005) identified twenty major disease/disease conditions affecting grass-cutter in captivity. Gastro-intestinal disorders, caused by helminthes parasite had also been identified in the grass-cutter (Awah-Ndukum *et al.*; 2001). Reports by Opara and Fagbemi (2008) showed that grass cutters can be infected with trypanosomes, although without obvious clinical disease.

Preliminary studies by Awah-Ndukum *et al.* (2001) on the captive grass-cutter in Cameroon showed the occurrence of ectoparasite such as fleas (*Xenopsylla sp*) and endoparasite like cestode (*Hymenolopsis sp*) and nematode (*Heterakis sp*). In another work by Yeboah and Simpson (2004) in Ghana, four species of tick namely *Rhipicephalus simpsoni*, *Ixodes aulacodi*, *Ixodes sp* and *Haemaphysalis parmata*, six species of helminthes comprising of 2 species of cestodes (*Furhmanella transvalensis*, *Railettina mahone*) and 4 species of nematodes (*Longistriata spira*, *Trachyphanyx natalensis*, *Paralibyostongylus vondwei* and *Trichuris paravispicularis*) were also found.

ROLE IN DISEASE TRANSMISSION

It is interesting to note that the grass-cutter had not been traced to harbor pathogens that can affect humans. Ebola virus disease for example, had been traced to chimpanzees, gorillas and bonobos, and currently spreading to humans when the meat is handled or consumed. Also,

gorillas and some other apes may also carry other diseases as simian foamy virus, smallpox, chicken pox, tuberculosis, measles, rubella, yellow fever and yaws (<http://en.wikipedia.org/wiki/Bushmeat>). It has occurred, on numerous occasions, that people who ate apes have caught such diseases, or even died (BBC News, 2009). Thus, apart from posing a significant risk to the people who eat the meat, it poses great risk to the human population as a whole, as it opens a doorway through which animal viruses can be transmitted to humans. Other bush meat like the African squirrels (*Heliosciurus*, *Funisciurus*) have also been implicated as reservoirs of the monkey pox virus in the Democratic Republic of Congo (<http://en.wikipedia.org/wiki/Bushmeat>), implying that the use of their meat may serve as a means of transmission of these viruses to humans. The commercial production of grass-cutter will therefore be of great benefit to lovers of bush meat since the animal had not been linked to pathogens.

CONCLUSION

This study has revealed that the grass-cutter, a wild African rodent can be successfully domesticated. The feeding and nutritional requirement, production system and housing, dentition, reproductive performance, anatomy and morphology, environmental issues and the benefits have all been studied. Detailed information on the challenges, including irregular supply of the animal, environmental issues, processing and marketing, feeding, poor producer training and education, infrastructural development, poor information dissemination, mortality and diseases among others have also been reviewed. This animal which provides juicy and palatable meat, and is highly preferred in meals of both Nigerians and foreigners, without any fear of disease transmission, therefore, offers suitable opportunities for enhancing livelihood and revenue generation in rural and urban areas in the sub-region. Consequently, this review creates more opportunity for the grass-cutter farmers and intending farmers to easily sustain their business, create more jobs, increase income and increase protein consumption of the growing populace while it serves to assure the consumers of bushmeat (commercial grasscutter in this case) of the safety of the meat.

REFERENCES

- Addo, P.G. (1997). Domesticating the wild grasscutter (*Thryonomys swinderianus* Temminck 1827) under laboratory conditions. Ph.D Thesis, Univ. of Ghana, Legon, Ghana.

- 363 Adebayo, A.O, Oke, B.O and Akinloye, A.K. (2009). The morphology and morphometry of
364 the male accessory sex gland in the greater cane rat (*Thryonomys swinderianus*
365 Temminck). Book of Abstract of the 46th Ann. Congress of the Nig. Vet. Med. Assos.
366 (NVMA). pp.6
367
- 368 Adekola, A. G. and Ogunsola, D. S. (2009). Determinants of Productivity level of
369 commercial grasscutter farming in Oyo state. Proc. Int'l. Conf. on Global Food
370 Crisis, 19th-24th April, Owerri, Nigeria. Pp. 15-21.
371
- 372 Adjanohoun, E. (1993). Quelques aspects du cycle sexual de l'aulocode () et leurs
373 consequences pratiques sur la conduit des elevages. L'er Conference Internationale
374 L'Aulocodicuture: Acquis et perspective (Schrage, R. and Yewaden, L.T. eds) pp.
375 111-118
376
- 377 Adu, J.A. (1995). Manual of Grasscutter Husbandry. Adagro Lagos. Pp 25-30
378
- 379 Adu, E.K. (2002). Dexieme Conference on promoting broadcast from the breeding of the
380 cane rats in Africa. Su DuSAHARA.
381
- 382
- 383 Adu E. K. (2005). Constraints to grass-cutter production in Ghana. Proc. of the Int'l Forum
384 on grass-cutter. Institute of Local Government Studies. Accra. Ghana, T. Antoh., R.
385 Weidinger., J. Ahiaba and A. Carrilo (eds). Ministry of Food and Agriculture. Accra.
386 Ghana. pp. 44-50.
387
- 388 Adu E. K. Wallace P. A. and Ocloo T. O. (2002). Efficacy of sex determination in the greater
389 cane rate, *Thryonomous swinderianus*, Temminck. Trop. J. Anim. Prod. 34 (1): 27-33.
390
- 391 Agbelusi, E. (1997). Ranching grasscutter (*Thryonomys swinderianus* Temminck) for
392 meat production in the humid forest zone of Nigeria. FAO Jul Publication
393
- 394 Ajayi, S. S. and Tewe, O. O (1980). Food preference and carcass composition of grass-cutter
395 (*Throyonomis swinderianus*) in captive. African. J. Eco. 18: 13-14.
396
- 397 Akinola, L.A.F. (2008). Grasscutter farming: A new initiative in protein supply. An Invited
398 paper presented at Agricultural Product Development Strategy Workshop organized
399 by Rivers State Sustainable Development Agency (RSSDA), held on 9-10th Sept.
400 2008 at The Elkan Terrace, Abacha Road, Port Harcourt, Rivers State, Nigeria
401
- 402 Akpan I. A. Wogar, G. S. I., Effiong, O. O. and Akpanenua, E. J. (2009). Growth
403 performance of grass-cutter fed diets treated with urea and urine solution. Proc. of
404 34th Ann. Conf. of Nig. Soc. for Anim. Uyo, Nigeria. Pp. 163-164
405
- 406 Alaogninouwa, T., Agba, K.C., Agossou, E. and Kpodekon, M. (1996). Anatomical,
407 histological and functional specificities of the digestive tract in the male grasscutter
408 (*Thryonomys swinderianus* Temminck). Anatomy, Histology, Embryol., 25:15-21
409
- 410 Amubode, F.O. (1991). Basic information for captive rearing of grasscutter. The
411 Conservation. 1: 33-34
412

- 413 Anieunam, A.S. (2005). The place of bush meat in the supply of animal protein in Nigeria.
414 Seminar Paper, Department of Animal Science, University of Calabar, Nigeria.
415
- 416 Asibey, E. O. A. and Addo, P. G. (2000). The grass-cutter, a poisoning animal for meat
417 production, in: African Perspective, practices and policies supporting sustainable
418 development (Turnham, D. Ed) scandinavian Seminar College, Denmok; in
419 Association with weaver press Harare. Zimbabwe. [www.cdr.dk/sscafrica/asad-gh.](http://www.cdr.dk/sscafrica/asad-gh.htm)
420 [htm](http://www.cdr.dk/sscafrica/asad-gh.htm).
- 421 Awah, A.A. (2000). Introduction to minilivestock development as a sustainable agricultural
422 business. Workshop Paper presented at Agric. Dev. Prog. (ADP) in Anambra, Imo,
423 Rivers and Cross Rivers States.
424
- 425 Awah-Ndukum, J., Tchoumboue. And Tong, J.C. (2001) Stomach impaction in grass-cutter
426 (*Thryonomys swinderianus*) in captivity: Case Report. Trop. Vet. 19(2): 60-62.
427
- 428 Ayodele, I.A and Meduna, A.J. (2007). Essentials of Grasscutter Farming. Hope Publications,
429 Ibadan, Nigeria. Pp. 37 BBC News (2009). Scientist find new strain of HIV. 2
430 August 2009
431
- 432 BBC News (2009). Scientist find new strain of HIV. 2 August 2009
433
- 434 Benjamin, U.U., Akinyemi, A.F. and Ijeomah, H.M. (2006). Problems and prospects of
435 grasscutter (*Thryonomys swinderianus*) farming in Ibadan, Nigeria. J. Agric.
436 Forestry and the Social Sci. (JOAFSS). 4(2):24-32.
437
- 438 Clotley J.A. (1981). Relation of physical body composition to meat yield in the grass-cutter
439 (*Thryonomys swinderianus* Temminck). Ghana Journal of Science.
440
- 441 Dyce, K.M., Sack, W.O. and Wensing, C.J.C. (2002). Textbook of Veterinary Anatomy. 2nd
442 Ed. W.B. Saunders
443
- 444 Ebenebe, C.I. (2005). Identification and analysis of some plants utilized by the grasscutter
445 (*Thryonomys swinderianus* Temminck) at Akpaka forest reserve, Onitsha, Nigeria.
446 Proc. of 10th Ann. Conf. of Anim. Sci. Assos. of Nig (ASAN). Univ. of Ado-Ekiti,
447 Nigeria. pp. 251-254
448
- 449 Fakoya, E.O., Sodiya, C.I., Alarima, C.I. and Omoare, A.M. (2008). Information needs of
450 farmers in grasscutter production for improving performance in Ona Ara Local
451 Government Area of Oyo State. Proc. of 33rd Ann. Conf. of Nig. Soc. for Anim. Prod.
452 Ayetoro, Ogun State, Nigeria. pp. 300-301
453
- 454 Fatokun, B.O., Ogunjobi, J.O., Olajide, O.B. and Ukandu, P. (2010). Clinical investigations
455 on the causes of mortality in grasscutter (*Thryonomys swinderianus*, T. 1827) held in
456 captive within Ibadan metropolis in Oyo State, Nigeria. Proc. of 35th Ann. Conf. of
457 Nig. Soc. for Anim. Prod. held at Univ. of Ibadan, Nigeria. pp 751-753.
458
- 459 Fayenuwo, J.O., Akande, M., Ogundola, F.I. and Oluokun, J.A. (2001). Moor J. Agric. Res.
460 (2): 141-146
461

- 462 Fayenuwo, J.O., Akande, M., Taiwo, A.A., Adebayo, A.O., Saka, J.O., Lawal, B.O.,
463 Tiamiyu, A.K. and Oyekan, P.O. (2003). Guidelines for Grasscutter Rearing.
464 Technical Bulletin. IAR&T Ibadan Nigeria. 38pp
465
- 466 Fayenuwo, J.O., Taiwo, A.A., Oluokun, J.A., Akande, M. and Adebowale, E.A. (2005).
467 Observations on teeth development in captive-bred grasscutter (*Thryonomys*
468 *swinderianus* Temminck). Proc. Of 10th Ann. Conf. of Anim. Sci. Assos. Of Nig.
469 Univ. of Ado-Ekiti, Nigeria. pp. 248-250
470
- 471 Fitzinger, F. (1995). Can Rats. In: Walker's Mammals of the World, Nowak, R (Ed). The
472 Hopkins Univ. Press, Maryland. pp. 1650-1651
473
- 474 Fonweban, J. N. and Njwe, R. M. (1990). Feed utilization of life weight gain by the Africa
475 giant rat (*Cricetomys gambianus*, water house) at Dischana in Cameroon.
476 Tropiculture, 8 (3) 118-120.
- 477 Gueye, E. F. (2009). The role of networks in information dissemination to family poultry
478 farmers. World's Poultry Science Journal 65: 115-124.
479
- 480 Heath, E. and Olusanya, S. (1985). Anatomy and Physiology of Tropical Livestock. Longman
481 Scientific and Technical. 138
482
- 483 Henry, A.J. (2010). Reproductive performance of grasscutter does at first parity reared in
484 humid tropical environment. Proc. of 35th Ann. Conf. of Nig. Soc. for Anim. Prod.
485 Univ. of Ibadan, Nigeria. pp. 155-158
486
- 487 Henry, A.J. and Njume, G.N. (2008). Effect of varied energy levels on the carcass
488 characteristics of grasscutter (*Thryonomys swinderianus*). Proc. Of 33rd Ann. Conf.
489 of the Nig. Soc. for Anim. Prod. Ayetoro, Ogun State, Nigeria. Pp 168-170.
490
- 491 Hill, D. (1992). Cattle and Buffalo Meat Production in the Tropics. Longman Scientific and
492 Technical. 210pp.
493
- 494 Holzer R. Mensah G. A. and Baptist R. (1986). Practical aspect of grass-cutter (*Thryonomys*
495 *swinderianus*) breeding, particulars of coprophagy. Rev. Elev. Med-vet. Pays. Trop:
496 39 (2): 247-252.
497
- 498 Juma, N. and Ondwasy, H.O. (2002). Improved management of indigenous chicken
499 sustainable technologies contributing to the socio-economic welfare of rural
500 household. Proc. Of the 8th Kenya Agricultural Research Institute Biennial Scientific
501 Conference, Nairobi, Kenya. pp 359-364
502
- 503 Kasali, O.B., Majaro, O.M. and Dipeolu, O.O. (1982). An outbreak of intestinal coccidiosis
504 in a colony of can rats in Nigeria. Nig. J. Vet. Med. 9(2): 3-5
505
- 506 Kilungo, J.K and Mghenyi, E. (2001). Factors limiting beef productivity and marketing in
507 Kenya. Working Paper, Tegoneo Institute, Egerton Univ., Nairobi, Kenya
508
- 509 Kusi, C., Tuah. A.K., Annor, S.Y. and Djang-Fordjour, K.T. (2012). Determination of dietary
510 crude protein level required for optimum growth of the grasscutter in captivity.

- 511 Livestock Res. for Rural Devt.. 24(10).www
512 Irrd.cipav.org.co/Irrd24/10/kusi24176.htm
513
- 514 Lameed, G.S.A. (2008). Grasscutter/ Canerat (*Thryonomys swinderianus*) Farming in the
515 Tropics: The possibilities and Prospects for would-be Farmers. Peerless Prints,
516 Nigeria. pp. 14-21
517
- 518 Majiyagbe, K.A. and Lamorde, A.G. (1997). Nationally Cordinated Research Programme on
519 Livestock Disease: Subsectoral goals, performance and medium-term research plan.
520 Tropical Veterinary. 15: 75-83
521
- 522 Martin, G.H. (1985). West Africa: Carcass composition and palatability of some animal
523 commonly used as food world animal review. (53): 40-44.
524
- 525 Massanyi, P., Jancova, A, and Uhrin, V. (2003). Morphometric study of male reproductive
526 organs in the rodent species *Apodemus sylvaticus* and *Apodemus flavicollis*. Bull. Vet.
527 Inst. Pulway. 47: 133-138
528
- 529 Meduna, A.J. (2002). Preliminary observation on cane rat (*Thryonomys swinderianus*,
530 Temminck) feeding and breeding. Proc. Of the 27th Ann. Conf of the Nig. Soc. For
531 Anim. Prod. Federal College of Tech., Akure. pp 304-305
532
- 533 Mensah G.A. (1993). *Futteraufnahme und Verdaulichkeit beim Grasnager (Thryonomys*
534 *swinderianus)*. Thèse de Doctorat, Institut 480, Université de Hohenheim, Allemagne,
535 107 pp.
- 536 Mensah G.A. 1995. Consommation et digestibilité alimentaire chez l'aulacode *Thryonomys*
537 *swinderianus*. *Tropicultura*, 13 (3):123-124.
538
- 539 Mensah, G.A. (2000). General presentation of the breeding of grass-cutter, history and
540 distribution in Africa. Proc. Of Int'l Seminar on Intensive Breeding of Wildlife
541 Animals for the Purpose of Food in Liberville, Gabon Project,
542 DGEG/VSF/ADIE/CARPE/UE., pp. 45-59
543
- 544 Mensah G. A. and Okeyo, A. M. (2005). Continued harvest of the divers African Animal
545 Genetic Resources from the wild through domestication as a strategy for sustainable
546 use: A case of the larger grass-cutter (*Thryonomys swinderianus*).
547 <http://agtr.ilri.cigar.org/Case-study/Mensah/Mensah.htm>.
548
- 549 National Research Council (1991). Microlivestock: Little known small Animals with a
550 Promising Economic Future, National Academy Press, Washington DC, USA
551
- 552 Ndi, M.J. (2004). Cane Rat Farming Manual. Cameroun Youth Dev. Centre. Promotion for
553 grasscutter production. No. 3 Series. pp. 7-200
554
- 555 Ntiamoa-Baidu, V. (1998). Sustainable use of bush meat. Wildlife Devt. Plan. 1998-2003
556
- 557 Ogunjobi, J.A. (2008). Reproductive performance of cane rat (*Thryonomys*
558 *swinderianus* Temminck 1827) breeding stocks and littesr rate of survival reared

- 559 using two common housing materials. Proc. of 33rd Ann. Conf. of Nig. Soc. for Anim.
560 Prod. Ayetoro, Ogun State, Nigeria. pp. 208-210
561
- 562 Ogunjobi, J.A. and Inah, E.I. (2008). Seasonal mortality among farmed cane rat reared using
563 two different housing materials. Proc. of 33rd Ann. Conf. of Nig. Soc. for Anim. Prod.
564 Ayetoro, Ogun State, Nigeria. pp 136-138
565
- 566 Ogunsanmi, A.O., Ozegbe, P.C., Ogunjobi, O., Taiwo, V.O. and Adu, J.O. (2002).
567 Haematological, plasma biochemistry and whole blood minerals of the captive adult
568 African grasscutter (*T. swinderianus* Temminck). Trop. Vet. 20:27
569
- 570 Olomu, J.M., Ezieshi, V.E. and Orheruata, A.M. (2003). Grasscutter Production in Nigeria.
571 Principles and Practice. Jachem Publishers. pp. 62
572
- 573 Olukole, S.G., Oyeyemi, M.O. and Oke, B.O. (2009). Biometrical observations on the testis
574 and epididymis of the domesticated adult African great cane rat (*Thryonomys*
575 *swinderianus*). Eur. J. Anat., 13(2): 71-75.
576
- 577 Olukole, S.G., Oyeyemi, M.O. and Oke, B.O. (2010). Gross anatomy of male reproductive
578 organs of the domesticated grasscutter (*Thryonomys swinderianus* Temminck).
579 Proc.of 25th Ann. Conf. of Nig. Soc. For Anim. Prod. Univ. of Ibadan, Nigeria. Pp
580 268-271
- 581 Omole,A.J., Ayodeji, I.O., Ashaye, O.A. and Tiamiyu, A.K. (2005). Effect of scalding and
582 flaming methods of processing on physiochemical and organoleptic properties of
583 grasscutter meat. J. Applied Sci. Res. 1:249-252
584
- 585 Onadeko, S.A. (1996). The reproductive ecology of the grasscutter (*Thryonomys*
586 *swinderianus*) in captivity. Ph.D Thesis, Dept. of Wildlife and Fisheries
587 Management, Univ. of Ibadan.
588
- 589 Onadeko, S.A. and Amubode, F.O. (2002). Reproductive indices and performance of captive
590 reared grasscutter *Thryonomys swinderianus* Temminck (). Nig. J. Anim. Prod. 29(1):
591 142-149.
592
- 593 Onyeausi, A.E. and Famoyin, J.B. (2005). Health care management of grass-cutter in
594 captivity: Assessment of causes of mortalities among rearing stock in Ibadan
595 Metropolis. J. of Forestry Research and Management. 2:58-66
596
- 597 Opara M. N. and Fagbemi, B. O (2009). Therapeutic effect of Berenil in experimental murine
598 trypanosomiasis using stocks isolated from apparently healthy wild grass-cutters.
599 (*Thryonomys swinderianus*). Proc. Intl. Conf. on Global Food Crisis, April 19th-24th,
600 Owerri, Nigeria pp. 31-37.
601
- 602 Opara,M.N. (2010). The grasscutter 1: A livestock of tomorrow. Res.J. Forestry. 4:119-135
603
- 604 Opara,M.N. and Fagbemi, B.O. (2008). Haematological and plasma biochemistry of the adult
605 wild African grass-cutter (*Thryonomys swinderianus*). Azoonosis factor in the tropical
606 humid rain forest of southeast Nigeria. Ann. N.Y.Acad. Sci. 1149:394-397.
607

- 608 Owen, O. J. Alawa J. P. Wekhe S. N. Isiriman N. O, Ngodigha E. M. and Amakiri A. O.
609 (2009) Incorporating poultry litter in rabbit feed: A solid waste management
610 strategy. Egyptian J. Anim. Prod. 46 (1) 63-68.
611
- 612 Owen, O. J. and Amakiri A. O. (2009). Rabbitary in Nigeria. The potentialities, opportunities
613 and challenges: A review Proc. Int'l. Conf. on Global Food Crisis, FUT, Owerri,
614 April 19-24th. Pp. 79-82.
615
- 616 Ozowa, V.N. (1995). Information needs of small scale farmers in Africa: The Nigerian
617 Example. Quarterly Bulletin of the International Association of Agricultural
618 Information Specialists. IAALD?CABI. 40(1).
619
- 620 Redford, K.H., Godhalk, R. and Asher, K. (1995). What about the wild animals? Wild animal
621 species in community forest in the tropics. Community Forestry Note 13. Food and
622 Agricultural Organization of the United Nations, Rome
623
- 624 Redford, K.H., Godshalk, R. and Asher, K. (1995). What about the wild animals (mammals)
625 of some economic importance: Problems and Prospects. Proc. of the National
626 Forestry Workshop on Strategies for the management of endangered environment and
627 species. R.D. Adegbehin and R.D. Gbadegesin eds. pp 161-178
628
- 629 Saha, A., Love, H.A. and Schwart, R. (1994). Adoption of emerging technologies under
630 output uncertainty, *American Journal of Agriculture Economics* 76: 836-846.
631
- 632 Schrage R. and Yewandan L. T. (1999) In: Raising Grass-cutters Deutsche Gesellschaft for
633 technische Zusammenarbeit (GTZ) GmbH, Eschborn, Germany, 99 pp.
634
- 635 Schrage, R. and Yewaden, L.T. (1995). Abrege D'aulacodiculture. Deutsche Gesellschaft for
636 Technische Zusammen Anbait (GTZ) GmbH. pp. 103
637
- 638 Taiwo A. A Fayenuwo J. O. Omole A. J. Fajimi A. K. Fapohunda J. B. and Adebowale E. A.
639 (2009). Nig. J. Anim Prod. 36 (1): 153-160.
640
- 641 Taiwo, O.W. (2006). Wealth Creation through Commercial Grass-cutter Farming. Production
642 Techniques for Good Performance. 67pp.
643
- 644 The Thy Consulting (2006). How to become your own boss through high profitable
645 Agricultural businesses. Grasscutter farming. 11p
646 <http://grasscutterfarming.tripod.com>
647
- 648 Van-Zyl, A., Van der Merwe, M. and Blignaut, A.S. (1999). Meat quality and carcass
649 characteristics of the Vondo, *Thryonomys swinderianus*. South African Journal of
650 Animal Science. 29:120-121
651
- 652 Vietmeyer N. D. (1991). Board on Science and Technology for International Development
653 National Research Council, National Academic Press. Washington DC. pp. 233-240.
654
- 655 Williams, O.S., Ola, S.I., Boladuro, B.A. and Badmus, R.T. (2011). Diurnal variation in
656 ambient temperature and humidity in a pit pen grass-cutter (*Thryonomys*

657 *swinderianus*) house in Ile- Ife. Proc. 36th Ann. Conf. of Nig. Soc. For Anim. Prod.
658 held at Univ. of Auja, Nigeria. pp. 111-113
659
660 Wogar, G.S.I. (2011). Performance of gestating grasscutter (*Thryonomys swinderianus*) fed
661 cassava-based diets with graded protein levels. Asain J. of Anim. Sci. 5:373-380 .
662
663 Wogar, G.S.I., Effiong, O.O. and Nsa, E.E. (2011). Performance of growing grasscutter
664 (*Thryonomys swinderianus*) fed diets with graded energy levels Journal of Agric.,
665 Biotechnology and Ecology. 4(3): 134-139
666
667 Wogar, G.S.I. (2012). Performance of growing grasscutter on different fibre sources. Pakistan
668 Journal of Nutrition. 11(1): 51-53
669
670 Yeboah, S. and Simpson, P.K. (2004). A preliminary survey of ecto and endo parasites of the
671 grass-cutter (*Thronomys swinderianus* Temminck) case study in Ekumfi Central
672 Region of Ghana. J. of the Ghana Sci. Assos. 3(3): 2-5.
673
674 Yehoah, S. and Adamu, E. K. (1995). The Cane Rate Biologist. 42 (2): 86-87
675
676 <http://en.wikipedia.org/wiki/Bushmeat> (Retrieved 16th September, 2014)
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706



Figure 1: A family of grass-cutter



Figure 2: A mature Grass-cutter

Table 1: Nutrient requirement of an adult grass-cutter

Component	% dry matter basis
Crude protein	12 to 18.5
Crude lipid	2.5 to 4.5
Crude fibre	25 to 45
Ash	2.5 to 4.5
Nitrogen free extract	45 to 65
Neutral detergent fibre	42 to 64
Acid detergent fibre	25 to 35
Acid detergent lignin	3 to 8

Source: Mensah (1993; 1995)

Table 2: Quantity of feed consumed by grass-cutter per day

Subject	Grass/forage (g)	Supplement (g)
Cutling (Young grass-cutter)	10 – 150	10 -50
Weaner/grower grass-cutter	152 – 250	51 – 100
Adult grass-cutter	251 – 450	101 – 200

Source: Fayenuwo *et al.* (2003)

725

726

727

728 Table 3: Cutting of teeth (teeth eruption) periods in grass-cutter

Age/Period	Teething per half jaw				No. of teeth (%)
	Incisor (I)	Canine (C)	Premolar (P)	Molar (M)	
Birth	P1 ¹ (n=51)	-	Op ¹ (n=51)	-	40
2 weeks	„	-	Ap ¹ (n=48)	Om ¹ (n=48)	60
2-4 months	„	-	Pp ¹ (n=42)	Pm ¹ (n=42)	60
5-8 months	„	-	„	Pm ² (n=35)	80
9 months	„	-	„	Pm ³ (n=33)	100

729 Source: Fayenuwo *et al.* (2005)

730 = Tooth present = Tooth absent

P1¹ = Presence of one incisor

Ap¹ = Appearance of the premolar

Om¹ = Opening of the 1st molar

Pm² = Presence of the 2nd molar

Dental formular:

I(1) C(0) P(1) M(3) = 20

Op¹ = Opening of premolar

Pp¹ = Presence of premolar

Pm¹ = Presence of 1st molar

Pm³ = Presence of 3rd molar

I =Incisor, C = Canine

P = Premolar, M = Molar

731

732 Table 4: Information needs of farmers in grass-cutter production

S/N	Information Needs	Scores
1	Rabbit housing pattern	2 nd
2	Cleaning of housing unit	7 th
3	Sources of stable grass-cutter breed	9 th
4	Types of feed available	12 th
5	Weaning	8 th
6	Equipment required for grass-cutter production	3 rd
7	Appropriate number of grass-cutter required in cages	15 th
8	Incentive on grass-cutter	12 th
9	Identifying various grass-cutter disease	1 st
10	Selection of foundation stock	11 th
11	Marketing of grass-cutter	5 th
12	Record keeping	4 th
13	Control of pests and diseases of grass-cutter	6 th
14	Method of mating	10 th
15	Ovulation and heat period	14 th

733 Source: Fakoya *et al.* (2008)