

Original Research Article

EFFECT OF COWDUNG, NPK AND THEIR COMBINATIONS ON SOIL PROPERTIES AND PERFORMANCE OF SWEET POTATO (*Ipomoea batatas* L.) IN SUDAN SAVANNA, NIGERIA

Abstract

In Nigeria, one of the major problems confronting agriculture is poor soil fertility replenishment strategy that could allow sustainable production. Two field trials were conducted simultaneously at Sokoto (UDUS, Vegetable Research farm) and Zamfara (Bakalori Irrigation Project) in Sudan-Savanna agro ecological zone of Nigeria with the aim of studying the effect of different rates of cowdung, NPK and their combination on soil properties and performance of sweet potato. The research was conducted during the 2011/2012 dry season using ten treatments: CD(H), CD(M), CD(L), NPK(H), NPK(M), NPK(L), CD+NPK(H) CD+NPK(M), CD+NPK(L) and control. The result was consistent with regard to locations and indicated no significant effect on soil pH. However, the treatments resulted to a decrease in the soil pH from 5.63 to (5.38-4.95) and 5.90 to (5.88-5.01) at UDUS and Bakalori respectively. No significant effect was also recorded on soil organic carbon, CEC, percentage nitrogen, available phosphorus, exchangeable calcium and potassium but was found increased due to treatments application as compared to the initial values. NPK(M) was determine to have significantly ($p<0.05$) the highest sweet potato performance at both location. This research therefore recommended the treatments used as important materials for soil nutrients replenishment and NPK(M) as the recommended rate of fertilizer for better sweet potato production at the study areas.

Key words; fertilizer, Rates, Soil properties and sweet potato

33

34

35 Introduction

36 Plant nutrients are essential for the production of qualitative crops to provide healthy food for the
37 world's expanding population. Plant nutrients are therefore, a vital component of sustainable
38 agriculture. Increased crop production largely relies on the type of fertilizer used to supplement
39 essential **nutrient** for plants. The nature and the characteristics of nutrients **release** by mineral and
40 organic fertilizer are different and each type of fertilizer has its advantages and disadvantages
41 with regard to crop growth and soil fertility. The sound management of fertilization must attempt
42 to ensure both an enhanced crop productivity and safeguarded environment. Therefore, a
43 balanced fertilization strategy that combines the use of mineral and organic fertilizers must be
44 developed and evaluated (Chen, 2006).

45 Sweet potato (*Ipomoea batatas*L.) belongs to the family convolvulaceae. The family has about 45
46 genera and 1000 spp, but only sweet potato (*Ipomoea batatas* L.) is of economic **important** as
47 food [Nwauzor et al., 2010). It is an important food and vegetable crop that is widely distributed
48 throughout the world particularly in the Tropics [Nwauzor et al., 2010). Sweet potato originated
49 from Tropical America and has been cultivated probably from B.C. Today, the crop remains one
50 of the three most important root crops in the world, following potato (*Solanumtuberosum*) and
51 cassava (*Manihotesculenta*) [Nwauzor et al., 2010).

52 In an effort to reduce over and under application as well as establishing appropriate types of
53 fertilizer that will enhance both nutrient availability to plant and the soil fertility, the need
54 therefore, arises for assessing the comparative effect of different rates of organic, mineral and
55 organo-mineral fertilizers on the soil, growth and yield of crops. This research was therefore,
56 aimed at assessing the effect of different rates of cow dung, NPK and their combinations on soil
57 properties, growth and yield of sweet potato in Sudan savanna agro ecological zone of Nigeria.

58 Materials and Method

59 The trials were conducted in Sokoto and Zamfara States, simultaneously. For Sokoto (L1); the
60 trial was conducted at the UsmanuDanfodiyo University, Vegetable Research Farm located at
61 Kwakwalawa village in dundaye district of Sokoto state. The area lies on latitude 13° 01'N:
62 longitude 5° 15'E and altitude of about 350 m above Sea Level (ASL) and located along the
63 main road to UsmanuDanfodiyo University, main campus, Sokoto. InZamfara (L2); the trial was
64 conducted at the Bakalori Irrigation Project. The area lies on both sides of the Sokoto river
65 between latitude 12° 30'-12° 50'N and longitude 5° 59'-6° 20'E near TalataMafara, about
66 110km from Sokoto on the Sokoto – Gusau road in Zamfara State.

67

68

69

70 The treatments consisted of three rates of cow dung, three rates of NPK (15:15:15), three rates of
71 cowdung and NPK combinations and control. The ten treatments were arranged in a randomized
72 complete block design (RCBD) replicated three times. The treatments were;

73 CD (H) = 3t/ha cowdung

NPK (L) 200kg/ha

74 CD (M) = 2t/ha cowdung

CD+NPK (H) = 2t cowdug + 100kgNPK/ha

75 CD (L) = 1t/ha cowdung

CD+NPK (M) = 1.5t cowdung +150kgNPK/ha

76 NPK (H) = 400kg/ha NPK

CD+NPK (L) = 1t cowdung + 200kgNPK/ha

77 NPK (M) = 300kg/ha NPK

CONTROL = No fertilizer applied

78 Cow dung was applied in full dose and worked into the soil one week before transplanting; NPK
79 was applied in split dose: first dose at two weeks after transplanting and the remaining dose at 50
80 days after the first application, using broadcasting method.

81 Data was collected for the following growth and yield parameters; Number of leaves and vines
82 per plant, vine length, number of tubers per plant, tuber weight and tuber yield. Soil samples
83 were collected using soil auger at depth of (0-30cm) before and after the experiment and
84 analyzed for soil pH, organic carbon, cation exchange capacity (CEC), total nitrogen and
85 available phosphorus. Data was analyzed using (ANOVA) and DNMRT for mean separation.

86 Results and Discussion

87 Chemical composition of the cowdung used

88 Table 1; Chemical composition of the cowdung used

89 Chemical properties	values
90 pH (H ₂ O) 1:1	8.07
91 Org. carbon (g/kg)	34.89
92 Total nitrogen (%)	0.66
93 Avail. Phosphorus (mg/kg)	0.18
94 Exchangeable K (cmol/kg)	0.50

95

Soil analysis before the experiment

Table 2: soil analysis before the experiment during 2011/2012 dry season at UDUS (kwalkwalawa) and Bakalori

Parameters	UDUS(kwalkwalawa)	Bakalori
pH (H ₂ O) 1:1	5.63	5.9
Organic carbon (g/kg)	0.33	0.45
Total nitrogen (%)	0.16	0.17
Avail. Phosphorus (mg/kg)	4.74	4.67
CEC (cmol/kg)	3.41	3.94
Exchangeable Ca (cmol/kg)	0.29	0.32
Exchangeable Mg (cmol/kg)	0.52	0.72
Exchangeable K (cmol/kg)	0.35	0.40

Result presented in table 2 showed the chemical composition of soils before the experiment at the two locations that, soil pH were moderately acidic, low organic carbon, CEC, available phosphorus and medium in total nitrogen.

Soil analysis after the experiment

Table 3: Effect of Cow dung, NPK and their Combinations on Soil pH, Organic Carbon and CEC at UDUS and Bakalori in 2011/2012 Dry Season.

Treatment	L1 (UDUS)					L2 (BAKALORI)					
	pH	O/C (%)	CEC (cmol/kg)	T/N(%)	P(mg/kg)	pH	O/C (%)	CEC (cmol/kg)	T/N(%)	P(mg/kg)	
CD(H)	5.33	0.66	5.03	0.23	4.60	5.80	0.61	4.76	0.18	4.37	
CD(M)	5.31	0.57	5.16	0.18	4.94	5.81	0.55	5.88	0.2	4.89	
CD(L)	5.27	0.34	4.70	0.16	4.22	5.81	0.47	3.70	0.21	4.52	
NPK(H)	4.95	0.43	5.00	0.19	4.76	5.01	0.35	5.40	0.18	4.42	
NPK(M)	5.15	0.49	4.93	0.15	4.66	5.59	0.27	5.35	0.16	4.71	
NPK(L)	5.12	0.55	4.73	0.15	4.56	5.43	0.39	4.83	0.23	4.30	
CD+NPK(H)	5.38	0.68	4.83	0.19	4.93	5.63	0.46	4.20	0.2	4.72	
CD+NPK(M)	5.23	0.41	4.73	0.2	4.80	5.88	0.54	5.46	0.2	4.36	
CD+NPK(L)	5.26	0.47	5.06	0.35	4.54	5.70	0.41	4.90	0.2	4.64	
Control	5.10	0.38	4.73	0.16	4.78	5.77	0.41	3.43	0.12	4.62	
Level of Sig.	ns	ns	ns	ns	ns	ns	ns	ns	ns	ns	
SE	0.05	0.05	0.08	0.01	0.05	0.06	0.03	0.26	0.02	0.06	

ns = Not significant at 5% level of probability

T/N = total nitrogen

P = phosphorus

Effect of cowdung, NPK and their combinations on soil chemical properties is presented In Table 3. Result indicated that, treatments had no significant effect ($p>0.05$) on soil pH, organic carbon, CEC, total nitrogen and available phosphorus at both the locations. This result may be due to the time frame to which the experiment was conducted as related to the rate of decomposition, mineralization and crop quality. However, as compared to the initial values, there were drop in pH, increase in organic carbon, CEC, total nitrogen and available phosphorus. With regard to drop in pH, this could be due to formation of organic acid during decomposition of

organic treatments and or accumulation of nitric acid from nitrification process with application of mineral fertilizer.

As stated by Lynch (1983), decrease in soil pH could be linked to accumulation of plant products which yield various organic acids. Similar observation was reported by Gattani *et al.*, (1976) in his study on the effect of continuous use of chemical fertilizers and Manures on soil properties for five years and reported that, the organic carbon level of the soil was increased to 0.24 from 0.19 percent by manure application. Mathur (1997) studied the long term effect of farm yard manure on soil properties and yield in cotton – wheat rotation at Northwest Ragasthan. The observation made was that, incorporation of 16 tonnes of farmyard manure accounted for the highest increase in organic carbon content of soil. The increase in CEC was associated with rise in organic matter content. Yaduvanshi *et al.*, (1985) also reported that, application of farmyard manure resulted in a significant increase in CEC. Patiram and Singh (1993) also found an increased in CEC through manure application and the lowest CEC of the soil was found in the treatment which received only N. Increase in total nitrogen could be attributed to direct addition of nitrogen through organic and inorganic fertilizers. As reported by Anderson and Peterson (1973) that, continuous addition of manure for 20 years increased the soil total and available nitrogen content significantly from 0.05 to 0.083 percent, while increase was only 0.05 percent in N fertilizer applied plots as large portion of N was removed by the crops. It was reported by Somani and Saxena (1975) that, an increase in phosphorus with incorporation of wheat crop residue and farmyard manures was observed while inorganic fertilizer alone decreased available phosphorus when compared with initial status. Similarly, Bharandary and Omanwar, 1994 observed that, farmyard manure treated plots showed more increase in available phosphorus than inorganic fertilizer which was due to coating of sesquioxides by organic materials that reduced phosphorus fixing capacity of the soil.

Sweet potato performance

Table 4: Effects of Cow dung, NPK and their Combinations on growth parameters of Sweet Potato at 14 WAT in UDUS and Bakalori during 2011/2012 Dry Season.

Treatment	L1 (UDUS)			L2 (BAKALORI)		
	No. of Leaves	Vine No.	Vine Length (cm)	No. of Leaves	Vine No.	Vine Length (cm)
CD(H)	197.633c	9.320c	48.433b	231.400cde	12.300cd	49.000cd
CD(M)	209.133c	9.700c	34.767c	221.000def	11.267d	38.000e
CD(L)	172.433e	9.420c	27.533d	183.100f	9.100e	53.600c
NPK(H)	329.400a	16.500a	65.000a	300.967ab	16.600a	71.533a
NPK(M)	187.500cde	11.430b	49.800b	327.300a	15.300b	64.267b
NPK(L)	180.933de	12.770b	48.733c	190.967ef	11.300d	63.000b
CD + NPK(H)	181.700de	9.330c	37.897c	212.833def	11.400d	43.600de
CD + NPK(M)	235.733b	10.700bc	51.000b	277.300bc	13.067c	62.000b
CD + NPK(L)	206.033cd	12.300b	49.500b	241.200cd	11.067d	48.300cd
Control	167.433e	9.600e	35.633c	237.000cde	9.300e	37.667e
Level of Sig.	*	*	*	*	*	*
SE	7.17	0.63	1.14	14.00	0.52	3.38

Mean followed by the same letter(s) within the same column are statistically different at 5% level of probability.

* = Significant at 5% level of probability.

Effect of cow dung, NPK and their combinations on growth parameters of sweet potato is presented in Table 4. Result indicated that, treatments had significantly different effect ($p < 0.05$) on all the considered growth parameters at both the locations. Where NPK(H) recorded the best sweet potato growth performance while the least performance in control plots. This was similarly reported by El-Ghamry (2011), who observed that, vegetative growth parameters of sweet potato tended to increase with increasing application of mineral fertilizer and different forms of organic manure. Najim et al, (2010) reported that, shoot dry matter, leaf area and plant high showed highly significant difference in relation to the rate of nitrogen fertilizer, as they increased significantly and linearly with increasing rate of nitrogen application. Hence, the result of this research could be due to high rate and availability of nitrogen in the best treatment (NPK(H)). It

216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242

was reported by many researchers (Tam and Magistad, 1935; Chapman and Barreto, 1997; Vos et-al, 1998; Janat, 2001; Abu Hussein et-al, 2003; Rashid et-al, 2005 ; and Zebarth et-al, 2009) that, increased concentration of nitrogen fertilizer can increase nitrogen uptake and this increase has a positive effect on chlorophyll concentration, the photosynthetic rates, the leaf expansion, the total

Table 5: Effects of Cow dung, NPK and their Combinations on Yield and Yield Components of Sweet Potato at UDUS and Bakaloriduring 2011/2012 Dry Season.

Treatment	L1 (UDUS)			L2 (BAKALORI)		
	No.T/plt	T.W/ plt (kg)	T yield (t/ha)	No.T/plt	T.W /plt (kg)	T yield (t /ha)
CD(H)	7.100abc	0.831bc	14.433c	5.917a	0.830b	18.733b
CD(M)	6.150abc	0.433de	8.280ef	5.750a	0.400de	8.580d
CD(L)	6.150abc	0.594d	10.727de	5.850a	0.332de	6.527d
NPK(H)	7.167abc	0.793c	15.263c	6.417a	1.173a	25.900a
NPK(M)	8.000a	1.353a	27.300a	6.583a	0.937b	22.733a
NPK(L)	6.200abc	0.483de	12.700cd	5.333a	0.543cd	12.533c
CD+NPK(H)	5.967bc	0.910bc	21.133b	5.000a	0.440d	12.533c
CD+NPK(M)	7.600ab	0.953b	19.433b	6.250a	0.630c	14.900c
CD+NPK(L)	5.533c	0.370e	9.633ef	5.000b	0.358de	9.267cd
Control	6.300abc	0.543d	8.050f	3.833b	0.250e	7.933d
Level of Sig.	*	*	*	*	*	*
SE	0.58	0.03	0.64	0.05	0.05	1.05

Mean followed by the same letter(s) within the same column are statistically the same at 5% level of significant.

* = Significant at 5% level of probability

T = tuber, T.W = tuber weight

243

244

245 Effect of cowdung, NPK and their combinations on yield and yield component of sweet potato is
246 presented in Table 5. Result indicated that, treatments had significantly different effect ($p < 0.05$)
247 on the yield and yield component at both the locations. Where NPK(M) recorded the highest
248 tuber yield at both location while control plots recorded the least tuber yield. The findings of this
249 research was similarly reported by Magagula (2010) who reported significant effect on the total
250 tuber yield of sweet potato with application of different rates of Chicken manure. This could be
251 due to an increase in nutrient level.

252 Conclusion

253 This research revealed that, application of different rates of cowdung, NPK their combination to
254 soil under sweet potato production have no significant effect on soil pH, organic carbon, CEC,
255 total nitrogen and available phosphorus. While performance of sweet potato was significantly
256 influenced by rates of cowdung, NPK and their combination and NPK(M) gave the best sweet
257 potato yield and therefore, recommended.

258

259 Reference

260 Anderson, F.N and G.A Peterson (1973). Effective continuous corn (zea mays L) manuring and N
261 fertilization on yield and protein content of the grain and on the soil N
262 content. *Agronomy Journal*, 65:697 – 700.

263 Abou–Hussein, S.D., T.E.L. Shorbagy, A.F. Abou-Hadid, and U.El -Behairy (2003). Effect of cattle
264 and chicken manure with or without mineral fertilizers on tuber quality and yield of
265 potato crop. *ISHS Acta Horticulture* 698:95-100.

266

267

268 Bharandary, V. and P.K Omanwar (1994). Long–term effect of continuous rational cropping
269 and fertilization on crop yield and soil properties – 11. Effect on EC, pH, organic matter and
270 available nutrient of soil. *Journal of Indian Society of Soil Science*, 42:387 – 392.

271

272 Chapman, S.C., and H.J Barreto (1997). Using a chlorophyll meter to estimate specific leaf
273 nitrogen of tropical maize during vegetative growth. *Agronomy Journal* 89:557-562.

- 274 Chen, J. (2006). The Combined use of chemical and organic fertilizer and/or biofertilizer for
275 crop growth and soil fertility. International workshop on sustained management of the soil
276 rhizosphere system for efficient crop production and fertilizer use. 16-20 October 2006. Level
277 Development Department Bangkok Thailand.
- 278
- 279 El-Glamry, A.M (2011). Soil fertility and potato production as affected by conventional and
280 organic systems. *Journal of Soil Science and Agricultural Engineering*, 2(2):141-156.
- 281 Gattani, P.D., S.V. Jain and S.P Seth (1976). Effect of continuous use of chemical properties.
282 *Journal of the Indian Society of Soil Science*, 24:284-289
- 283 Janat, M. (2001). Efficiency of nitrogen fertilizer for potato under fertigation utilization a
284 nitrogen traler techniques. *Communication in Soil Science and Plant Analyses*, 38:2401-2422.
- 285 Lynch, J.M. (1983). *Soil Biotechnology. Microbiology factor in crop protection*. London, U.K
286 Blackwell 981pp.
- 287 Magagula, N.E.M., E.M. Ossom, R.L. Rhykerd and C.L. Rhykerd (2010). Effect of chicken
288 manure on soil properties under sweet potato (*Ipomoea batatas* (L) Lam) culture in Swaziland.
289 *American – Eurasian journal of Agronomy*, 3(2):36-43.
- 290 Mathur, G.M. (1997). Effect of long term application of fertilizer ad manure on soil properties
291 and yield under cotton wheat rotation in North-West Rayasthan. *Journal of the Indian Society of*
292 *Soil Science*, 45:288 – 292.
- 293 Najim, A.A, M. R. Hay SeyedHadi, F. Fazali, M. TaghiDarzi and R. Shamorady (2010). Effect of
294 utilization of organic and inorganic nitrogen source on the potato shoots, dry matter, leaf area
295 index and plant height, during middle stage of growth. *International Journal of Agriculture and*
296 *Biological Science* 1:1
- 297 Nwauzor, E.C, J. K. U. Emehute, E. A. O. Okorochoa, A. O. Njoku, J. Afuape and D. Korieocha
298 (2010). National Root Crops and Research Institute, Umudike. <http://www.nreri.org>.
- 299 Rashid, M.Y. P. Voroney, and G. Parkin (2005). Predicting nitrogen fertilizers requirement for
300 corbbychrophyllmetter under different N availability condition. *Canadian Journal of Soil*
301 *Science*, 85:149-159.
- 302 Somani, L.L and S.N Saxena (1975). Effect of some organic matter sources on nutrient availability
303 in humus buildup, soil physical properties and wheat yield under field condition. *Analys of*
304 *Arid Zone*, 14:149-158.
- 305 Tam, R.K., and O.C Magustad (1935). Relationship between nitrogen fertilization and chlorophyll
306 content in pineapple leaves, *Plant physiology*, 10:159-168.

307 Vos.J., and P.E.L. Van Der Putten (1998). Effect of Nitrogen supply on leaf growth, leaf nitrogen
308 economy and photosynthetic capacity of potato.*Field Crops Research*, 59:63-72

309 Yaduvanshi, H.S. , B. R. Tripathi and B. S Kanwar (1985). Effect of continuous manuring on
310 soil properties of an alfisol .*Journal of the Indian Society Of Soil Science*, 33: 700 – 703

311

312 Zebarth , B.J. and C.J Rosen (2009). Research perspective on nitrogen BMP development for
313 potato.*American Journal of Potato Research*. 84:3-18

314