

Original Research Article

Callus Induction and Plant Regeneration from Shoot Tips of Ginger (*Zingiberofficinale*Rosc.)

Abstract

A protocol was described to regenerate plants from callus culture of ginger (*Zingiberofficinale*Rosc.). Callus cultures were induced from ginger shoot tips inverted cultured on semisolid MS medium supplemented with 2,4-D (0.5-1.0 mg L⁻¹) and/or BA (0.5- 1.0 mg L⁻¹). Maximum callus induction was obtained on MS medium supplemented with BA either alone or with 2,4-D, while the highest callus weight was on MS medium supplemented with BA (0.5 mg L⁻¹) and 2,4-D (0.5 mg L⁻¹). Induced callus was tested for regeneration on MS medium supplemented with different concentrations of 2,4-D and BA. The best plant regeneration was observed on a medium containing 2.5 mg L⁻¹ BA. The regenerated plants were rooted and successfully established in the field after few days of acclimatization.

KEYWORDS: Ginger, Callus induction, Plant regeneration, Shoot tips.

Introduction

*Zingiberofficinale*Rosc. (Ginger) of the family *Zingiberaceae* is an important tropical horticultural plant, values all over the world as an important spices for its medicinal properties. The *Zingiberaceae* is an herbaceous moderate sized family of relatively advanced monocotyledonous plant of the order Zingiberales. Zingiberaceous plants are rhizomatous, perennial and aromatic herbs often of large size, bearing flowers either terminally on aerial leaf shoots or from ground level (Kambaska and Santilata 2009). Ginger's pungent aromatic rhizome is consumed all over the world as a spice, culinary herb, condiment, home remedy, and medicinal agent (Guan *et al.*, 2010). It is vegetatively propagated through underground rhizomes, with a very low multiplication rate (Sharma and Singh 1997). The common problems in increasing ginger production are soil-borne disease infection and degeneration of rhizomes (Guan *et al.*, 2010), and the breeding of ginger has been severely hampered by poor flowering and seed set (Kackar *et al.*, 1993), and even the rhizome cannot be stored for long time as it is susceptible to fungal diseases, which affect the quality of tubers (Jamil *et al.*, 2007). ~~Therefore, other propagation techniques should be conducted to overcome such problems.~~ *In vitro* culture techniques, especially meristem culture, provide an alternative way of plant propagation and maintenance of disease free germplasm banks (Routet *et al.*, 2001). There are ~~few~~ many reports on ginger tissue culture. ~~The first was by~~ Hoski and Sagawa (1977) ~~who~~ reported ~~that~~ maximum six shoots per bud from *in vitro* culture, with low survival under field conditions. Organ culture (Sharma and Singh 1997), somatic embryogenesis (Kackar *et al.*, 1993) and (Lincy *et al.* 2009), organogenesis (Babuet *et al.*, 1992), protoplast culture (Geetha *et al.*, 2000), microrhizom production (Zheng *et al.*, 2008), shoot tips culture (Ayenew *et al.*, 2012), rhizome buds cultured at different size (Sathyagowri and Thayamini 2011) and germplasm preservation (Dekker *et al.*, 1991) have been conducted. Ginger seed production also was suggested by Uozumiet *et al.*, (1994) by *in vitro* propagation

Comment [DU1]: Indicate the concentrations exhibited the highest percentage callus formed

Comment [DU2]: Indicate the concentration?

UNDER PEER REVIEW

methods, ~~but~~ however these methods are not efficient, expensive and do not meet ginger germplasm conservation requirements.

The present work aimed to describe an efficient protocol for; calli production; shoot ~~lets~~ regeneration, keeping in view the potential action against various ailments and its regeneration problems in natural environment. Micropropagation of this medicinal important plant was accomplished to provide stable supply of the *in vitro* raised plantlets for the mass production for this important medicinal plant in Iraqi Kurdistan Region and to become a powerful tool for future studies.

Materials and Methods

Callus Induction

This investigation was conducted at Plant Tissue Culture Lab at the Scientific Research Center, Faculty of Science at the University of Duhok. Shoot tips of *in vitro* cultured ginger on MS medium supplemented with 1.0 mg/l BA were cut and placed in inverted form on the culture medium. Various concentrations of 2,4-D and BA (0.0, 0.5, 1.0 mg /l) in MS medium containing 3% sucrose, 100 mg/l inositol were evaluated for inducing callus. The pH of the medium was adjusted to 5.7 ± 0.1 with 1N NaOH or HCl, prior to the addition of agar (7.0) g/l. The medium was brought up to the final volume, ~~then subsequently, 25 ml of medium was~~ dispensed ~~at 25 ml rates,~~ into 250 ml Mason jars and capped with colorless PVP covers and fitted with rubber bands. The medium was ~~then~~ sterilized by autoclaving ~~for~~ 15 min. under 1.04 kg/cm² and allowed to solidify under room temperature.

Comment [DU3]: Indicate the temperature used

Plant Regeneration

Two months old callus cultures were transferred to MS medium supplemented with BA (0.0, 2.5, and 5 mg/l) and 2,4-D (0.0, 0.2 and 0.4 mg/l) either alone or in combination. After eight weeks of culture the shoots length, number of shoots and leaves/ explant were recorded. The plantlets were separated from the callus and cultured on MS medium supplemented with 2.5 mg/L kinetin for both multiplication and rooting stages (Tomaet *et al.*, 2012).

Eventually, for hardening stage, a number of well and successfully rooted plantlets were removed from culture vessels and their roots were washed with distilled water and immersed in Benlate fungicide (0.1% for 10 minutes). Finally, they were planted in pots containing steam sterilized soil mixture (Peatmoss+ loam+ Styrofoam

1:1:0.5, v: v: v) under tightly controlled atmosphere of the greenhouse.

Data Analysis

Three explants were cultured in each vessel with five replications. The cultures were incubated in the culture room under $25 \pm 1^\circ\text{C}$ temperature and 16 hours daily exposure to 1000 Lux cool white light, followed by 8 hours of darkness. The results were recorded after 8 weeks of culture.

All the comparison between means was carried out according to Duncan's multiple range test ($P < 0.05$) using a computerized program of SAS (SAS, 2001).

Results and Discussion

After 8-10 days of culture, there was a notable enlargement of inverted shoot tips with a whitish to brown [color](#) and fragile texture (Figure 6, A). The callus growth was slow and highly sensitive to mechanical force or fragile, the surface was covered with small hairy structure and the contamination rate was rare.

The caulogenic response of the shoot tips to various concentrations and combinations of cytokinins and auxins is shown in (Figure 1, Figure 2). For callus induction the highest and best callus induction (100%) was noticed on MS medium containing BA either alone or in combination with 2,4-D (Fig. 6, B), while those devoid of BA showed lowest response. Similar kind of results were obtained by (Malamu *et al.*, 1991), when they observed that BA and 2,4-D combinations were best for callus induction. (Babu *et al.*, 1992) emphasized that 2, 4-D is the best auxin for callus induction in ginger and turmeric (Jala, 2013).




Figure (12): Effects of different concentrations of BA and 2, 4-D on callus formation from shoot tips of *Zingiber officinale*.

The highest callus weight (1.43 g) showed in (Figure 23) observed on MS medium supplemented with 0.5 mg/l from BA and 2,4-D (Fig. 6, C) followed by MS medium containing only BA at concentration 0.5 mg/l (1.34 g), but the statistical analysis of the experimental data revealed that there is no significance differences between them.

While the least weight (0.17 g) was observed on MS medium devoid of growth regulators.




Figure (23): Effects of different concentrations of BA and 2, 4-D on callus weight from shoot tips of *Zingiberofficinale*.

The well developed, white and fragile callus was transferred to regeneration medium (Figure 34). The highest mean number of regenerated shoots (6.0 shoots/explants) was observed on MS media containing BA 2.5 mg/l (Fig. 61, D) followed by MS medium supplemented with BA 5.0 mg/l which was 5.0 shoots/explant, but they were not different significantly. Similar results were recorded by (Jala, 2013) they observed that 5.0 mg/l BA produced the highest number of shoots form callus in *Curcuma attenuate* and, (Yapinget *al.*, 2013) recorded that BA combination with NAA and TDZ was important for anther callus regeneration in *Curcuma attenuate*, while the lowest mean number of shoots was 0.67 shoots/ explant on MS medium devoid of growth regulators. The MS medium fortified with BA and 2,4-D showed less mean numbers of shoots. Similar results were recorded by Malamuget *al.* (1991). They observed that 1-3 mg/l of BA was the best concentration for regeneration from callus tissues, while (Babuget *al.*, 1992) confirmed that 2, 4-D enhances organogenesis when added in small concentrations. Since there is no significant difference between 2.5 and 5.0 mg/l of BA, one can use the lower concentration of BA (2.5 mg/l) in order to lessen the mass production cost.





Figure (34): Effects of different concentrations of BA and 2, 4-D in No. of shoot \ explant from callus culture of *Zingiber officinale*.

These results emphasize the importance of cytokinins in shoot regeneration in *in vitro* plant propagation. Cytokinins activate RNA synthesis; stimulate protein synthesis and the activities of some enzymes. Generally, herbaceous plants are highly responsive to BA treatments and most herbaceous species produce well formed shoots suitable for further shoot proliferation (Debergh and Zimmerman, 1991).

On the other hand, the highest mean of shoot length (6.00 cm) (Figure 45) and highest mean of leaves number (5.0 leaves/ explant) (Figure 56) was found in a medium containing 0.2 mg/l 2,4-D only without adding any cytokinins and this was highly significant than all other means.






Figure (45): Effects of different concentrations of BA and 2, 4-D in shoot length (cm) from callus culture of *Zingiber officinale*.




Figure (56): Effects of different concentrations of BA and 2, 4-D in No. of leaves\ shoots from callus culture of *Zingiber officinale*.

Then the callus regenerated explants transferred to multiplication medium containing 2.5 mg/l kinetin and the plantlets rooted in the same medium. For rooting stage, the plantlets rooted without auxins in the medium; these records are in agreement with those found by Tomaet *al.* (2012) and Kavyashree (2009) which indicates that the endogenous auxin level in the plant itself is so sufficient that can form roots without applying any exogenous auxins. Plantlets were successfully transplanted into pots

UNDER PEER REVIEW

after sufficient development of roots to establish the field condition and rhizome formation in the soil (Fig. 61, E, F).




Figure-61 : A. shoot tip culture in inverted in MS medium, B. Callus induction in MS medium supplemented with BA and 2,4-D after 8 weeks, C. High callus weight in MS medium supplemented with 0.5 mg/l BA and 2,4-D , D. plantlets regenerated from the callus , E Acclimatization of plantlets in the soil, F. Hardened plant ready for transfer to field with formed rhizome .

Comment [DU4]: Should be considered as figure 1

References

UNDER PEER REVIEW

- Ayenew, B.; Tefera, W. and Kassahun, B. (2012). *In vitro* propagation of Ethiopian Ginger (*Zingiberofficinale*Rosc.) cultivars: evaluation of explants types and hormone combinations. Afr. J. Biotechnol., 11(16): 3911-3918.
- Babu, K.N.; Samsudeen, K. and Ravindran, P.N. (1992). Direct regeneration of plantlets from immature inflorescences of ginger (*Zingiberofficinale*Rosc.) by tissue culture. J. Spices Aromat. Crops, 1: 43–48.
- Dekker, A.J.; Rao, A.N. and Gob, C.J. (1991). *In vitro* storage of multiple shoot cultures of ginger at ambient temperatures of 24–29 C°. Sci. Hort., 47:157–167.
- Geetha, S.P.; Babu, K.N. and Rema, J. (2000). Isolation of protoplasts from cardamom (*Elettariacardamomum*Maton.) and ginger (*Zingiberofficinale*Rosc.). J. Spices Aromat. Crops, 9:23–30.
- Guan, Q.; Guo, Y.; Wei, Y.; Meng, F. and Zhang, Z. (2010). Regeneration of somatic hybrids of ginger via chemical protoplast fusion. Plant Cell Tiss. Organ Cult., 102:279–284.
- Hosoki, T. and Sagawa, Y. (1977). Clonal propagation of ginger (*Zingiberofficinale*Rosc.) through tissue culture. Hort Science, 12:451–452.
- Jamil, M.; Kim, J.K.; Akram, Z.; Ajmal, S. U. and Rha, E. S. (2007). Regeneration of ginger plant from callus culture through organogenesis and effect of CO₂ enrichment on the differentiation of regenerated plant. Biotechnology, 6(1): 101-104.
- Jala, A. (2013). The effect of the 2,4-Dichlorophenoxy acetic acid, benzyl adenine and paclobutrazol, on vegetative tissue derived somatic embryogenesis in turmeric (*Curcuma*VarChattip). Int. Trans. J. Eng. Manag. Sci. Tech., 4(2):105-110.
- Kackar, A.; Bhat, S.R.; Chandel, K.P.S. and Malik, S.K. (1993). Plant regeneration via somatic embryogenesis in ginger. Plant Cell, Tissue and Organ Culture, 32: 289-292.
- Kambaska, K.B. and Santilata, S. (2009). Effect of plant growth regulator on micropropagation of ginger (*Zingiberofficinale*Rosc.) cv- Suprava and Suruchi. Journal of Agricultural Technology , 5(2): 271-280.
- Kavyashree, R. (2009). An efficient *in vitro* protocol for clonal multiplication of ginger - var. Varada. Indian Journal Biotechnology, 8:328-331.

UNDER PEER REVIEW

Malamug, J. F.; Inden, A. and Asahira, T. (1991). Plantlet regeneration and propagation from ginger callus. *Sci. Hortic* , 48: 89 – 97.

Murashige, T. and Skoog, F. (1962). A revised medium for rapid growth and bioassay with tobacco tissue cultures. *Physiol. Plant*,15: 473-497.

Routs, G. R.; Palai, S. K.; Samantaray, S. and Das, P. (2001). Effect of growth regulator and culture conditions on shoot multiplication and rhizome formation in ginger (*Zingiberofficinale*Rosc.) *in vitro*. *In Vitro Cell. Dev. Biol.-Plant*, 37:814-819.

Sharma, T. R. and Singh, B. M. (1997). High-frequency *in vitro* multiplication of disease-free *Zingiberofficinale*Rosc. *Plant Cell Reports*, 17: 68–72.

SAS. (2001). SAS/ STAT User's Guide for Personal Computers. Release 6.12. SAS Institute Inc. Cary, NC, USA.

Sathyagowri, S. and Thayamini, H. (2011). *In vitro* plant regeneration of ginger (*Zingiberofficinale*Rosci.) with emphasis of initial culture establishment. *Int. J. Med.Arom. Plants*,1(3):195-202.

Toma, S. R.; Danial, G. H. and Rashad, R. H. (2012). *In vitro* propagation of ginger (*Zingiberofficinale*Rosc.) from a single bud explant as affected by kinetin and benzyl adenine. *Journal of Duhok University*, 15 (1): 423-427.

Uozumi, M.; Asamo, Y. and Kobayashi, T.K. (1994). Micropropagation of horseradish hairy root by means of adventitious shoot primordia. *Plant Cell Tissue Org. Cult.* 36:183–190.

Yaping, K.; Guohua, M.; Jaime, A. and Teixeira, S. (2013). Callus induction and shoot organogenesis from anther cultures of *Curcuma attenuate* Wall. *Plant Cell Tiss. Organ Cult.*, 112:1-7.

Zheng, Y.; Liu, Y. Ma, M. and Xu, K. (2008). Increasing *in vitro* microrhizome production of ginger (*Zingiberofficinale* Roscoe). *ActaPhysiol Plant.* 30:513–519.