

Original Research Article

Estimation of Nutritional and Starch Characteristics of *Dioscorea alata* (Water Yam) Varieties Commonly Cultivated in the South-Eastern Nigeria.

Abstract:

Background: Studies on the nutritional and mineral compositions of staple foods will provide evidence to regulate intake of certain common foods for optimal health condition.

Aim: To estimate the proximate, mineral and starch characteristics of five local varieties, *D. purpurea*, *D. atropurpurea*, *D. liliopsida* (purple yam), *D. vilgaris*, and *D. villosa* of *Dioscorea alata* grown in Izzi Area of Ebonyi State, Nigeria.

Methodology: Five (5) commonly cultivated varieties of *D. alata* were sampled, each variety in triplicates, and prepared for analysis by peeling, dicing, drying and milling into flour. The proximate, nutritional and starch characteristics were determined by standard methods. Minerals were quantified by Atomic Absorption Spectrophotometric Method.

Results: The results showed that the varieties has moisture content 9.20-10.30%, ash content 2.48-3.53%, fiber content 3.31-3.53%, fat content 1.62-2.41%, protein content 8.40-10.46%, and carbohydrate content 70.88-73.90%. There were significant differences ($P<.05$) in the proximate compositions within the varieties. The ranges of minerals in mg per kg (dry weight) were Na 16.38-24.84, K 97.78-141.14, Ca 79.99-269.75, Mg 18.55-31.53, P 114.65-211.63, Fe 15.18-30.86. Glycaemic index ranged from 35.56-41.31, amylose content 12.42-16.11% and gelatinization temperature 85.00-87.00°C

Conclusion: The results indicate that the protein and fiber contents of *D. alata* varieties estimated in this study were high. The low glycemic index confirmed the yam varieties as low glycemic index food. It was concluded that the yam varieties are good sources of protein nutrient and suitable staple food for the diabetics. However, low mineral compositions were reported. Therefore, there is a need for intense cultivation of improved varieties for improved mineral intake from the yam by the consumers.

Key words: *Dioscorea alata*, proximate composition, mineral composition, nutritional quality, starch

34 1. Introduction

35 Nutritional quality and mineral composition of primary agricultural and cultural foods in
36 developing countries is a major health problem which has manifested as diet-associated
37 disorders among the African population. Food and nutrition are known important modifiers of
38 disease initiation and development. The major causes of illness and death appear to be the
39 chronic degenerative diseases, such as cancer, heart disease, arthritis, respiratory diseases,
40 diabetes, hypertension, cognitive impairment and various toxic states, which could be averted
41 with proper nutrition and diet. Beside good quality of water, a well-balanced diet of essential
42 macronutrients and micronutrients is required to achieve optimal health condition [1]. In recent
43 years, there has been a tremendous interest in the field of nutrition on the assessment of
44 nutrients, vitamins and mineral composition of staple foods considered to be economically,
45 socially and culturally important in many tropic and subtropic regions of the world. In these
46 regions, yam is being investigated for its whole nutritional quality. It is traditionally known that
47 yam tubers have potential ability to provide one of the cheapest sources of dietary energy in
48 the form of carbohydrate [2, 3].

49 Yam (*Dioscorea species*) is a climbing, vigorously twining herbaceous plant that coils swiftly
50 around the stake. They are perennial through root system but are grown as annual crops [4].
51 Yams are of great economic importance and nourishment to the people of Africa, the
52 Caribbean, Asia and America [5]. They are well integrated into the social, cultural and religious
53 lifestyle of the people who cultivate and consume them [6]. Large quantity consumption has
54 been reported to provide the body not only with carbohydrate but also with vitamins and
55 minerals [7]. It is second to cassava in tropical root crops but first in higher vitamin C content
56 (40-120mg/g edible portion) and crude protein content (40-140g/kg dry matter) [8]. Yam
57 belongs to a family of *Dioscorea* with over 600 species in which only few are cultivated for food
58 and medicines [9]. The most cultivated species in Nigeria are the white yam (*D. rotundata*),
59 yellow yam (*D. cayenensis*), water yam (*D. alata*) and trifoliate yam (*D. dumentorum*) [10].

60 *Dioscorea alata*, commonly called water yam or ten month yam is the most widespread yam
61 species and more important as food in West Africa and the Caribbean than in Asia and in
62 America where it originated and has been competing with the most important native species,
63 *D. rotundata*. *D. alata* is an excellent source of starch, which provides calorific energy and
64 protein three times more superior than the one of cassava and sweet potato [11,12]. It has low
65 sugar content necessary for diabetic patients [13]. Cultivars of *D. alata* possess higher contents
66 of protein, vitamin C and low lipids than *D. cayenensis*, *D. rotundata* and *D. trifida* [14]. It has
67 comparative advantage for sustainable production due to its better agronomic characteristics
68 such as ease of propagation and yield, high nutritive value and longer storage of the fresh
69 tubers [5]. It is considered to be among the most primitive of the angiosperms and contains

many species of which only about few are considered edible. *D. alata* is popular and prevalent within Abakaliki agro-ecological zone of Ebonyi State where it is called “ Mbala or Nvula” (Igbo names) [4]. Cultivated varieties of *Dioscorea alata* in this area includes, *D. purpurea* (Okwale nkata), *D. atropurpurea* (Nvula ite), *D. liliopsida* (Nvula eke) or purple yam, *D. vilgaris* (Nvula ji), *D. villosa* (Akpuru akputo), *D. harmiltonii*, *D. persimilis* etc [9].

The objective of this study was to determine the nutritional and starch characteristics of five major varieties of *D. alata* commonly cultivated by the Igbo people of Ebonyi State. Results from the study will provide scientific evidence for the nutritional quality of the varieties, and possibly the need for improved varieties for cultivation in the area.

2. Materials and Methods

2.1 Sample Collection

Five (5) variety samples of *D. alata*, sample A (*D. purpurea*), sample B (*D. atropurpurea*), sample C (*D. liliopsida*), sample D (*D. vilgaris*) and sample E (*D. villosa*) were obtained from different parts of Izzi Local Government Area of Ebonyi State, Nigeria. The yam tubers were carefully selected to avoid rot, bruises or sign of spoilage.

2.2 Sample Preparation for Analysis

The yam tubers were cut into two, thoroughly washed, peeled and cut into small pieces (cubes). It was sun-dried for 3 to 4 days, packaged and labeled sample A-E. Each sample variety was prepared in triplicates. The dried pieces were milled into powdered samples and sieved through 250µm mesh and stored in plastic container for nutritional and mineral analysis.

2.3 Determination of Proximate Compositions and Starch Characteristics

The moisture and crude fibre content of each variety were determined by drying the sample in an oven at 105⁰C for 6hrs and values expressed in percentage. The ash content was determined by incinerating in a muffle furnace 2g of each sample at 550⁰C for 2hrs [15]. Protein content, expressed in percentage, was obtained using Kjeldhal method ($N \times 6.25$) by AOAC [15]. Crude fat content was estimated by Soxhlet extraction technique using petroleum ether (40-50⁰C) (16). The carbohydrate content was determined by difference (% carbohydrate = 100% - sum of percentages of moisture, fat, ash, fibre and crude protein contents). Iodo-colorimetric method [17] based on the chemical reaction of starch and iodine was used to determine amylose content. Glycemic Index was calculated from the hydrolysis index expressed in percentage (area under curve procedure) as described by Hettiaratchi et al., [18]. To determine the gelatinization temperature, 10% suspension of the flour sample was prepared in a test tube. The aqueous suspension was heated in a boiling water bath, with continuous stirring. Gelatinization was observed after 30s and measured accurately as the gelatinization temperature.

2.4 Determination of Mineral Content

Sodium, potassium, calcium, magnesium, phosphorus and iron were analysed after digestion with HNO_3 and H_2SO_4 , using Atomic Absorption Spectrophotometer (Model Buck 2006, Buck Scientific, USA). Phosphorus content was determined with the use of phosphomolybdate complex method.

2.5 Statistical Analyses

All measurements were carried out in triplicates and the mean triplicate values of the five varieties were subjected to One-Way ANOVA Duncan's t-test. Differences considered significant at 5% ($P < .05$).

3. Results and Discussion

Table 1: Proximate Composition of *Dioscorea alata* varieties (% dry weight) *

Proximate Composition	Sample Varieties of <i>Dioscorea alata</i> (water yam)				
	A <i>D. purpurea</i>	B <i>D. atropurpurea</i>	C <i>D. liliopsida</i>	D <i>D. vilgaris</i>	E <i>D. villosa</i>
Moisture content	9.90 ± 0.14 ^b	9.20 ± 0.00 ^d	9.90 ± 0.14 ^b	10.30 ± 0.14 ^a	9.5 ± 0.14 ^c
Ash content	3.40 ± 0.14 ^a	3.45 ± 0.07 ^a	3.53 ± 0.04 ^c	2.48 ± 0.04 ^c	3.14 ± 0.04 ^b
Crude fibre content	3.38 ± 0.03 ^b	3.50 ± 0.00 ^a	3.53 ± 0.01 ^a	3.31 ± 0.01 ^b	3.33 ± 0.04 ^b
Fat content	1.99 ± 0.01 ^d	2.15 ± 0.07 ^c	2.39 ± 0.01 ^b	1.62 ± 0.03 ^e	2.41 ± 0.01 ^a
Protein content	10.46 ± 0.06 ^a	10.15 ± 0.13 ^b	8.71 ± 0.06 ^c	8.40 ± 0.13 ^d	10.02 ± 0.06 ^b
Carbohydrate content	70.88 ± 0.78 ^c	71.55 ± 0.27 ^b	71.95 ± 0.24 ^b	73.90 ± 0.04 ^a	71.57 ± 0.15 ^b

* Values are Mean ± SD of triplicate determinations; means along a row with the same superscript are not significantly different, $P < .05$

Table 2: Starch Characteristics of *Dioscorea alata* varieties

Starch Characteristics	Sample Varieties of <i>Dioscorea alata</i> (water yam)				
	A <i>D. purpurea</i>	B <i>D. atropurpurea</i>	C <i>D. liliopsida</i>	D <i>D. vilgaris</i>	E <i>D. villosa</i>
Glycemic index(%)	35.56 ± 0.64 ^a	36.47 ± 1.09 ^a	38.90 ± 1.09 ^a	40.88 ± 0.45 ^a	41.31 ± 1.77 ^a
Amylose	12.42 ± 0.03 ^c	15.22 ± 0.02 ^a	13.68 ± 0.60 ^b	16.11 ± 0.56 ^a	13.99 ± 0.13 ^b

content (%)

Gelatinization 85.00± 1.40^a 85.00 ± 0. 00^a 85.50 ± 2.10^a 87.00 ± 1.40^a 87.00± 1.40^a
temp. (°C)

Values are Mean±SD of triplicate determinations; means along a row with the same superscript are not significantly different, $P < .05$

Table 3: Mineral Composition of *Dioscorea alata* Varieties (mg/kg)*

Minerals	Sample Varieties of <i>Dioscorea alata</i> (Water yam)				
	A	B	C	D	E
	<i>D. purpurea</i>	<i>D.atropurpurea</i>	<i>D. liliopsida</i>	<i>D. vilgaris</i>	<i>D. villosa</i>
Sodium	18.38 ± 0.18 ^c	16.38 ± 0.46 ^d	16.99 ± 0.02 ^d	24.84 ± 0.37 ^a	21.06 ± 0.77 ^b
Potassium	134.68±1.38 ^b	141.14± 1.35 ^a	127.04± 2.74 ^c	111.30±1.06 ^d	97.78 ± 2.58 ^e
Calcium	79.99 ± 0.15 ^e	269.75± 1.77 ^a	168.09± 2.74 ^c	196.90±0.28 ^b	140.07±0.41 ^d
Magnesium	25.58 ± 0.67 ^b	29.77 ± 0.38 ^a	21.04 ±0.06 ^c	18.55 ± 1.56 ^d	31.53 ± 0.76 ^a
Phosphorus	114.65±0.43 ^d	211.63± 2.02 ^a	117.10±0.64 ^b	205.10±7.78 ^a	169.76±1.05 ^c
Iron	20.61 ± 0.09 ^b	19.25± 0.64 ^c	16.86 ±0.23 ^d	30.86± 0.29 ^a	15.18 ± 0.61 ^e

* Values are Mean±SD of triplicate determinations; means along a row with the same superscript are not significantly different, $P < .05$

Table 4: Ratios of Na to K and Ca to P of *Dioscorea alata* Varieties

Ratio	Sample Varieties of <i>Dioscorea alata</i> (Water yam)				
	A	B	C	D	E
	<i>D. purpurea</i>	<i>D.atropurpurea</i>	<i>D. liliopsida</i>	<i>D. vilgaris</i>	<i>D. villosa</i>
Na/K	0.14	0.12	0.13	0.22	0.22
Ca/P	0.70	1.27	1.44	0.96	0.83

3.1 Proximate Compositions of *D. alata* Varieties

The proximate composition of *D. alata* varieties is presented in Table 1. The moisture content of the local varieties ranged from 9.2% for *D. atropurpurea* to 10.30% for *D. vilgaris*, with a mean of 9.76%. The mean values for moisture content showed significant differences ($P<.05$). Moisture content is an important factor in food quality, preservation, and resistant to deterioration. It is related to many properties that affect the quality and shelf life of food and food products, including bacterial growth and texture [19]. Low moisture content of a food indicates its stability for prolonged storage or permissive rate of spoilage. In this study, local varieties of *D. alata* have lower moisture content than those reported in the literature [5,8,21,22]. Comparable to our result, Udensi et al, [4] have reported moisture content of 6.05% for local best variety of *D. alata*. In the same study, five high yielding and disease

resistant water yam varieties gave low moisture content. The moisture content of sweet potato, 8.72% and red cocoyam, 9.02% corroborated the moisture content of *D. alata* varieties in this study [20]. These local varieties of *D. alata* would be suitable for prolonged storage and more stable for industrial food product processing.

The crude fiber content ranged between 3.31 to 3.53% with a mean value of 3.41%. There was no significant difference ($P>.05$) between *D. atropurpurea* and *D. liliopsida* fibre contents, however, significant difference ($P<.05$) was observed between these varieties and others. Fibers in foods are plant non-digestible carbohydrate and lignin and have been shown to have health benefits. Studies have shown that increase in fiber consumption in foods reduces the incidence of obesity, cardiovascular disease, type 2 diabetes, digestive disorders and some cancers [23]. Fiber consumption also softens stools and lowers plasma cholesterol level in the body [24]. The mean fiber content in the current study is comparable to the 3.96% reported by Shajeela et al, [25]. Lower values have been reported for other varieties of *D. alata* [4], *D. rotundata* (white yam) and *Colocasia esculenta* (white cocoyam) [26]. The result indicates that the local varieties of *D. alata* investigated could be a source of dietary fiber with nutritional benefits.

The range of crude protein content (8.40-10.46%) was in agreement with the earlier studies [4,5,21,25,27]. Strikingly, a study by Alinnor and Akalezi, [26] reported crude protein content of 0.087% for *D. rotundata*, a widely cultivated and the most preferred specie of yam in Nigeria and West Africa. In earlier study carried out by Oyenuga [28], he found the protein content of *D. alata* to be in the range of 7.26-8.10 while that of *D. rotundata* 4.42-5.87. This suggests a higher nutritional potential for *D.alata*, at least in terms of protein content than *D. rotunda*. Intake of staple foods with low protein content may lead to several impaired biological processes in the body. This shows that *D.alata* is rich in protein and can provide this nutrient to the consumers.

The highest carbohydrate content in this study was 73.90% with no significance difference ($P>.05$) within the varieties. The result is in line with the findings of studies [27, 29] which reported 76.57% and 75.65% respectively for *D.alata* but lower when compared with the carbohydrate contents reported for red cocoyam, 86.69% and sweet potato, 86.90% [29]. Improved varieties of *D.alata* contained a higher carbohydrate content ranged between 81.53 to 87.64% [4]. However, lower carbohydrate content has been reported for commonly consumed root tubers in Nigeria. Alinnor and Akalezi, [26] reported 40.61% and 57.78% carbohydrate contents for white yam (*D. rotundata*) and white cocoyam (*Colocasia esculenta*) respectively. *D.alata* varieties in this study may be considered as tolerable sources of energy and this may be related to the recommendation of water yam and some other cocoyam varieties for the diabetics.

3.2 Starch Characteristics of *D. alata* Varieties

Table 2 presents the starch characteristics of the yam varieties. The glycemic indices of *D. alata* investigated ranged from 35.56 to 41.31, with mean value of 38.62. The glycemic index (GI) concept is based on the difference in blood glucose response after ingestion of the same amount of carbohydrates from different foods, and possible implications of these differences for health [30]. Based on this, previous studies have been able to provide evidence for the classification of foods as low (≤ 55), intermediate (56 to 59) and high (≥ 70) GI foods. The glycemic index values of yam reported in literature ranged from low to high and generally the species and varieties were unspecified [30,36]. However, low GI foods have been associated with reduced risk for chronic and metabolic diseases including diabetes [31-34]. The GI phenomenon expresses the glycemic response to a food after ingestion of 50g of available carbohydrate and the possible implications on health[30]. Foods that raise blood glucose level quickly have high GI levels, whereas foods with low GI will raise blood glucose levels slowly [35]. The reported GI in the current study is classified as low for the local yam varieties specified in Table 2. A study reported a GI of 67 for white yam (*D. rotundata*), 56 for three-leaved yam (*D. domentorum*) and 24 for water yam (*D. alata*) [21]. Among the Chinese starch foods, a GI of 52 was obtained for yam, although the specie unspecified [37]. It is noteworthy that the *D. alata* varieties investigated in the current study are low-GI foods. Inconsistent in GI for yam may be due largely to different species. The yam varieties may be considered in diet and more importantly among the diabetics who are concerned about their postprandial blood glucose levels for effective diet management.

The amylose content of the varieties varied significantly ($P<.05$) and *D. purpurea* was found to have the lowest amylose content ($12.42\pm0.03\%$) while *D. vilgaris* had the highest ($16.11\pm0.56\%$). A similar result was obtained in a study that demonstrated inverse correlation between amylose content and swelling power of different food starches, reporting 12.24% amylose for *Dioscorea esculenta* [38]. However, previous studies have reported comparatively higher values for amylose content in *D. alata*, although varieties investigated were not specified [39,40]. Riley et al, [41] analysed eight *D. alata* cultivars grown in Jamaica and reported amylose content that ranged from 20.117 to 23.001%. The difference in amylose content observed may be attributed to genetic variations among the varieties, as it was found that the amylose content of starches is affected by the expression of the amylose extender gene [42]. The gelatinization temperature range of starch in the varieties was 85.00 to 87.00°C. The gelatinization temperature obtained was considerably higher than for wheat starch (55.6 to 63.0°C) according to Lineback and Ke (43). This high temperature suggests slow swelling property of the starch granules which may be connected with protein-amylose formation [41]. In biotechnological application of foods for value-added products, starch with lower gelatinization temperature to minimize energy cost and for shorter cooking time may be more preferred.

3.3 Mineral Compositions of *D. alata* Varieties

Minerals are biological components of diets which perform biochemical and physiological functions in living cells through synergistic interactions and/or independent modulation of biological reactions. The mineral levels of the *D. alata* varieties are shown in Table 3. The low mineral levels indicate that these varieties may not be an important source of minerals to the consumers. This may be due to the species and the prevailing environmental factors. Considering the specialized functions of minerals in the human body, which include maintenance of acid-base balance, neurotransmission, cofactors of enzymes, bone and blood formation, energy transduction among others, there is need for improved varieties of *D.alata* for cultivation. The sodium to potassium ratio in the body is of great concern for the prevention of hypertension [26,44]. According to the World Health Organization [45], the sodium-to-potassium ratio should be $\leq$ one. The consumption of the varieties regarding sodium-to-potassium ratio would not promote high blood pressure. Food is considered “good” if Ca/P ratio is above one and “poor” if the ratio is less than 0.5, while the Ca/P ratio above two helps to increase the absorption of calcium in the intestine [46]. The Ca/P ratios of *D. atropurpurea* and *D. liliopsida* were above the recommended value of 1.00. However, higher ratios have been reported for *Dioscorea rotundata* and *Colocasia esculenta* [26].

Conclusions

From the present study, it could be summarized that *D. alata* varieties investigated had nutritional proximate compositions with immense health benefits. The average moisture content was lower compared to the reported in the literature, an important food property for long-term food storage and industrial food processing. The varieties are good sources of protein, fiber and carbohydrate. *D. purpurea*, *D. atropurpurea* and *D. villosa* in particular could be selected for high protein level for intense cultivation in other parts of Nigeria and other regions of the world growing *D. alata*. Based on the standard criteria, *D. alata* varieties with mean glycemic index of 38.62 in this study are low GI foods. It is therefore relevant, noting the health benefits of these varieties to the consumers in prevention of chronic diseases, including type 2 diabetes mellitus. The tuber varieties had low mineral levels. There is a need for intense breeding of improved varieties for enhancing mineral quality of these varieties without losing the enviable nutritional qualities. Taken together, *D.atropurpurea* may be regarded as the best variety according to the findings in this study.

References

1. Enwonwu CO, Philips RS, Falkler WA Jr. Nutrition and Oral Infectious Diseases: State of the Science. Comp. Cont. Educ. Dent 2002; 23: 431—448
2. Behera KK, Maharana T, Sahoo S, Prusti A. Biochemical quantification of protein, fat, starch, crude fiber, ash and dry matter content in different collection of greater yam (*Dioscorea alata* L) found in Orissa. Nature and Science 2009; 7: 24-32

3. Ugwu FM. The potentials of roots and tubers as weaning foods. *Pakistan Journal of Nutrition* 2009; 8: 1701-1705
4. Udensi EA, Oselebe HO, Iweala OO. The investigation of Chemical Composition and functional properties of water yam (*Dioscorea alata*): Effect of Varietal Differences. *Pakistan Journal of Nutrition* 2008; 7(2): 342-344
5. Baah FD, Maziya-Dixon B, Ashiedu R, Oduro I, Ellis WO. Nutritional and biochemical composition of *Dioscorea alata* (*Dioscorea* spp) tubers. *J of Food, Agriculture and Environment* 2009; 7(2): 373—378
6. Hahn SK, Osiru DSO, Akoroda MO, Otoo JA. Yam production and its future prospects. *Outlook on Agriculture* 1987; 16(3): 105-110
7. Scot GJ, Rosegrant M, Ringler C. Roots and Tubers for 21st Century: Trends, Projections and Policy Options. Food and Agriculture and Environment Discussion. International Food Policy Research Institute (IFPRI) and International Potato Centre (CIP), Washington DC, USA, 2000.
8. Opera LU. Yam Storage. In: Bakker-Arkema, FW (ed), *CIGR Handbook of Agricultural Engineering Vol IV Agro Processing*. The American Society of Agricultural Engineers, St Joseph, MI, USA, 1999, pp 182-214
9. IITA. "Yam Research Review". International Institute of Tropical Agriculture, Ibadan, Nigeria, 2006.
10. Amusa NA, Adegbite AA, Muhammed S, Baiyewu RA. Yam diseases and its management in Nigeria. *African J Biotechnol* 2003; 2(12): 497-502
11. Coursey DG. Cassava as food: Toxicity and Technology. In: Nestel B and MacIntyre (eds). *Chronic Cassava Toxicity*, Ottawa, Canada, IDRC, IDRC-10e, 1973, pp 27-36
12. Bourret-Cortadellas D. Ethnobotanic study of the *Dioscorea*, Yams New Caledonia. Doctorate 3rd cycle, Faculty of Science, Paris, France, 1973, pp: 135
13. Udensi EA, Oselebe HO, Onuoha AU. Antinutritional Assessment of *D. alata* varieties. *Pak J Nutr* 2010; 9(2): 179—181
14. Muzac-Tucker I, Asemota HN, Ahmad MH. Biochemical composition and storage of Jamaica yams (*Dioscorea* sp). *J Sci. Food Agric* 1993; 62 (3): 219-224
15. AOAC. Official Methods of Analysis (18th edn). Association of Official Analytical Chemists, Washington DC, 2005
16. Pearson D, Egan H, Kirk RS, Sawyer R. *Pearson's chemical analysis of food*. Edinburgh: Churchill Livingstone, 1981.
17. Juliano BO, Perez CM, Blakeney AB, Castillo T, Kongserree N, Laignelet B, Lapis ET, et al. International testing on the amylase content of milled rice. *Starch/Starke* 1981; 33: 157-162
18. Hettiaratchi UPK, Ekanayake S, Welihinda J. Prediction of glycaemic Index (GI) of meals by starch hydrolysis indices. *Int Food Research Journal* 2012; 19(3): 1153-1159

19. Parasoglou P, Parrott EPJ, Zeitler JA, Rasburn J, Powell H, Gladden LF, Johns ML. Quantitative Water Content Measurements in Food Wafers using Terahertz Radiation. *Terahertz Science and Technology* 2010; 3(4): 172-182
20. Oladebeye AO, Oshodi AA, Oladebeye AA. *Proceedings of International Conference of Chemical Society of Nigeria*, 2008a.
21. Ihediohanma NC, Onuegbu NC, Peter-Ikechukwu AI, Ojimba NC. A comparative study and determination of glycemic indices of three yams cultivars (*Dioscorea rotundata*, *Dioscorea alata* and *Dioscorea domentorum*). *Pakistan Journal of Nutrition* 2012; 11(6) 547-552.
22. Ogbo CF, Agu CK. Proximate composition, physiological changes during storage, and shelf life of some Nigerian varieties of yam (*Dioscorea* species). *Journal of Scientific Research and Reports* 2014; 3(4): 553-562
23. Turner ND, Lupton JR. Dietary fiber. *Adv in Nutr* 2011; 2: 151-152
24. Norman NP, Joseph HH. *Food Science*, 5th edn, Chapman and Hall Publishers, New York, 1995, pp 55
25. Shajeela PS, Mohan VR, Jesudas LL, Soris PT. Nutritional and antinutritional evaluation of wild yam (*Dioscorea* spp). *Tropical and Subtropical Agroecosystems* 2011; 14: 723-730
26. Alinnor JI, Akalezi CO. Proximate and mineral composition of *Dioscorea rotundata* (white yam) and *Colocasia esculenta* (white cocoyam). *Pakistan Journal of Nutrition* 2010; 9(10): 998-1001
27. Ezeocha VC, Ojimelukwe PC. The impact of cooking on the proximate composition and antinutritional factors of water yam (*Dioscorea alata*). *Journal of Stored Products and Postharvest Research* 2012; 3(13): 172-176
28. Oyenuga GO. *Nigeria's food and food stuffs*. Ibadan University Press, Ibadan, Nigeria, 1968, pp 99
29. Longe OG. Energy content of some Tropical starchy crops. *Nig. Agric. J* 1986; 21: 136
30. Arvidsson-Lenner R, Asp N, Axelsen M, Bryngelsson S, Haapa E, Jarvi A et al. Glycemic Index, Relevance for health, dietary recommendations and food labeling. *Scandinavian Journal of Nutrition* 2004; 48(2): 84-94
31. Salmeron J, Manson JE, Stampfer MJ, Colditz GA, Wing AL, Willet WC. Dietary fiber, glycemic load and the risk of non-insulin dependent diabetes mellitus in women. *JAMA* 1997; 277: 472-7
32. Liu S, Willet WC, Stampfer MJ, Hu FB, Franz M, Sampson L et al. A prospective study of dietary glycemic load, carbohydrate intake and risk of coronary heart disease in US women. *Am J Clin Nutr* 2004; 71: 1455-61
33. McKeown NM, Meigs JB, Liu S, Saltzman E, Wilson WPF, Jacques PF. Carbohydrate nutrition, insulin resistance, and the prevalence of the metabolic syndrome in the Framingham Offspring cohort. *Diabetes Care* 2004; 27: 538—546

34. Liu S, Manson JE, Buring JE, Stampfer MJ, Willet NC, Rikder PM. Relationship between a diet with a high glycemic load and plasma concentrations of high-sensitivity C-reactive protein in middle-aged women. *Am J Clin Nutr* 2002; 75: 492-8
35. Rimbawan dan A. Siagian . Indeks Glikemik Pangan. Penebar Swadaya, Jakarta 2004. 124 hlm.
36. Foster-Powell K, Host SHA, Brand-Miller JC. International table of glycemic index and been worth glycemic load. *Am J Clin Nutr* 2002; 76: 5-56
37. Amanda M, Wu M, Lu S, Lin J. Glycemic index, glycemic load and insulinemic index of Chinese starchy foods. *World Journal of Gastroenterology* 2010; 16(39): 4973-4979
38. Nattapulwat N, Purkkao N, Suwithayapanth O. Evaluation of Native and carboxymethyl Yam (*Dioscorea esculenta*) starches as Tablets disintegrants. *Silpakorn U Science and Tech J* 2008; 2(2): 18-25
39. Rugchati O, Mahawongwiriya K, Thanacharoenchanaphas K. Some characteristics of biodegradable film substituted by yam (*Dioscorea alata*) starch from Thailand. *International Journal of Biological, Veterinary, Agricultural and Food Engineering* 2013; 7(9): 499-502
40. Freibas RA, Paula RC, Feitosa JPA, Rocha S, Sierakowski MR. Amylose content, rheological properties, gelatinization kinetics of yam (*Dioscorea alata*) and Cassava (*Manihot utilissima*) starches. *Carbohydrate Polymers* 2004; 55(1): 3-8
41. Riley CK, Wheatley AO, Asemota HN. Isolation and characterization of starch from eight *Dioscorea alata* cultivars grown in Jamaica. *African Journal of Biotechnology* 2006; 5(17): 1528-1536
42. Noda T, Takahata Y, Nagata T . Developmental changes in the properties of sweet potatoe starches. *Starch/Starke* 1992; 44: 405-409
43. Lineback DR, Ke CH. Starches and low molecular weight carbohydrate from chick pea and horse bean flours. *Cereal Chem* 1975; 52: 334-347
44. Zhang F, Cogswell ME, Gillespie C, Fang J, Loustalot F, Dai S et al. Association between usual sodium and potassium intake and blood pressure and hypertension among US adults: NHANES 2005-2010. *PLoS ONE* 2013; 8(10): e75289. DOI:10.1371/journal.pone.0075289
45. WHO Website. Available: http://whqlibdoc.who.int/trs/who_trs_916.pdf
46. National Research Council (NRC). Recommended Dietary Allowance, 10th edn, National Academic Press, Washington DC, USA, 1989.