

Variations in $\delta^{13}\text{C}$ rates and Crassulacean acid metabolism of six different *Coleus* species

ABSTRACT

Photosynthetic adaptations and biomass productivity of six *Coleus* species *Coleus aromaticus*, *barbatus*, *blumei*, *zeylanicus*, *forskholii*, and *ambonicus* were studied by determining their photosynthetic rates, Crassulacean acid metabolism (CAM) and carbohydrate levels. CAM metabolism was investigated by studying the activities of phosphoenol pyruvate carboxylase (PEPC), PEP carboxykinase and diurnal fluctuations of malate dehydrogenase along with levels of total organic acids, malic acid and citric acids. Carbohydrates such as sucrose, starch, glucose and fructose were assayed spectrophotometrically. Significant differences among the six *Coleus* species were observed in photosynthesis levels and CAM metabolic pathway enzymes whose activities were correlated with the aerial biomass. Our study indicates that *Coleus* is a CAM plant with Nicotineamine Adenine Di nucleotide (Phosphate) (NAD (P)) as a preponderant malic enzyme and diurnal fluctuations in the levels of carbohydrates reveals that *Coleus* utilizes reservoir of starch for the synthesis of malic acid. $\Delta^{13}\text{C}$ accumulation is more in *Coleus aromaticus* and *ambonicus* whose aerial biomass values are also high indicating the correlation between carbon fixation and biomass accumulation.

KEY WORDS: Aerial biomass; CAM path way; carbohydrates; *Coleus* species; photosynthetic carbon fixation; organic acids.

ABBREVIATIONS

CAM: Crassulacean acid metabolism; PEPC: Phosphoenol pyruvate carboxylase; PEPCCK: Phosphoenol pyruvate carboxykinase; MDH: Malate dehydrogenase; NAD(P): Nicotineamine adenine di nucleotide (phosphate)

1. INTRODUCTION

Crassulacean Acid Metabolism (CAM) is one of the three forms of the photosynthetic carbon assimilation and is considered as one of the important models for probing the evolution of complex traits in response to changing environment [1]. CAM form of carbon assimilation has been characterized in 16,000 species of 328 genera in 33 families including Crassulaceae, Lamiaceae, Polypodiaceae Polypodiaceae etc. [2]. The mechanism of CAM is typically characterized by the acquisition of nocturnal CO_2 via

protein kinase activated phosphoenol pyruvate carboxylase (PEPC), leading to the formation of organic acids mainly malic acid, which are stored in the large central cell sap vacuoles. In the subsequent light period, organic acid is released from the vacuole, decarboxylated to release CO₂ for assimilation in the Calvin cycle behind closed stomata [3]. Based on the differences in the decarboxylation of malic acid, ~~two~~ three subtypes of CAM plants are known, NAD-ME type, NAD(P)-ME type and PEPC type [4]. CAM plants perform nocturnal recycling of CO₂ through malic acid for the purpose of photosynthesis. CAM plants ~~are possessing~~ possess high efficiency in water utilization by losing 50-100g of water per 1 g of CO₂ fixed in comparison to the 250-300g of water by the C₄ and 400-500 gm of water by the C₃ plants respectively [5]. ~~They have~~ CAM plants contain scotoactive stomata, which opens ~~only~~ during night facilitating the CO₂ ~~during the night time and leading to reduced photorespiration and leading to reduce photorespiration.~~ CAM plants are also provided with various morphological, anatomical features such as thick cuticles, low surface-to-volume ratios, large cells and vacuoles with enhanced water storage capacity (succulence), reduced stomatal size and frequency, which minimizes water loss [6].

~~Genus~~ *Coleus* (Lamiaceae) is an important plant used traditionally in folk medicine in India ~~belongs to family Labiatae and is popular CAM plant in India~~ due to its potential medicinal value as an expectorant, for the treatment of bronchitis, eczema and rheumatism. *Coleus aromaticus* and *Coleus zeylanicus* are aromatic perennial herbs that are native to Indonesia. *C. aromaticus* is used for chronic cough and asthma. It is considered to be antispasmodic, stimulant and stomachic and is used for the treatment of headache, fever, epilepsy and dyspepsia. *Coleus blumei* is a perennial native to Java, but now spread over many parts of tropical Africa, Europe and Asia, popular as an ornamental plant. *Coleus forskohlii*, an Indian herb, produces the labdane diterpenoid forskolin in its tuberous roots, which has been shown to be a hypertensive agent with spasmolytic, cardiotonic and platelet aggregation inhibitory activity, antiglaucomic in nature and a potent stimulator of the enzyme adenylate cyclase, which increases the levels of cAMP affecting cardiac muscle contraction, blood and intraocular pressure, cancer, eczema, rheumatism and obesity [7] (Duke and Mary, 2002). The medicinal properties of this plant has been well studied but the photosynthetic performance of this CAM plant was not studied. ~~This plant is mainly cultivated in hilly regions where the water resources are limiting and is capable of surviving under a variety of agro climatic conditions. However the photosynthesis mechanisms were not studied in this medicinally important plant species.~~ Previous studies have mentioned the role of environment on gas exchange and CAM mechanisms of *Plectranthus marrubioides*, *P. parviflorus* and *P. prostratus* (Lamiaceae) [8,9].~~the present study~~ Here, we report the photosynthesis and CAM mechanisms of six *Coleus* species, *C. aromaticus*, *C. barbatus*, *C. blumei*, *C. zeylanicus*, *C. forskohlii*, and *C. ambonicus* and their aerial biomass levels. Characterizing the *Coleus* species depending on the mechanism of CAM photosynthesis, which utilizes either malate or ~~PEPC~~ ~~carboxykinase~~ and identification of starch or sucrose carbohydrate reservoir pool utilized for the carbon fixation, are few of the important aspects of the present study.

2. MATERIALS AND METHODS

2.1 Plant Material and Growth Conditions

59 Six *Coleus* species namely ~~aromaticus, barbatus, blumei, zeylanicus, forskholii, and ambonicus~~ *C. aromaticus*, *C. barbatus*, *C.*
60 *blumei*, *C. zeylanicus*, *C. forskholii*, and *C. ambonicus* were propagated in the GITAM University botanical garden,
61 Visakhapatnam, India in 30 cm pots under 12h natural photoperiod [~~Irradiance (400-700 nm) of~~ 1600-1800 μ moles $\text{m}^{-2} \text{s}^{-1}$]
62 with day/night temperatures of 30 °C/23 °C and approximate relative air humidity of 60%. The plants were well watered with
63 normal tap water. Five pots were used for each *Coleus* species and one plant is grown per pot. Third or fourth fully expanded
64 leaf from the top of the 3-month-old plant was collected for all physiological experiments and aerial biomass measurements
65 made by measuring the mass of the above ground plant.

66 2.2 Carbon Isotope Determination

67 Samples were prepared by drying the *Coleus* leaves in a hot air oven at 80 °C for 72h. Completely dried leaves were grinded
68 into fine powder in a mortar with pestle. 2mg dry powder of each *Coleus* species leaf was subjected in the Flash Elemental
69 Analyzer (NA 1112, Carlo Erba, Italy) interfaced to an Isotope Ratio Mass Spectrometer (IRMS; Delta-Plus, Thermo-Finnigan,
70 Bremen, Germany) via a continuous flow device (ConFlo-III). The carbon isotopic composition of plant samples ($\delta^{13}\text{C}_p$) was
71 determined with an analytical precision of less than 0.1%. Carbon isotope discrimination ($\Delta^{13}\text{C}$) was computed according to
72 [10] assuming the isotopic composition of atmospheric air ($\delta^{13}\text{C}_a$) to be -8%.

$$73 \quad \Delta^{13}\text{C} (\text{‰}) = [\delta^{13}\text{C}_a - \delta^{13}\text{C}_p] / [1 + \delta^{13}\text{C}_p / 1000].$$

74 2.3 Extraction and Assay of Enzymes

75 All extractions were performed at 4°C. The leaf blades (10 g) were homogenized in a grinding medium consisting of 100 mM
76 Tris-HCl (pH 7.8), 5 mM DTT, 10 mM MgCl_2 , 1 mM EDTA, 5 mM magnesium acetate and 1.5% PVP at 4°C. The
77 homogenate was squeezed through four layers of cheesecloth and then centrifuged at 10000 g for 10min. The solution was
78 filtered to remove the cellulose and washed thrice with the extraction medium. The protein was precipitated with 75% (m/v)
79 ammonium sulphate and spun at 30000 g for 30 min and the precipitate was dissolved in 50 mM Tris-HCl buffer (pH 8.0)
80 containing 1 mM DTT and 0.2 mM NADPH.

81 ~~PEP~~Carboxylase (EC 4.1.1.31) was assayed spectrophotometrically at 340 nm at 30°C in a final volume of 3ml following the
82 reduction of NADH as described by [11]. Assay buffer comprises 0.1M Tris-HCl (pH 7.8), 10 μ M MgCl_2 , 10 mM NaHCO_3 ,
83 5mM PEP, 0.5 μ M NADH and 0.2 ml of enzyme extract. Malate dehydrogenase (EC 1.1.1.37) was assayed according to [12].
84 Assay buffer comprise of 0.1M Tris-HCl (pH 7.8), 0.5 μ M oxaloacetic acid, 10 μ M of MgCl_2 , 0.4 μ M NADH and 0.2 ml of

85 enzyme extract. PEP carboxykinase (EC 4.1.1.49) was assayed spectrophotometrically according to [13]. Carboxylation
86 activity was measured following the oxidation of NADH by malate dehydrogenase at 340 nm using 100 mM HEPES (pH
87 7.0), 4% 2-mercapto ethanol, 100 mM KCl, 90 mM KHCO₃, 5mM PEP, 1 mM ADP, 10 µm MnCl₂, 4 mM MgCl₂, 0.14 mM
88 NADH and 6 units of malate dehydrogenase.

89 NAD-ME (EC 1.1.1.38) was assayed spectrophotometrically at 340 nm according to [14] using 5mM malate, 25mM HEPES-
90 KOH of pH 7.2, 2 mM NAD, 2.5 mM MnCl₂, 5mM DTT and 500 µM fructose bis phosphate and 0.2 ml enzyme extract in a
91 volume of 1 ml assay mixture. NADP-ME (EC 1.1.1.40) was assayed spectrophotometrically at 340 nm according to [15].
92 Assay mixture contained 50 mM Tris-HCl (pH 7.3), 10 mM MgCl₂, 0.5 mM NADP and 10 mM l-malate with 0.2 ml of
93 enzyme extract.

94 2.4 Titratable Acidity, Malic Acid and Citric Acid Contents

95 Day and night oscillations of titratable acidity were measured using fully expanded *Coleus* leaves according to collected
96 during 8A.M. and 8 P.M. [16]. 1g of leaf tissue was cut into small pieces was, boiled in a beaker with 50 ml of distilled water
97 for 30 minutes and the slurry was filtered. 2.5 ml of extract was neutralized with 5 ml of 1N NaOH for 15 minutes and 1 drop
98 of phenolphthalein was added. The contents were titrated against 1N HCl till the solution becomes colour less [17].

99 Content of malic acid was determined according to [18]. 1 gram leaf sample was homogenized in a mortar and pestle with
100 0.6N perchloric acid. The sample was centrifuged at 15000g for 5 minutes. 0.9 1 ml of supernatant solution was collected and
101 neutralized with 80 µl of 5M K₂CO₃, centrifuged at 15000 g for 5 min. 0.5 ml of supernatant was incubated with 1.5 ml of
102 assay buffer containing 0.5 M glycine, 0.4 M hydrazine (pH 9.0) and 0.1 ml of 40 mM NAD. The reaction mixture was pre-
103 incubated for 30 min at room temperature before the addition of 5µl of malate dehydrogenase and the increase in the
104 absorbance at 340 nm due to the production of NADH was measured.

105 2.5 Carbohydrate Contents

106 The contents of starch and sucrose in the leaf tissues were estimated enzymatically according to the method of [19]. For
107 estimation of starch, eight leaf discs of 1 cm diameter were extracted with 80% ethanol 4-5 times at 80 °C. Extracts were
108 pooled and stored for the sucrose estimation. Ethanol extracted leaf discs were suspended in 1 cm³ of 0.2 M KOH and boiled
109 for 30 minutes. Tubes were cooled to room temperature and 0.2 cm³ of 1 M acetic acid was added to each tube and incubated at
110 55 °C for 30 minutes to hydrolyze starch. Reaction was stopped by boiling the tubes for 1 minute. 0.2 cm³ of the extract was
111 added to 0.2 cm³ of distilled water and 1cm³ of glucose enzyme reagent (Sigma). Tubes were incubated at 37 °C for 20 minutes

and the absorbance was read at 492 nm. For estimating the sucrose contents, ethanolic extracts were used. 0.2 cm³ of the extract was added to 0.3 cm³ of glucose reagent and incubated at 37 °C for 30 minutes. 75 mm³ of invertase was added and incubated at 37 °C for 30 minutes. The absorbance was read at 492 nm. The reading was proportional to the original plus glucose liberated due to the invertase action on sucrose. To determine the sucrose contents, the assays were run simultaneously without invertase for glucose contents. Sucrose standards were determined by comparing the difference in the absorbance of the two samples with that obtained from sucrose standards.

The glucose and fructose contents in the 80% ethanolic extract were determined according to [20] and [21] respectively. 500 mg of leaf tissue was homogenized in 80% ethanol and centrifuged at 5,000 rpm for 10 min. For glucose estimation, 1 ml of supernatant was added to 5 ml of anthrone reagent and incubated in boiling water bath for 30 minutes. The tubes were cooled to room temperature and the absorbance was measured at 620 nm. Fructose was estimated by adding 1 ml of resorcinol reagent to 2 ml of the ethanolic extract and incubating the tubes in boiling water bath for 10 minutes. After cooling, absorbance of the sample was measured at 520 nm. Glucose and fructose concentrations were calculated using a standard curve.

2.6. Statistical Analysis

Statistical analysis of absolute values was performed by analysis of variance (ANOVA). The values are the mean $\pm$ SD for five samples in each group. The significance of the data was tested by student's *t* test. *P* values greater than 0.05 were considered as significant.

3. RESULTS & DISCUSSION

Among the six *Coleus* species considered for this study, *C. aromaticus* and *C. ambonicus* possessed significantly higher $\Delta^{13}\text{C}$ values and biomass yields than other species (Table 1). A significant variation in the stable isotope levels was observed among the *Coleus* species which is correlated with the PEP Carboxylase and malate dehydrogenase activities (Fig. 1 and 2). PEP Carboxykinase activities of six *Coleus* species were compared with NAD-ME and NADP-ME (Fig. 3) in which, *Coleus aromaticus* and *ambonicus* were found to possess high NADP-ME enzyme activity where as *barbatus*, *blumei*, *zeylanicus*, *forskholii*, have shown NAD-ME activity indicating *Coleus* plants are exhibiting NAD(P)-ME type of photosynthesis carbon fixation. Day- night studies of the organic acid levels in six *Coleus* species reveals that concentrations of titrable acids, malic acid and citric acid were high during night time which were reduced during day, correlating with malate dehydrogenase activities (Fig. 4, 5 and 6). Levels of carbohydrates such as glucose and sucrose were higher than that of starch in all *Coleus* species considered for this study and the sugar levels were high during night than day time (Fig. 7). Our studies on the overall photosynthesis activities, accumulation of organic acids and carbohydrate levels have shown that *C. aromaticus* and

141 *C.ambonicus* were possessing better showing higher photosynthetic performance and biomass yields compared with other four
142 species.

143 CAM photosynthesis is one of the remarkable examples of ecophysiological adaptations of plants to arid conditions [22]. CAM
144 is an important elaboration of photosynthetic carbon fixation that facilitates nocturnal CO₂ fixation initially using PEPC in the
145 cytosol leading to the formation of C4 organic acids, which gets stored in the vacuole followed by subsequent decarboxylation
146 during day time behind closed stomata, generating an internal CO₂ source that is re-assimilated by Rubisco in the chloroplast.
147 In the present study, the $\Delta^{13}\text{C}$ accumulation of carbon in the leaves of six *Coleus* species was studied, whose values are ranging
148 from -22.94 ‰ to -20.09 ‰ respectively (Table 1). $\Delta^{13}\text{C}$ values in the CAM plants *Tillandsia usneoides*, *Didierea*
149 *madagascariensis* and *Microsorium punctatum* were reported to be -19.8‰ [23], -21.2‰ [24], and -22.6‰ [25] respectively.
150 ^{13}C values in CAM plants serves as an indicator of their biomass derived from nocturnal CO₂ fixation compared to that of
151 diurnal CO₂ fixation because PEP carboxylase that fixes nocturnal CO₂ discriminates less slightly higher against ^{13}C than
152 Rubisco, which is responsible for the day time CO₂ uptake [26]. CAM plants gain maximum CO₂ during night by PEP
153 carboxylase activity, their ^{13}C values and biomass levels will be similar to that of C4 plants. The aerial biomass levels were
154 measured in six *Coleus* species, which shows that *aromaticus* and *ambonicus* possessed high aerial biomass values than
155 *blumei*, *barbatus*, *forskholii* and *zeylanicus* respectively (Table 1).

156 CAM is a well known water conserving modification of photosynthesis characterized by the nocturnal CO₂ assimilation
157 catalysed by the enzyme PEPCarboxylase leading to the synthesis of organic acids [27]. Activity of PEPCarboxylase
158 estimated in the six *Coleus* species was depicted in Figure 1, which shows that *C.aromaticus* and *ambonicus* were possessing
159 high PEPC activities when compared with the other *Coleus* species. Malate dehydrogenase (MDH) is a ubiquitous enzyme,
160 which serves in a wide range of metabolic processes of plants. MDH participation has been identified in key metabolic
161 pathways such as Krebs cycle, photorespiration and metabolite shuttling [28]. MDH plays a remarkable role in the CAM
162 plant's photosynthesis carbon fixation by reducing oxalo-acetic oxaloacetic acid to malic acid in CAM plants [29]. In the
163 present study, day and night activities of MDH in six *Coleus* species were studied (Fig. 2). The night time activity of MDH was
164 high in *aromaticus* (331.65 μ moles mg protein min⁻¹) and *ambonicus* (307.4 μ moles mg protein min⁻¹) compared to that of the
165 other four species. The day time activities were less among all *Coleus* species without much difference in their activity (Fig.
166 2), which shows the limited role of MDH during day time in *Coleus*. NAD-MDH has been isolated, purified and its day and
167 night fluctuation has been studied in *Ananas comosus*, a PEPCK type of CAM plant [30]

168 Malic acid released from the vacuoles during day time immediately undergoes decarboxylation reaction specific to NAD or
 169 NADP in malic enzyme type CAM plants, yielding pyruvate and CO₂. In ~~PEPCK –phosphoenolpyruvate carboxykinase~~ type
 170 CAM plants, malic acid will be converted to OAA prior to its decarboxylation, yielding PEP and CO₂. In the present study the
 171 activities of NADP-ME, NAD-ME and PEPCK were studied in six *Coleus* species (Fig. 3). *C. aromaticus* and *ambonicus* were
 172 possessing high rates of NADP-ME activity where as *C. blumei*, *barbatus*, *forskholii* and *zeylanicus* were exhibiting high
 173 NAD-ME activity indicating that *Coleus* is a CAM plant with NAD(P)-ME as preponderant enzyme (Fig. 3). The activities of
 174 NADP-ME were high in *Coleus*, when compared to that of NAD-ME and PEPCK. Species variation with respect to PEPCK
 175 enzyme was observed in *A.comosus* and *C.lycopodioides* with lower oxidation of malic acid [31].
 176 Malic acid is a ubiquitous vacuolar anion, synthesized nocturnally as a result of CO₂ fixation and stored in the vacuoles. Malic
 177 acid is a central intermediary in the process of CAM photosynthesis. Nocturnal malic and citric acid accumulation decreases
 178 the osmotic potential and ~~facilitates~~ enhances ~~the~~ water uptake [32]. Daytime decarboxylation of organic acids will increase
 179 internal CO₂ concentration enabling CAM plants to close stomata, at least partially and facilitating the maintenance of high
 180 photosynthesis rates [33]. In the present study the day time and night time levels of organic acids such as ~~titrable acid number~~
 181 ~~(TAN)~~, malic acid and citric acid were estimated in six *Coleus* species (Fig. 4, 5 and 6). *Coleus aromaticus* and *C. ambonicus*
 182 accumulated high levels of organic acid ~~accumulation~~. Such higher levels of organic acid accumulation can be observed more
 183 in NADP-ME type CAM plants since their oxidation capacity is higher in contrast to PEPCK type plants. Nocturnal synthesis
 184 of citric acid in CAM plants might serve as the reservoir for the CO₂. High concentrations of citric acid exceeding 200 mM has
 185 been reported in CAM plants such as *Bryophyllum calycinum*, *Clusia minor*, *Clusia rosea*, *Notonia petrea* and *Cissus*
 186 *quadrangularis* [34].
 187 Carbohydrates plays pivotal role in balancing the day/night CO₂ fixation of CAM plants [35]. The night time uptake of CO₂ in
 188 CAM plants is supported by the degradation of carbohydrate reserve pool for the synthesizing the PEP substrate and 20% of
 189 leaf biomass is utilized for ~~this purpose the synthesis~~. Utilization of carbohydrate reserves for the synthesis of organic acids is
 190 the unique feature of CAM plants which is 20-170 times the carbohydrates catabolized for the respiration [36]. ~~In the present~~
 191 ~~study, The~~ day and night levels of carbohydrate reserves were determined in six *Coleus* species by estimating the foliar
 192 glucose, sucrose and starch contents (Fig. 7). The levels of glucose and sucrose were higher in all *Coleus* during night time
 193 than day except starch contents whose day time concentrations were higher than night time. Formation of nocturnal malate in
 194 CAM plants will reduce the carbohydrate levels as evidenced in the analysis of their sap [35]. In the present study, the levels of
 195 nocturnal starch were lower than that of glucose and sucrose indicating that starch might serve as the carbohydrate reservoir for

196 carbon fixation in *Coleus*. Among the six *Coleus* plants considered for this study, *aromaticus* and *ambonicus* were possessing
197 high levels of glucose, sucrose, fructose and low levels of starch. Starch is the source of malate synthesis in the members of
198 crassulaceae and cactaceae [37, 38]. In *M. crystallinum*, carbon allocated to starch is maintained at the expense of the soluble
199 sugar pool during acclimation to salinity [39]. Deficiency in leaf starch and plastidic phosphoglucomutase (PGM) in CAM-
200 deficient mutants of *M. crystallinum* indicated the role of starch in providing substrate for nocturnal carboxylation [8].

201 4. CONCLUSION

202 CAM plants are more efficient in water conservation and are having the unique mechanism of nocturnal and diurnal CO₂
203 fixation. In the present study the photosynthetic performance and CAM pathway of *Coleus* has been demonstrated in six
204 species among which *C. aromaticus* and *C. ambonicus* have shown to possess high photosynthetic efficiency and biomass
205 levels, which might be due to the presence of high NADP-ME enzyme activity when compared with *barbatus*, *blumei*,
206 *zeylanicus*, *forskholii*, which are possessing NAD-ME activity. Overall *Coleus* plants following NAD(P)-ME type of
207 photosynthesis carbon fixation. Such studies can be useful in the utilization of the low water lands for the growth of
208 medicinally important plants like *Coleus* which are capable of surviving with less water availability. Further studies are in
209 progress for studying the response of these plants to abiotic stress which will give more information regarding the behavior of
210 these plants.

211 COMPETING INTERESTS

212 There are no Competing Interests behalf of authors.

213 AUTHORS' CONTRIBUTION

214 All authors contributed equally for this study. Authors affirm that this work is in genuine.

215 REFERENCES

- 216 1. Yamori W, Kouki H, Danielle AW. Temperature response of photosynthesis in C₃, C₄, and CAM plants: temperature
217 acclimation and temperature adaptation. *Photosynthesis Research*.2014; 119:101-117.
- 218 2. Herrera A. Crassulacean acid metabolism and fitness under water deficit stress: if not for carbon gain, what is
219 facultative CAM good for? *Annals of Botany*.2009;103: 645–653.
- 220 2. Fernanda R, Robert EB. Evolutionary aspects of crassulacean acid metabolism. *Oecologia Australis*. 2010; 14: 359-
221 368.
- 222 3. Luttge U. CO₂-concentrating: consequences in crassulacean acid metabolism. *Journal of Experimental Botany*. 2002;
223 53: 2131-2142.

4. Yamori W, Masumoto C, Fukayama H, Makino A. Rubisco activase is a key regulator of non steady state photosynthesis at any leaf temperature and, to a lesser extent, of steady state photosynthesis at high temperature. The Plant Journal.2012;71: 871–880.
4. Christopher JT, HOLTUM JAM. Patterns of carbon partitioning in leaves of crassulacean acid metabolism species during deacidification. Plant Physiology. 1996; 112: 393-399.
5. Rathore A, Jasrai TY. Growth and chlorophyll levels of selected plants with varying photosynthetic pathways (C3, C4 and CAM). International Journal of Scientific & Engineering Research.2013; 4:1–4.
5. Sayed OH. Crassulacean acid metabolism 1975-2000 a check list. Photosynthetica. 2001; 39: 339-352.
6. Cushman JC, Agarie S, Albino RL, Elliot SM, Taybi T, Borland AM . Isolation and characterization of mutants of common ice plant deficient in crassulacean acid metabolism. Plant physiology. 2008; 147:228-238.
7. Girish KR, Dayananda S and Attipalli R Reddy. Free radical quenching activity and polyphenols in three species of *Coleus*. Journal of Medicinal Plants Research. 2008; 2: 285-291.
8. Herppich WB, Herppich M, von Willert DJ. Ecophysiological investigations on plants of the genus *Plectranthus* (*Lamiaceae*). Influence of environment and leaf age on CAM, gas exchange and leaf water relations in *Plectranthus marrubioides* Benth. Flora. 1998; 193:99–109.
9. Nobel PS. Achievable productivities of CAM plants: basis for high values compared with C3 and C4 plants. New Phytologist. 1991; 119:183-205.
10. Bindumadhava H, Sheshshayee MS, Shashidhar GP, Prasad TG, Udayakumar M (2005) The ratio of stable carbon and oxygen isotope discrimination ($\Delta^{13}\text{C}/\Delta^{18}\text{O}$) reflects the variability in leaf intrinsic carboxylation efficiency in plants. Current Science 89: 1256 – 1258.
11. Chu C, Dai Z, Ku MSB, Edwards GE .Induction of Crassulacean acid metabolism in the facultative halophyte *Mesembryanthemum crystallinum* by abscisic acid. Plant Physiology.1990; 93: 1253–1260.
12. Gnanam A, Francis K. Distribution of photosynthetic enzymes in the leaf cells types of *Sorghum vulgare*. Plant Biochemical Journal.1976;3:11-23.
13. Walker RP, Leegood RC. Phosphorylation of phosphoenolpyruvate carboxykinase in plants: studies in plants with C₄ photosynthesis and Crassulacean acid metabolism and in germinating seeds. Biochemical Journal. 1996;317: 653–658.
14. Murata T, Oshugi R, Matsuoka M, Nakamoto M. Purification and characterization of NAD ME from leaves of *Eleusine coracana* and *Panicum dichotomiflorum*. Plant Physiology.1989;89: 316–324.
15. Ferreyra MLF, Andreo CS, Podesta FE .Purification and physical and kinetic characterization of a photosynthetic NADP-dependent malic enzyme from the CAM plant *Aptenia cordifolia*. Plant Science. 2003;164:95–102.
16. Mawson BT, Zaugg MW. Modulation of light-dependent stomatal opening in isolated epidermis following induction of crassulacean acid metabolism in *Mesembryanthemum crystallinum* L. Plant Physiology.1994;144:740-746.

17. Hazell T, Johnson IT. Effects of food processing and fruit juices on in-vitro estimated iron availability from cereals, vegetables and fruits. *Journal of the Science, Food and Agriculture*. 1987;38:73-82.
18. Delhaize E, Craig S, Beaton CD, Bennet RJ, Jagadish VC, Randall PJ .Aluminum tolerance in wheat (*Triticum aestivum* L.) II. Aluminum stimulated excretion of malic acid from root apices. *Plant Physiology*. 1993;103:695-702.
19. Ramachandra Reddy A, Reddy KR, Hodges HF. Mepiquat chloride (PIX)-induced changes in photosynthesis and growth of cotton. *Plant Growth Regulations*. 1996; 20: 179-183.
20. Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F .Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*. 1956;28:350-356.
21. Ashwell G *Methods in Enzymology*, Vol. 3, pp 75. In: Colowick S J and Kaplan N O(eds). Academic Press, New York; 1957.
22. Ramachandra Reddy A, Sundar D, Gnanam A. Photosynthetic flexibility in *Pedilanthus titihymaloides* Poit, a CAM plant. *Journal of Plant Physiology*.2003;160:75-80.
23. Griffiths H,Smith JAC. Photosynthesis pathways in the Bromeliaceae of Trinidad: Relations between life-forms, habitat preferences and the occurrence of CAM. *Oecologia*. 1983; 60: 176-184.
24. Winter K. $\delta^{13}\text{C}$ values of some succulent plants from Madagascar. *Oecologia*.1979;40: 103 -112.
25. Holtum JAM, Winter K. Degrees of Crassulacean acid metabolism in tropical epiphytic and lithophytic ferns. *Australian Journal of Plant Physiology*.1999;26:749–757.
26. Winter K, Holtum JAM. Environment or development? Lifetime net CO_2 exchange and control of the expression of crassulacean acid metabolism in *Mesembryanthemum crystallinum*. *Plant Physiology*.2007;143: 98–107.
27. Dodd AN, Borland AM, Richard PH, Griffiths H, Maxwell K .Crassulacean acid metabolism: plastic, fantastic. *Journal of Experimental Botany*. 2002;53:569-580.
28. Gietl C. Malate dehydrogenase isoenzymes: cellular locations and role in the flow of metabolites between the cytoplasm and cell organelles. *Biochim. Biophys. Acta*. 1992; 1100: 217–234.
29. Karina EJT, Podestá FE. Purification and characterization of an NAD-dependent malate dehydrogenase from leaves of the crassulacean acid metabolism plant. *Aptenia cordifolia*. *Plant Physiology and Biochemistry*. 2003; 41: 97–105.
30. Cuevas I, Podestá F.E. Purification and physical and kinetic characterization of an NAD-dependent malate dehydrogenase from leaves of pineapple (*Ananas comosus* L.). *Physiol. Plant*. 2000; 108: 240–248.
31. Peckman K, Dieter JVW, Craig EM, Werner BH. Mitochondrial respiration in ME-CAM, PEPCK-CAM, and C_3 succulents: comparative operation of the cytochrome, alternative, and rotenone-resistant pathways. *Journal of Experimental Botany*. 2012; 63:2909–2919.
32. Von Willert DJ, Eller BM, Werger MJA, Brinckmann E, Ihlenfeld HD .Life strategies of succulents in deserts with special reference to the Namib desert. Cambridge University Press; 1992.

33. Chheffings CM, Pantoja O, Aschroft FM, Smith JAC. Malatetransport and vacuolar ion channels in CAM plants. Journal of Experimental Botany 1997;48:623-631.

34. López-Bucio J, Martínez de la Vega O, Guevara-García A, Herrera EL. Enhanced phosphorus uptake in transgenic tobacco plants that overproduce citrate. Nature Biotechnology. 2000; 18:450–453.

35. Ceusters J, Anne MB, Ceusters N, Veerle V, Godts C, Maurice Proft PDe .Seasonal influences on carbohydrate metabolism in the CAM bromeliad Aechmea ‘Maya’: consequences for carbohydrate partitioning and growth. Annals of Botany. 2010; 105: 301–309.


36. Carnal NW, Black CC .Soluble carbohydrates as the carbohydrate reserve for CAM in pineapple leaves. Implications for the role of pyrophosphate:6-phosphofructokinase in glycolysis. Plant Physiology. 1989; 90: 91-100.

37. Sutton BG. The path of carbon in CAM plants at night. Aust. J. Plant Physiol. 1975; 2: 377-387.

38. Whiting BH, Van De Venter HA, Small JGC. Crassulacean acid metabolism in jointed cactus (*Opuntia aurantiaca*) Lindley. Agroplanta. 1979; 11: 41-43.

39. Borland AM, Dodd AN .Carbohydrate partitioning in crassulacean acid metabolism plants: reconciling potential conflicts of interest. Functional Plant Biology. 2002; 29: 707–716.

Fig.1



317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339

Fig. 1. PEPcarboxylase activity in 6 different *Coleus* species. Each point is an average of five independent determinations $\pm$ SE, ($t_{(4)} = 9.2$, $P < 0.05$).

Fig. 2




Fig. 2. Day and night activities of Malate dehydrogenase in 6 different *Coleus* species. Each point is an average of five independent determinations $\pm$ SE, ($t_{(4)} = 8.4$, $P < 0.05$).

Fig. 3




Fig. 3. Activities of malic enzymes NAD-ME, NADP-ME and PEPCK in 6 different *Coleus* species. Each point is an average of five independent determinations $\pm$ SE, ($t_{(4)} = 6.2$, $P < 0.05$).

Fig. 4




Fig. 4. Total organic acid content among six *Coleus* species during day and night. Each point is an average of at least five independent determinations $\pm$ SE, ($t_{(4)} = 11.2$, $P < 0.05$).




Fig. 5

385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406

Fig. 5. Variations in the day and night Malic acid contents of six *Coleus* species. Each point is an average of at least five independent determinations $\pm$ SE, ($t_{(4)} = 4.9, P < 0.05$).

Fig. 6




Fig. 6. Day and night variations in the citric acid contents of six different *Coleus* species. Each point is an average of at least five independent determinations $\pm$ SE, ($t_{(4)} = 5.6$, $P < 0.05$).

Fig. 7




Fig. 7. Carbohydrate levels in six different *Coleus* species during day and night. Each point is an average of at least five independent determinations $\pm$ SE, ($t_{(4)} = 10.2$, $P < 0.05$).

Table 1. $\Delta^{13}\text{C}$ (‰) Carbon isotope levels and aerial biomass contents in 6 *Coleus* Species. Each point is value represents an average of five independent determinations $\pm$ SE, ($t_{(4)} = 3.2$, $P < 0.05$) and ($t_{(4)} = 7.4$, $P < 0.05$).

428	Name of the plant	$\Delta^{13}\text{C}$ (‰)	Aerial biomass (g)
429			
430	<i>Coleus aromaticus</i>	-22.94 $\pm$ 0.65	793 $\pm$ 1.12
431	<i>Coleus ambonicus</i>	-22.75 $\pm$ 0.81	721 $\pm$ 0.89
432	<i>Coleus blumei</i>	-22.35 $\pm$ 0.10	672 $\pm$ 0.55
433	<i>Coleus barbatus</i>	-22.06 $\pm$ 0.23	599 $\pm$ 1.47
434	<i>Coleus forskholii</i>	-20.37 $\pm$ 0.45	591 $\pm$ 0.84
435	<i>Coleus zeylanicus</i>	-20.09 $\pm$ 0.63	572 $\pm$ 0.45
436			
437			
438			
439			