

Original Research Article

A Class of Implicit Six Step Hybrid Backward Differentiation Formulas for the Solution of Second Order Differential Equations.

Umaru MOHAMMED^{1*} and Raphael Babatunde ADENIYI²

¹ *Department of Mathematics and Statistics, Federal University of Technology, Minna, Niger State, Nigeria,*

² *Department of Mathematics, University of ILorin, ILorin, Nigeria*

E-mail(s): digitalumar@yahoo.com; umaru.mohd@futminna.edu.ng;
raphade@unilorin.edu.ng

Abstract

In this paper, we propose a class implicit six step Hybrid Backward Differentiation Formulas (HBDF) for the solution of second order Initial Value Problems (IVPs). The method is derived by the interpolation and collocation of the assumed approximate solution. The single continuous formulation derived is evaluated at grid point of $x = x_{n+k}$ and its second derivative at $x = x_{n+j}$, $j = 1, 2, \dots, k-1$ and $x = x_{n+\mu}$ respectively, where k is the step number of the methods. The interpolation and collocation procedures lead to a system of $(k+1)$ equations, which are solved to determine the unknown coefficients. The resulting coefficients are used to construct the approximate continuous solution from which the Multiple Finite Difference Methods (MFDMs) are obtained and simultaneously applied to provide the direct solution to IVPs. Numerical examples are given to show the efficiency of the method.

Keywords

Hybrid method; Backward Differentiation Formulas; Collocation; Interpolation; Second Order; Multiple Finite Difference.

Introduction

In recent times, the integration of Ordinary Differential Equations (ODEs) are investigated using some kind of block methods. This paper discusses the family of implicit Linear Multistep Method (LMM) for numerical integration of general second order ODEs which arise frequently in the area of science and engineering especially mechanical system, control theory and celestial mechanics and are generally written as: $y'' = f(x, y', y)$, $y(a) = y_0$, $y'(a) = \eta_0$ (1)

In practice the problems are reduced to systems of first order equations and any method for first order equations is used to solve them see Awoyomi [1]. It has been extensively discussed that due to the dimension of the problem after it has been reduced to a system of first order equations also, more often the reduced systems of ordinary differential equations (ODEs) is not well posed, unlike the given problem. The approach waste a lot of Computer time and human efforts, hence there is a need to develop algorithms to handle these classes of problems directly without any reduction to system of first order ODEs .

Development of LMM for solving ODE can be generated using methods such as Taylor's series, numerical interpolation, numerical integration and collocation method, which are restricted by an assumed order of convergence. In this paper we will consider the contribution of multi step collocation technique introduced by Onumayi et al [2] by deriving our new method. Some researchers have attempted the solution of directly using linear multistep methods without reduction to system of first order ordinary differential equations the include Mohammed et al [3], Yusuf and Onumayi [4] and Onumayi et al [5].

Block methods for solving ODEs have initially been proposed by Milne [6]. The Milne's idea of proceeding in blocks was developed by Rosser [7] for Runge-Kutta method. Also block Backward Differentiation Formulas (BDF) methods are discussed and developed by many researchers [8-16]. The method of collocation and interpolation of the power series

approximate solution to generate continuous LMM has been adopted by many researchers among them are (Houwen et al. [17], Fatunla [18], Jiaxiang [19]).

In this paper we are suggested a construction of six step HBDF method, it is self-starting and can be applied for the numerical solution of IVPs (Cauchy problem) for second-order ODEs.

Material and Method

We seek an approximation of the form

$$Y(x) = \sum_{j=0}^{r+s-1} \ell_j x^j \quad (2)$$

Where ℓ_j are unknown coefficients to be determined and $k < r$ and $s > 0$ are the number of interpolation and collocation points respectively. We then construct our continuous approximation by imposing the following conditions

$$Y(x) = y_{n+j}, \quad j = 0, 1, 2, \dots, k-1 \quad (3)$$

$$Y''(x_{n+k}) = f_{n+k} \quad (4)$$

We note that $y_{n+\mu}$ is the numerical approximation to the analytical solution $y(x_{n+\mu})$, $f_{n+\mu} = f(x_{n+\mu}, y_{n+\mu}, y'_{n+\mu})$.

Equations (3) and (4) lead to a system of $(k+1)$ equations which is solved by Cramer's rule to obtain ℓ_j . Our continuous approximation is constructed by substituting the values ℓ_j into equation (2). After some manipulation, the continuous method is expressed as

$$Y(x) = \sum_{j=0}^{k-1} \alpha_j(x) y_{n+j} + \alpha_\mu(x) y_{n+\mu} + h^2 \beta_k(x) f_{n+k} \quad (5)$$

Where $\alpha_j(x)$, $\beta_k(x)$ and $\alpha_\mu(x)$ are continuous coefficients. We note that since the general second order ordinary differential equation involves the first derivative, the first derivative formula

$$Y'(x) = \frac{1}{h} \left(\sum_{j=0}^{k-1} \alpha'_j(x) y_{n+j} + \alpha'_\mu(x) y_{n+\mu} + h^2 \beta'_k(x) f_{n+k} \right) \quad (6)$$

$$Y'(x) = \delta(x) \quad (7)$$

$$Y'(a) = \delta_0 \quad (8)$$

2.1 Specification of Methods

2.1.1 Six Step Methods with one- off -step point at interpolation.

To derive this methods, we use Eq.(5) to obtained a continuous 5-step HBDF method with the following specification : $r=7, s=1, k=6$. We also express $\alpha_j(x), \alpha_\mu(x)$ and $\beta_k(x)$ as a

functions of t , where $t = \frac{x - x_n}{h}$ to obtain the continuous form as follows:

$$y(x) = \alpha_0 y_n + \alpha_1 y_{n+1} + \alpha_2 y_{n+2} + \alpha_3 y_{n+3} + \alpha_4 y_{n+4} + \alpha_5 y_{n+5} + \alpha_{\frac{11}{2}} y_{n+\frac{11}{2}} + h^2 \beta_6 f_{n+6} \quad (9)$$

Where

$$\begin{aligned} \alpha_0(t) &= 1 - \frac{2223121}{850080}t + \frac{3014973}{1133440}t^2 - \frac{3157433}{2266880}t^3 + \frac{186435}{453376}t^4 - \frac{156729}{2266880}t^5 + \frac{13999}{2266880}t^6 - \frac{773}{3400320}t^7 \\ \alpha_1(t) &= \frac{27775}{3864}t - \frac{179425}{15456}t^2 + \frac{2088853}{278208}t^3 - \frac{231881}{92736}t^4 + \frac{126277}{278208}t^5 - \frac{3961}{92736}t^6 + \frac{227}{139104}t^7 \\ \alpha_2(t) &= -\frac{4125}{368}t - \frac{244245}{10304}t^2 - \frac{1109779}{61824}t^3 + \frac{409541}{61824}t^4 - \frac{26609}{20608}t^5 + \frac{7909}{61824}t^6 - \frac{157}{30912}t^7 \\ \alpha_3(t) &= \frac{25619}{1932}t - \frac{389547}{12880}t^2 + \frac{648047}{25760}t^3 - \frac{51789}{5152}t^4 + \frac{10755}{5152}t^5 - \frac{5601}{25760}t^6 + \frac{347}{38640}t^7 \\ \alpha_4(t) &= -\frac{57805}{5152}t - \frac{546379}{20608}t^2 - \frac{8589991}{370944}t^3 + \frac{403209}{41216}t^4 - \frac{794791}{370944}t^5 + \frac{9641}{41216}t^6 - \frac{1865}{185472}t^7 \\ \alpha_5(t) &= \frac{2211}{280}t - \frac{21309}{1120}t^2 + \frac{114827}{6720}t^3 - \frac{10049}{1344}t^4 + \frac{3817}{2240}t^5 - \frac{1301}{6720}t^6 + \frac{29}{3360}t^7 \\ \alpha_{\frac{11}{2}}(t) &= -\frac{17516}{5313}t - \frac{30368}{3795}t^2 - \frac{247696}{34155}t^3 + \frac{7312}{2277}t^4 - \frac{5072}{6831}t^5 + \frac{976}{11385}t^6 - \frac{928}{239085}t^7 \\ \beta_6(t) &= \frac{1}{10304}(1320t - 3254t^2 + 3023t^3 - 1385t^4 + 335t^5 - 41t^6 + 2t^7) \end{aligned}$$

Evaluating (9) at $x = x_{n+6}$ yields Hybrid Six step implicit method

$$y_{n+6} = \frac{257}{28336}y_n - \frac{51}{644}y_{n+1} + \frac{405}{1288}y_{n+2} - \frac{247}{322}y_{n+3} + \frac{3555}{2576}y_{n+4} - \frac{81}{28}y_{n+5} + \frac{768}{253}y_{n+\frac{11}{2}} + \frac{45}{644}h^2 f_{n+6} \quad (10)$$

Taking the second derivative of equation of equation (9), thereafter, evaluating the resulting continuous polynomial solution at $x = x_{n+2}, x = x_{n+3}, x = x_{n+4}, x = x_{n+5}, x = x_{n+\frac{11}{2}}$ we

generate five additional methods

$$y_{n+2} = -\frac{16829}{579590}y_n + \frac{81682}{142263}y_{n+1} + \frac{7724}{26345}y_{n+3} + \frac{30685}{94842}y_{n+4} - \frac{7498}{26345}y_{n+5} + \frac{960512}{7824465}y_{n+\frac{11}{2}} - \frac{2576}{5269}h^2f_{n+2} - \frac{26}{5269}h^2f_{n+6} \quad (11)$$

$$y_{n+3} = \frac{353}{85976}y_n - \frac{2915}{52758}y_{n+1} + \frac{6455}{11724}y_{n+2} + \frac{39035}{70344}y_{n+4} - \frac{391}{5862}y_{n+5} - \frac{3584}{290169}y_{n+\frac{11}{2}} - \frac{3220}{8793}h^2f_{n+3} + \frac{5}{17586}h^2f_{n+6} \quad (12)$$

$$y_{n+4} = -\frac{11601}{15281585}y_n + \frac{8468}{833541}y_{n+1} - \frac{20754}{277847}y_{n+2} + \frac{774072}{1389235}y_{n+3} + \frac{908316}{1389235}y_{n+5} - \frac{6680576}{45844755}y_{n+\frac{11}{2}} - \frac{92736}{277847}h^2f_{n+4} - \frac{540}{277847}h^2f_{n+6} \quad (13)$$

$$y_{n+5} = \frac{2663}{3217148}y_n - \frac{16075}{1974159}y_{n+1} + \frac{5675}{146234}y_{n+2} - \frac{9514}{73117}y_{n+3} + \frac{1266205}{2632212}y_{n+4} + \frac{13411328}{21715749}y_{n+\frac{11}{2}} - \frac{560}{3179}h^2f_{n+5} - \frac{5}{4301}h^2f_{n+6} \quad (14)$$

$$y_{n+\frac{11}{2}} = -\frac{5200659}{563806208}y_n + \frac{11352935}{140951552}y_{n+1} - \frac{90099405}{281903104}y_{n+2} + \frac{54767691}{70475776}y_{n+3} - \frac{773233395}{563806208}y_{n+4} + \frac{259721451}{140951552}y_{n+5} + \frac{34155}{157312}h^2f_{n+\frac{11}{2}} - \frac{7248285}{140951552}h^2f_{n+6} \quad (15)$$

Since our method is design to simultaneously provide the values of

$y_{n+1}, y_{n+2}, y_{n+3}, y_{n+4}, y_{n+5}, y_{n+\frac{11}{2}}, y_{n+6}$ at a block point $x_{n+1}, x_{n+2}, x_{n+3}, x_{n+4}, x_{n+5}, x_{n+\frac{11}{2}}, x_{n+6}$, the six

equations (10)-(15) are not sufficient to provide the solution for seven unknown

$y_{n+1}, y_{n+2}, y_{n+3}, y_{n+4}, y_{n+5}, y_{n+\frac{11}{2}}, y_{n+6}$. Thus, we obtain an additional method from (8), given by

$$850080h\delta_0 + 2223121y_0 - 6110500y_1 + 9528750y_2 - 11272360y_3 + 9537825y_4 - 6712596y_5 + 2805760y_{\frac{11}{2}} = 108900h^2f_6 \quad (16)$$

The derivatives are obtained from (7) by imposing that

$\delta(x_{n+\mu}) = \delta_{n+\mu}, \mu = \{j, v\}, j = 0, \dots, 6$, thus, we have

$$\begin{aligned}
h\delta_{n+1} &= -\frac{39411}{283360}y_n - \frac{17177}{11592}y_{n+1} + \frac{8037}{2576}y_{n+2} - \frac{8919}{3220}y_{n+3} + \frac{10747}{5152}y_{n+4} - \frac{387}{280}y_{n+5} + \frac{45184}{79695}y_{n+\frac{11}{2}} \\
&\quad - \frac{27}{1288}h^2f_{n+6} \\
h\delta_{n+2} &= \frac{1803}{80960}y_n - \frac{1061}{3312}y_{n+1} - \frac{12865}{15456}y_{n+2} + \frac{327}{184}y_{n+3} - \frac{4441}{4416}y_{n+4} + \frac{143}{240}y_{n+5} - \frac{3776}{15939}y_{n+\frac{11}{2}} + \\
&\quad \frac{3}{368}h^2f_{n+6} \\
h\delta_{n+3} &= -\frac{2833}{340032}y_n + \frac{695}{7728}y_{n+1} - \frac{2895}{5152}y_{n+2} - \frac{6493}{19320}y_{n+3} + \frac{11695}{10304}y_{n+4} - \frac{57}{112}y_{n+5} + \frac{5056}{26565}y_{n+\frac{11}{2}} \\
&\quad - \frac{15}{2576}h^2f_{n+6} \\
h\delta_{n+4} &= \frac{309}{56672}y_n - \frac{607}{11592}y_{n+1} + \frac{633}{2576}y_{n+2} - \frac{2823}{3220}y_{n+3} + \frac{845}{15456}y_{n+4} + \frac{51}{56}y_{n+5} - \frac{22912}{79695}y_{n+\frac{11}{2}} + \frac{9}{1288}h^2f_{n+6} \\
h\delta_{n+5} &= -\frac{117}{25760}y_n + \frac{475}{11592}y_{n+1} - \frac{1325}{7728}y_{n+2} + \frac{297}{644}y_{n+3} - \frac{16435}{15456}y_{n+4} - \frac{197}{840}y_{n+5} + \frac{1408}{1449}y_{n+\frac{11}{2}} - \frac{15}{1288}h^2f_{n+6} \\
h\delta_{n+6} &= \frac{42859}{1700160}y_n - \frac{1693}{7728}y_{n+1} + \frac{4449}{5152}y_{n+2} - \frac{40189}{19320}y_{n+3} + \frac{37519}{10304}y_{n+4} - \frac{3849}{560}y_{n+5} + \frac{123328}{26565}y_{n+\frac{11}{2}} + \\
&\quad \frac{801}{2576}h^2f_{n+6} \\
h\delta_{n+\frac{11}{2}} &= \frac{16209}{2072576}y_n - \frac{29315}{423936}y_{n+1} + \frac{184635}{659456}y_{n+2} - \frac{83259}{117760}y_{n+3} + \frac{258995}{188416}y_{n+4} - \frac{8217}{2048}y_{n+5} + \frac{498251}{159390}y_{n+\frac{11}{2}} + \\
&\quad \frac{1485}{47104}h^2f_{n+6}
\end{aligned}$$

Error Analysis and Zero Stability

Following Fatunla [18] and Lambert [20] we define the local truncation error associated with the conventional form of (5) to be the linear difference operator

$$L[y(x); h] = \sum_{j=0}^k \{\alpha_j y(x + jh)\} + \alpha_v y(x + vh) + h^2 \beta_v y''(x + jh) \quad (17)$$

Assuming that $y(x)$ is sufficiently differentiable, we can expand the terms in (17) as a Taylor series about the point x to obtain the expression

$$L[y(x); h] = C_0 y(x) + C_1 h y' + \dots + C_q h^q y^{(q)}(x) + \dots, \quad (18)$$

where the constant coefficients C_q , $q = 0, 1, \dots$ are given as follows: C_q , $q = 0, 1, \dots$

$$C_1 = \sum_{j=1}^k j\alpha_j,$$

•

According to Henrici [21], method (5) has order p if

Therefore, C_{p+2} is the error constant and $C_{p+2}h^{p+2}y^{(p+2)}(x_n)$ the principal local truncation error at the point x_n . It is established from our calculations that the HBDF have higher order and relatively small error constants as displayed in the Table 1.

$$\rho(R) = \det(RA^{(0)} - A^{(1)}) = R^k(R-1) \quad (19)$$

Case $k=6$. It is easily shown that (9)-(16) are normalized to give the first characteristic polynomial $\rho(R)$ given by

Where $A^{(0)}$ an identity matrix of is dimension seven and $A^{(1)}$ is a matrix of dimension seven given by

[illegible]

Following Fatunla [18] the block method by combining $k+1$ HBDF is zero-stable, since from (19),

$\rho(R)=0$ satisfy $|R_j| \leq 1$ $j=1, \dots, k$ and for those roots with $|R_j|=1$, the multiplicity does not exceed 2. The block method by combining $k+1$ HBDF is consistent since HBDF have order $P > 1$. According to Henrici [21], we can safely ascertain the convergence of HBDF method.

Results

We report here a numerical example taken from the literature.

Problem 1

$$y'' - y' = 0, \quad y(0) = 0, \quad y'(0) = -1, \quad h = 0.1$$

$$\text{Exact Solution } y(x) = 1 - e^x$$

Source: Mohammed [8]

Problems 2

$$y'' + y = 0, \quad y(0) = 1, \quad y'(0) = 1, \quad h = 0.1$$

$$\text{Exact Solution } y(x) = \cos(x) + \sin(x)$$

Source: Awari [22]

Problems 3

$$y'' = x(y')^2, \quad y(0) = 1, \quad y'(0) = \frac{1}{2}, \quad h = \frac{1}{30}$$

$$\text{Exact Solution } y(x) = 1 + \frac{1}{2} \ln \left(\frac{2+x}{2-x} \right)$$

Source: Badmus and Yahaya [23]

Table 1: Order and Error Constants For the HBDF methods

Step number	Method	order	Error constant
6	(9)	6	$\frac{801}{144256}$
	(10)	6	$\frac{3446421}{56}$
	(11)	6	$\frac{87945}{56}$
	(12)	6	$\frac{141471}{56}$
	(13)	6	$\frac{1378311}{56}$
	(14)	6	$\frac{179315}{6924288}$
	(15)	6	$\frac{31273}{288512}$

Table 2: Showing exact solutions and the computed results from the proposed methods for problem 1

x	Exact Solution	Proposed Method	Error in Proposed Method	Error in Mohammed [8]
0	0	0	1.4800000E-08	2.1980000E-05
0.1	-0.105170918	-0.1051709032	3.8100000E-08	6.0704000E-06
0.2	-0.221402758	-0.2214027199	6.2400000E-08	1.0051000E-05
0.3	-0.349858808	-0.3498587456	8.6200003E-08	1.4025300E-05
0.4	-0.491824698	-0.4918246118	1.1030000E-07	1.7993400E-05
0.5	-0.648721271	-0.6487211607	1.3360000E-07	2.1616200E-05
0.6	-0.822118800	-0.8221186664	1.5400000E-07	2.7993000E-05
0.7	-1.013752707	-1.013752553	1.8200000E-07	3.4561000E-05
0.8	-1.225540928	-1.225540746	2.1000000E-07	4.1114000E-05
0.9	-1.459603111	-1.459602901	2.3800000E-07	4.7656000E-05
1.0	-1.718281828	-1.718281590	1.4800000E-08	2.1980000E-05

Table 3: Showing exact solutions and the computed results from the proposed methods for problem 2

x	Exact Solution	Proposed Method	Error in Proposed Method	Error in Awari [22]
0	1	1	0.0000000E+00	0.0000E-00
0.1	1.094837582	1.094837655	7.300000E-08	1.1570E-07
0.2	1.178735909	1.178736102	1.9300000E-07	3.0990E-07
0.3	1.250856696	1.250857010	3.1400000E-07	5.0550E-07
0.4	1.310479336	1.310479768	4.3200000E-07	6.9570E-07
0.5	1.357008100	1.357008646	5.4599999E-07	8.7890E-07
0.6	1.389978088	1.389978742	6.5400002E-07	1.0540E-06
0.7	1.409059874	1.409060598	7.2400000E-07	1.0080E-06
0.8	1.414062800	1.414063636	8.3600000E-07	9.2260E-07
0.9	1.404936878	1.404937018	1.4000000E-07	8.2610E-07
1.0	1.38177329	1.381774327	1.0370000E-07	7.2160E-07

Table 4: Showing exact solutions and the computed results from the proposed methods for problem 3

X		Exact Value	Approx value	Present Error	Yahaya and Badmus [23]
0.1		1.050041729	1.050041724	5.00E-10	5.891E-06
0.2		1	1.100318692	1.67E-06	8.2399E-05
0.3		1.151140436	1.151028384	1.12E-05	3.46421E-04
0.4		1.202732554	1.202585545	1.47E-05	7.52101E-04
0.5		1.255412817	1.255265756	1.47E-05	1.380283E-03

Discussion

The HBDF methods are implemented as simultaneous numerical integration for IVPs without requiring starting values and predictors (see Tables 2,3 and 4). We proceed by explicitly obtaining initial conditions at x_{n+k} , $n=0,k,\dots,N-k$ using the computed values $Y(x_{n-k})=y_{n+k}$ and $\delta(x_{n-k})=\delta_{n+k}$ over sub-intervals $[x_0, x_k], \dots, [x_{N-K}, x_N]$ which do not overlap. We give examples to illustrate the efficiency of the methods.

We report here a numerical example taken from the literature.

Conclusion(s)

In this paper we developed a uniform order 1-block 6 –point integrators of orders (6,6,6,6,6,6) and the resultant numerical integrators posses the following desirable properties.

- (I) Zero-stability i.e stability at the origin
- (II) Facility to generate the solution at six point simultaneously
- (III) It is a convergence schemes

Hence, an improvement over other cited works.

References

1. Awoyemi D.O., *A class of continuous linear multistep method for general second order initial value problem in ordinary differential equation*, Intern. J. Comp. Math.,1999, 72, p.29–37.
2. Onumanyi P., Awoyemi D.O., Jator S.N., Sirisena U.W., *New linear Multistep with Continuous Coefficients for First Order Initial Value Problems*, J. Nig. Math. Soc.,1994, 13, p.37-51.
3. Mohammed U., Jiya M., Mohammed A.A., *A class of six step block method for solution of general second order ordinary differential equations*, Pacific Journal of Science and Technology, 2010, 11(2),p.273-277.
4. Yusuph Y., Onumayi P., *New Multiple FDMS through multi step collocation for $y'' = f(x, y)$* , Abacus, 2002, 29(2),p. 92-100.
5. Onumanyi P., Sirisena U.W and Jator S.N., *Continous Finite difference Approximation for Solving Differential Equations*, Inter.J. Comp Maths,1999, 72(1), p. 15-27.
6. Milne W.E., *Numerical Solution of Differential Equations*, John Wiley & Sons: New York, USA,1953
7. Rosser J.B., *A Runge-Kutta for all seasons*. Siam Rev, 1967,9(3), p.417–452.

8. Mohammed U., *A class of implicit five step block method for general second order ordinary differential equations*. *Journal of Nigerian Mathematical Society (JNMS)*, 30, p 25-39
9. Ibrahim Z.B., Othman K.I., Suleiman M., *Implicit r-point block backward differentiation formula for solving first-order stiff ODEs*, *Applied Mathematics and Computation*, 2007, 186, p.558–565.
10. Majid Z.A., Suleiman M.B., *Implementation of four-point fully implicit block method for solving ordinary differential equations*, *Applied Mathematics and Computation*, 2007, 184(2), p.514–522.
11. Akinfenwa O., Jator S., Yoa N., *An eighth order Backward Differentiation Formula with Continuous Coefficients for Stiff Ordinary Differential Equations*, *International Journal of Mathematical and Computer Sciences*, 2011, 7(4), p.171-176.
12. Akinfenwa O.A., Jator S.N., Yao N.M., *Continuous block backward differentiation formula for solving stiff ordinary differential equations*, *Computers and Mathematics with Applications*, 2013, 65, p.996–1005.
13. Yahaya Y.A., Mohammed U., *A Reformulation of Implicit Five Step Backward Differentiation Formulae in Continuous Form for Solution of First Order Initial Value Problem*, *Journal of General Studies*, 2009, 1(2), p. 134-144.
14. Yahaya Y.A., Mohammed U., *Fully Implicit Three Point Backward Differentiation Formulae for Solution of First Order Initial Value Problems*, *International Journal of Numerical Mathematics*, 2010, 5(3), p.384-398.
15. Mohammad U., Yahaya Y.A., *Fully Implicit four Point Block Backward Differentiation Formulae for Solution of First Order Initial Value Problems*, *Leonardo Journal of Sciences*, 2010, 16, p. 21-30.

16. Semenov D.E., Mohammed U., Semenov M.E., *Continuous multistep methods for solving first order ordinary differential equations*, Proceeding of Third Postgraduate Consortium International Workshop Innovations in Information and Communication Science and Technology IICST , Tomsk. Russia, 2013, p.165-170.
17. Van der Houwen P.J., Sommeijer B.P., Cong Nguyen Huu., *Stability of collocation-based Runge-Kutta-Nyström methods*, BIT, 1991,31, p.469–481.
18. Fatunla S.O., *Block Method for Second Order Differential Equation*. International Journal Computer Mathematics, 1991, 41, p. 55-63.
19. Jiaxiang X., Cameron I.T., *Numerical solution of DAE systems using block BDF methods*, Journal of Computation and Applied Mathematics, 1995, 62, p. 255-266.
20. Lambert J.D., *Computational Methods in Ordinary Differential Equations*, New York, John Wiley and Sons, 1973.
21. Henrici P.,*Discrete Variable Methods in Ordinary Differential Equations*. John Wiley & Sons: New York, USA.
22. Awari Y.S.,*Derivation and application of six point linear multistep numerical methods for solution of second order initial value problems*, ISOR Journal of Mathematics, 2013, 7(2), p.23–29.
23. Yahaya, Y.A and Badmus, A. M (2009): A class of Collocation Methods for General Second Order Differential equation. Africa Journal of Mathematical and Computer Science Research, **2(4)**, 69-71