

***Effect of L-arginine on the expression of endothelial
nitric oxide synthase (eNOS) in heart tissues of
diabetic rats***

Ahmed Mohamed M. Okasha, Hend Mohamed Abdel Ghany

Biochemistry Department – Faculty of Medicine, Minia University,
61511 Egypt.

***Correspondence to:**

Hend Mohamed Abdel Ghany

Lecturer, Department of Biochemistry, Faculty of Medicine, Minia University, Minia
61511, Egypt

Email: Mostafa.gedan@gmail.com

Abstract

Aims: This study aimed to determine the role of L-arginine on eNOS expression and on the level of some antioxidant as reduced glutathione and catalase in diabetes mellitus.

Study design: Ninety adult male albino rats were divided into 3 groups: Group (I): control group (30 rats). Group (II): Diabetic group (30 rats). Group (III): 30 rats were given L-arginine. Then the second and the third groups were made diabetic by intraperitoneal injection of 80 mg/kg alloxan monohydrate intraperitoneally at once. The animals were sacrificed and blood collected for determination of biochemical and antioxidant markers. L-arginine given orally for one week in dose 100 mg/kg in sterile (0.9%) NaCl before alloxan injection then L-arginine was given for another four

weeks. Heart tissues were homogenized for determination of expression of eNOS gene by RT-PCR.

Results: The study showed that L arginine increases HDL, reduces glutathione, catalase and decrease LDL, TAG, Cholesterol, MDA, also increase expression of eNOS in heart tissues of diabetic rat.

Key words: Antioxidant, Diabetes mellitus, e NOS gene, L-arginine.

Introduction

Diabetes mellitus is a serious metabolic disorder with micro and macrovascular complications that results in significant morbidity and mortality. The increasing size of the ageing population, consumption of a caloric rich diet, increasing rates of obesity and sedentary life style have led to an increase in the number of diabetic's worldwide [1].

Alloxan and streptozotocin selectively damage pancreatic β -cells and exhibit the most potent diabetogenicity. They have been widely used for induction of experimental diabetes mellitus type I. The mechanism increasingly accepted is the fragmentation of nuclear DNA of pancreas β -cells. Which seems to be important for the development of diabetes and is supposed to result from the accumulation of superoxide or hydroxyl radicals in the case of alloxan [2].

Oxidative stress is a constant feature of uncontrolled diabetes in humans and animals. Antioxidants such as vitamin C and E are decreased in diabetes, suggesting that oxidative stress in diabetes is at least in part, due to impaired antioxidant system, additionally reports from various areas show decrease activities of catalase and reduced glutathione [3].

L-Arginine is an α -amino acid. It was first isolated in 1886. In mammals, arginine is classified as a semiessential or conditionally essential amino acid, depending on the developmental stage and health status of the individual [3].

Nitric oxide (NO) is an important cellular signaling molecule involved in many physiological and pathological processes. It is a powerful vasodilator with a short half-life of a few seconds in the blood. In the presence of cardiovascular risk factors, endothelial dysfunction frequently is encountered. Several molecular defects could account for reductions in endothelium dependent vascular relaxation. Endothelial dysfunction could be due to decreased eNOS expression. However, several studies have shown that cardiovascular risk factors are associated with an increase rather than a decrease in eNOS expression. The increased expression of eNOS in vascular disease is likely to be a consequence of an excess production of H_2O_2 [4]. Nitric oxide is synthesized from L-arginine in the presence of oxygen and NADPH in a reaction catalyzed by NO synthase (NOS). Three NOS isoforms are expressed in mammalian cells: neuronal NOS (nNOS), endothelial NOS (eNOS), and inducible NOS (iNOS). The eNOS is constitutively expressed in endocardial cells, endothelial cells, ventricular myocytes, and other myocardial cells [5]. NO-mediate regulation of beating rate and vascular permeability, respectively. [6].

In diabetes, endothelial injury may be a consequence of glucometabolic and peroxidative stress. Experimental evidence suggests that the endothelium dependent relaxation is greatly reduced in the aorta of diabetic rats [6].

Materials and Methods

Experimental animals:

Ninety adult male healthy albino rats (obtained from the National Research Center, Giza, Egypt) weights (200 – 220 gm). The experiments were conducted according to the ethical norm approved by Institutional Animal Ethics Committee (IAEC). They were kept at the constant environmental condition for acclimatization, food and water were provided ad libitum for one week before the experiment.

Animals were given standard rat diet from (El- Nasr, cairo, Egypt) (protein 18 %, lipid 2.7 %, fibers 10.06 % and energy 2600 K cal) and tap water.

Animals were divided into three groups:

1. Group I: (N = 30) animals served as control group.
2. Group II (Diabetic group): (N = 30). Animals served as positive control that were starved over night before alloxan injection.
3. Group III (Treated group): (N = 30). Animals were given L-arginine orally for one week in dose 100 mg/kg in sterile (0.9%) NaCl before alloxan injection then L-arginine was given for another four weeks.

After that, the second group and the third group were injected by alloxan monohydrate in dose 80 mg/kg in sterile (0.9%) NaCl intraperitoneally (I.P.) at once. Alloxan has led to type I diabetes mellitus. After 72 hour of alloxan injection, blood samples were collected from the tail vein of rats and blood glucose was determined by one touch method.

At the end of the experiment, animals were scarified and the blood was collected and centrifuged to obtain serum which collected for determination of the following parameters (glucose, serum triglycerides, cholesterol, HDL-cholesterol, LDL-cholesterol, plasma catalase, reduced glutathione and lipid peroxide by colorimetric method according manufacture instruction. Total RNA was extracted

by Ribozol reagent from heart tissues for determination of expression of eNOS gene by RT-PCR.

Drugs, chemicals, and kits; Alloxan (Biotechnology Co., Egypt).

- Catalase, reduced glutathione, cholesterol, triglycerides, HDL, LDL and malondialdehyde kit (Biodiagnostic Co., Egypt).
- Glucose kit (BioMed Diagnostic Co., Egypt).
- L-arginine (Sigma-Aldrich Co., Egypt).

Detection of the level of NO

Nitric oxide colorimetric assay kit (cat No K262-200). Bio Vision (USA) according to manufacture instruction. Nitric Oxide Colorimetric Assay Kit provides an accurate, convenient measure of total nitrate/nitrite in a simple two-step process. The first step converts nitrate to nitrite utilizing nitrate reductase. The second step uses Griess. Reagents to convert nitrite to a deep purple azo compound. The amount of the azochromophore accurately reflects nitric oxide amount in samples. The detection limit of the assay is approximately 0.1 nmol nitrite/well, or 1 μ M. Calculations: Absorbance was plotted at 540 nm as a function of nitrate and/or nitrite concentration (It is possible to use plate readers with filters rather far from 540 nm; e.g. 570 nm).

$$\text{Nitrate/nitrite concentration} = \left[\frac{\text{Abs sample} - \text{blank Abs}}{\text{Slope of Std curve}} \right] / \mu\text{L of sample}$$

mRNA analysis (rt-PCR).

Total RNA was extracted from homogenates of heart tissue using RiboZol solution (Amresco, Solon, USA) according to the manufacturer's instructions. RNA samples were stored at -80°C until analysis.

RT-PCR was performed according to manufacture instructions (Qiagen one step). Reaction tube of RT-PCR was containing 5 µl of RT-PCR buffer 5× 10 pM of specific primers for eNOS or GADPH, 5 µg of total RNA template, 1 µl dNTPs mix (100 mM), 1 µl enzyme mix (Hot Star Taq DNA polymerase, Omniscript and Sensiscript reverse transcriptases).

Polymerase chain reaction (PCR) amplification was performed at the following temperature steps: 5 min, 95 °C; 94 °C for 20 sec; 55 °C for 30 sec; 72 °C for 40 sec (30 cycles); and 72 °C for 5 min. The MgCl₂ concentration used for eNOS cDNA amplification was 2.0 mM. The following primer pairs were used: 5'-CGA GAT ATC TTC AGT CCC AAG C-3' (sense) and 5'-GTG GAT TTG CTG CTC TCT AGG-3' (antisense) for rat eNOS. PCR was performed with rat GADPH as an internal standard (Thermo scientific) USA .Sense5'-CAGTGCCAGCCTCGTCTCAT-3', antisense 5'-GGGGCCATCCACA GTCTTC-3', sample for rat GADPH has same condition for rat eNOS except the annealing temperature was 54 °C, in a thermal cycler (Progene, Techne (Cambridge) LTD., UK).

Detection of PCR product.

Polymerase chain reaction products (5 µl) were detected by electrophoresis on a 1.5 % agarose gel containing ethidium bromide. Location of a predicted product was confirmed by using 100 bp ladder (Gene Ruler, Thermo scientific, USA)) as a

standard size marker. The negative control tubes containing all component as in RT-PCR reaction tubes except it was not containing total RNA template.

Statistical analysis

The data were expressed with SPSS version 13 software as the mean $\pm$ standard error (SE), analysis of variance (ANOVA) indicated a significant at ($P < 0.05$). Independent t-test for parametric quantitative data.

The density of PCR product was measured using Scion Image J software (Scio Cooperation, Fredrick, Maryland) and expressed as mean $\pm$ SEM

Results

The result shows that L-arginine reduces blood glucose level by 49% from 266.7 mg/dl in alloxan group to 135.1 mg/dl in treated group (Table 1). Also the result shows that L-arginine reduces the triglyceride level by 25.5% from 200.8 mg/dl in group I to 149.6 mg/dl in treated group and LDL level by 22% from 210.2 mg/dl in group I to 164.4 mg/dl in treated group. Finally the result shows that L-arginine improves HDL level by 97% from 17.5 mg/dl in group I to 34.5 mg/dl in treated group (Table 1). L-arginine decrease cholesterol level from 291.2 mg/dl in group I to 241.7 mg/dl in treated group which means L-arginine reduced triglyceride level by 17%.

The result shows that L-arginine increase the reduced glutathione level by 80% from 0.5 mg/dl in alloxan group to 0.9 mg/dl in treated group (Table 1 and Fig 1).

The result shows that L-arginine reduces the MDA level by 47% from 56.9 nmol/ml in alloxan group to 30.3 nmol/ml in treated group. (Table 1). Also there was significant decrease in cardiac NO level in diabetic group (21 ± 9 nmol/mg protein)

and significant increase in treated group to become (25 ± 6) at P value <0.001 as compared with control group.

Also there was negative correlation between catalase, reduced glutathione and blood glucose (Fig. 2, 3)

Fig. (4) Shows mRNA eNOS RT-PCR analysis. The band of mRNA for eNOS in the diabetic heart (Group II) appears significantly decreased in comparison to that in the control lane (with GAPDH as internal standard). Also there was significant increase in expression in treated group (III) as compared with group I.

Table (1): Shows the average values of each group separately.

	Group I (control)		Group II (Diabetic)		Group III (Treated)		P value
	Mean $\pm$	SE	Mean $\pm$	SE	Mean $\pm$	SE	
Glucose (mg/dl)	94.2 $\pm$	2.6	266.7 $\pm$	7.6	135.1 $\pm$	1.4	<0.001
TG (mg/dl)	3.4 $\pm$	5.6	200.8 $\pm$	3.9	49.6 $\pm$	1.8	<0.001
LDL (mg/dl)	81.9 $\pm$	4.3	210.2 $\pm$	3.4	120.4 $\pm$	2.03	<0.001
HDL (mg/dl)	53.9 $\pm$	2.8	17.5 $\pm$	4.9	34.5 $\pm$	1.2	<0.001
CHL (mg/dl)	177.8 $\pm$	3.7	291.2 $\pm$	2.4	241.7 $\pm$	1.4	<0.001
Glutatione (mg/dl)	1.2 $\pm$	0.01	0.5 $\pm$	0.01	0.9 $\pm$	0.01	<0.001
Catalase (U/l)	28.4 $\pm$	0.5	11.9 $\pm$	0.3	18.5 $\pm$	0.3	<0.001
MDA (nmol/ml)	56.2 $\pm$	0.2	56.9 $\pm$	0.4	30.3 $\pm$	0.3	<0.001
Cardiac NO (nmol/mg protein)	43 $\pm$ 18		21 $\pm$ 9		25 $\pm$ 6		<0.001




Figure (1): Positive correlation between MDA and blood glucose




Figure (2): Negative correlation between catalase and blood glucose




Figure (3): Show negative correlation between reduced glutathione and blood glucose.




Figure 4: Agarose gel electrophoresis of RT-PCR product showing rat eNOS expression (135 bp) in heart tissue of different groups.

Lane 1: DNA ladder (100 bp), lane 2: negative control, lane 3: control group, Lane 4: group II (diabetic group). Lane 5: group III (L-arginine treated group), with GADPH as internal standard at 595 bp

Discussion

This study reveals a number of important findings on the relationship between diabetes and the consumption of L-arginine. Poorly controlled long standing metabolic disorders diabetes mellitus is frequently results in cardiovascular complications [9].

L-arginine administration reduces lipid peroxide levels in diabetic patients, due to direct scavenger of superoxide anions. L-arginine might enhance relaxation by

increasing NO release, which in turn scavenges enhanced rates of superoxide anion production that are elevated in diabetic arteries [10].

The results which were obtained showed that treatment of diabetic rats with the L-arginine leads to significant reduction in blood glucose level and this could be attributed to its effect on the enzymes that digest carbohydrates and hence decrease glucose absorbance from intestine and these data were in line agreement with Broca et al 2004 [11].

The level of MDA was significantly reduced which indicate that the reduction in the lipid preoxidation because of the treatment. In addition, the antioxidant enzymes, catalase and reduced glutathione were significantly increased as result of the treatment indicating the reduction of oxidative stress and ROS in the treated groups which may be attributed an antioxidation effect of the L-arginine [12].

Lipid profile results show high efficacy of L-arginine in reducing triglyceride and LDL levels and improving HDL level and these data were in line with Fu et al 2005 [13].

Antioxidant enzymes as reduced glutathione and catalase were improved by L-arginine according to result. L-arginine increase reduced glutathione. from 0.5 mg/dl in diabetic group to 0.9 mg/dl in treated group, which means that after treatment by L-arginine reduced glutathione improved by 80% and these data were in line with Li et al 1994 [14].

L-arginine increase catalase by 55.5% from 11.9 U/L in diabetic group to 18.5 U/L in treated group and these data were in line with Li et al 1994 [14].

A defect in nitric oxide function develops very early in diabetes mellitus and leads to altered vascular reactivity [15]. The association of diabetes mellitus with

atherosclerotic cardiovascular disease may be a consequence, in part of the decrease of eNOS protein in endothelial cells.

NO has important cardiovascular effects and modulates cardiac function. The contraction of cardiac muscle responds to chemical transmitters, oxygen tension and physical factors by modifying transmembrane calcium currents, intra-cellular calcium storage and the contractile proteins [16].

Reports indicate that diabetes induces alterations of calcium handling and impairs nitric oxide production in coronary endothelial cells, due to tetrahydrobiopterin deficiency and in the contractile proteins that limit myocardial function independent of altered metabolism [17].

Although dysfunction of the NO pathway may promote atherogenesis, NO supplied exogenously to activated endothelium down regulates pro-inflammatory activity such as chemokine secretion and functional activity such as transendothelial migration of neutrophils [18]

:[1R]Comment

Conclusion

The study show that L arginine increase HDL, reduced glutathione, catalase and decrease LDL, TAG, Cholesterol, MDA, also increase expression of eNOS in heart tissues of diabetic rat.




Figure 5: eNOS expression in heart tissues in different group (show significant decrease in diabetic group as compared to control group at P value <0.001, and significant increase in treated group as compared with diabetic group at P value <0.001).

References

1. Wild S, Roglic G, Green A, Sicree R. and King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004; 27 (5): 1047–53.

2. Renold AE. Streptozotocin and alloxan in experimental diabetes: Comparison of the two models in rats **Diabetes Annu.** 1988; 4: 592-608.
3. Maritim AC, Sanders RA, and Watkins JB. Diabetes, oxidative stress and antioxidant: a review. *J Biochem Mol Toxicol* 2003; 17: 24 – 38.
4. Tapiero. L-Arginine". *Biomedicine and Pharmacotherapy* 2002; 56 (9): 439–445.
5. Förstermann U, and Münzel T. Endothelial nitric oxide synthase in vascular disease: *Circulation.* 2006; 113:1708-1714.
6. Balligand JL, and Cannon PJ. (Nitric oxide synthases and cardiac muscle. Autocrine and paracrine influences. *Arterioscler Thromb Vasc Biol* 1997; 17:1846-1858.
7. Sikorski BW, Hodgson WC, and King RG. Effect of haemoglobin and N-nitro-L-arginine on constrictor and dilator response of aortic rings from streptozotocin diabetic rats. *Eur J Pharmacol* 1993; 242:275-282.
8. Felaco M, Grilli A, Gorbunov N, Di Napoli P, De Lutiis MA, Di Giulio C, and Conti P. Endothelial NOS expression and ischemia-reperfusion in isolated working rat heart from hypoxic and hyperoxic conditions. *Biochim Biophys Acta* 2000; 1524:203-211.
9. Innis M, Gelfand D, and Sninsky J. PCR Protocols, a Guide to methods and applications , *Academic Press, San Diego* 1990; pp 1-482.

10. Ritz E, Fliser D and Nowicki M. Hypertension and vascular disease as complications of diabetes. *laragh JV, Bernner BM eds. New York, NY: Raven press Ltd* 1995; 2321 – 2334.
11. Caballero F, Gerez E, Batlle A and Vasquez. Preventive aspirin treatment of streptozotocin induced diabetes: blockage of oxidative status and prevention of heme enzymes inhibition. *Chem biol Interact* 2000; 126:215-225.
12. Broca C, Breil V and Cruciani GC. Aminoacid Support in the prevention of diabetes and diabetic complications: *Am. J. physiol. Endocrinol .Metab* 2004; 287:463.
13. Vaziri ND (2002): Pathogenesis of led – induced hypertension: role of oxidative stress. *J Hypertens Supp1*; 3: S15 – S20.
14. Fu WJ, Haynes TE, Kohli R, Hu J, Shi W, Spencer TE, and Meininger CJ. Dietary L-arginine supplementation reduces fat mass in Zucker diabetic fatty rats. *J Nutr* 2005; 135(4):714-21.
15. Li R, Sundran R, Joseph W., Kerrie D. and Chris R, (1994): Mutation of an arginine residue in the human glycine receptor transforms β -alanine and taurine from agonists into competitive antagonists. *Neuron*. 14, p169–175
16. Chan NN, Vallance P, and Colhoun HM. Nitric oxide and vascular responses in Type I diabetes. *Diabetologia* 2000; 43:137-147.
17. Mohan P, Brutsaert DL, and Sys SU. Myocardial performance is modulated by interaction of cardiac endothelium derived nitric oxide and prostaglandins. *Cardiovasc. Res* 1995; 29:637-640

18. Meininger CJ, Marinos RS, Hatakeiama K, Martinez and Wu G. Impaired nitric oxide production in coronary endothelial cells of the spontaneously diabetic BB rat is due to tetrahydrobiopterin deficiency. *Biochem* 2000; J; 349:353-356.
19. Fowler AA III, Fisher BJ, Sweeney LB, Wallace TJ, and Ghosh S. Nitric oxide regulates interleukin 8-gene expression in activated endothelium by inhibiting NF-kappa B binding to DNA: effects on endothelial function. *Biochem Cell Biol* 1999; 77:201-208.