

Original Research Article**Evaluation of the effects of methanolic extract of alligator pepper seed on serum electrolyte and renal function markers of rats.****Abstract**

The effect of methanolic extract of *Aframomum melegueta* (AME) on the serum creatinine, blood urea nitrogen (BUN) and electrolytes were investigated in albino rats. The extraction was done by cold maceration of pulverized seeds of alligator pepper in absolute methanol for 48 hours. The extract was filtered and concentrated *in vacuo* in a rotary evaporator at 40 °C. Twenty male albino Wistar rats were randomly divided into four groups of five animals each. Group A served as the control and received 10 ml/kg of 5 % Tween 20 while Group B-D received 100, 200 and 400 mg/kg respectively, of the AME. The animals were dosed daily for 21 days. Twenty four hours later, blood was collected from the rats through direct cardiac puncture. The separated serum was used to determine the serum creatinine, BUN and electrolyte assay. The extract produced no significant ($p > 0.05$) difference in mean serum creatinine, BUN and electrolytes level in the treated rats when compared to the control group rats. In conclusion the extract did not exhibit any adverse effect on the kidney at the doses used and for the period of experiment.

Keywords:- *Aframomum melegueta*, creatinine, Blood urea nitrogen, electrolyte

1. INTRODUCTION

Plants have provided a source of inspiration for novel drug compounds as plant derived medicines have made large contribution to human health and well being. Despite the achievements recorded in drug discovery and development from plant sources, phytomedicine continues to be highly valuable for developing synthetic drugs [1, 2]. According to the World Health Organization (WHO), about 80% of the world's population depends wholly or partly on pharmaceuticals of plant origin [3]. Thus, the need for toxicity evaluation of these medicinal plants, cannot be underscored [4], such research could advance the

utilization of these indigenous herbal medicine for orthodox treatment [5] or to serve as new dietary supplements which consist of minerals that may be used for the treatment of organ related diseases. Kidney is the major organ of excretion of drug and its metabolite from the system. Some of the toxic effect of drugs manifest clinically in the kidney by either enhanced or reduced reabsorption of electrolyte by the kidney tubules [6].

Aframomum melegueta Schum (Zingiberaceae) also known as alligator pepper (indigenous names include: Atare in Yoruba, Ose-oji in Igbo and Citta in Hausa) is used in Nigeria and some other parts of West Africa, as a spicy during entertainment and have a wide range of folkloric uses in traditional medicine. They are used as a remedy for treating stomach ache, diarrhoea and snakebite [7, 8]. The seed extract have been evaluated for antinociceptive, anti-ulcer, anti-microbial, anti-inflammatory, anti-oxidant and sexual performance enhancing activities [7-11]. There is paucity of information on the effect on the kidney tubule. The present study was aimed at evaluation of the serum electrolyte and renal function markers of methanolic extract of alligator pepper seed.

2. Materials and methods

2.1. Plant

The freshly harvested fruit of *Aframomum melegueta* (Alligator Pepper) were bought from Ngoro market, Ogoro in Ikwo LGA of Abia State in the month of July 2013 and were authenticated by Dr. I. C. Okwulehie of the Department of Plant Science and Biotechnology, Michael Okpara University of Agriculture, Umudike and the voucher specimen catalogued MOUAU/CVM/VPP/2013/03 was kept in the departmental herbarium for reference purpose.

2.2. Preparation of the extract

The dehaulled, dried and pulverized seeds of *Aframomum melegueta* were extracted by cold maceration method for 48 hours at room temperature using absolute methanol in a Winchester bottle. The *Aframomum melegueta* extract (AME) was filtered with Whatmann No 1 filter paper. The filtrate was concentrated *in-vacuo* using vacuum rotary evaporator at 40 °C and was later concentrated to dryness in a hot air oven at 40 °C. The extract was stored in a refrigerator at 4 °C throughout the duration of this study.

2.3. Determination of the yield of the AME

An empty clean and dry beaker was weighed and later the extract was poured into it. The beaker was weighed after the extract has been concentrated to constant weight. The weight of the extract was calculated as follow:

$$\text{The percentage yield of extract (\% w/w)} = \frac{\text{weight of beaker and extract} - \text{weight of empty beaker}}{\text{weight of plant material}} \times \frac{100}{1}$$

2.4. Animals

Twenty four male Wistar albino rats weighing between 120 - 170 g were obtained from Department of Zoology, University of Nigeria, Nsukka and kept in the Animal house of the Biochemistry Department, Michael Okpara University of Agriculture, Umudike. The animals were allowed access to feed and water *ad libitum* and were allowed two weeks of acclimatization before the commencement of the experiment. The animals were kept in well ventilated aluminium cages at room temperature and under natural light/darkness cycles. They were maintained in accordance with the recommendation of the *Guide for the care and use of laboratory animals* [12]. The experiment was approved by the University Animal Ethics Committee with ref. MOUAU/CVM/EAEC/2013/201.

2.5. Phytochemical spot test

The AME was tested for the presence of alkaloids, flavonoids, tannins, glycosides, saponins, terpenes/sterols, carbohydrates, and starch using the standard procedures as described by Trease and Evans [13].

2.6. Experimental design

Twenty male albino Wistar rats were randomly divided into four groups of five animals each. Group A served as the control and received 10 ml/kg of 5 % Tween 20 while Group B-D received 100, 200 and 400 mg/kg respectively, of the AME. The animals were dosed daily for 21 days and were observed daily for changes and other signs of toxicity and death throughout the period of study. Twenty four hours after the last treatment blood obtained through direct cardiac puncture was used to assay for effects of AME on serum electrolyte, urea and creatinine level.

2.7. Body weight and body weight gain

Individual body weights were recorded at study initiation (day 0), and weekly thereafter. Mean body weight gains were calculated for each group at each interval and for the overall (days 0–21) testing interval. Animals were also weighed immediately prior to killing (fasted body weight) for calculation of relative organ weight.

2.8. Biochemical analysis

After 21 days of extract administration the rats were fasted overnight, only distilled water was made available *ad libitum*. Blood samples were collected through direct cardiac puncture under chloroform anesthesia using a 5 ml syringe into plain sample bottles. The blood sample was centrifuged at 4000 rpm for 20 mins in order to separate the serum. The clean sera were aspirated and stored frozen for serum biochemical analysis. Serum electrolytes (Na^+ , K^+ , Cl^- , HCO_3^-), urea and creatinine were determined using standard ready-to-use kits (Randox Ltd, UK) following the manufacturer's instructions.

2.9. Statistical analysis

Data obtained were analyzed using one-way analysis of variance (ANOVA) and the variant means were separated by least significant difference (LSD) of the different groups. Significance was accepted at the level of $p < 0.05$.

3. RESULTS AND DISCUSSION

3.1 Results

The methanolic extraction of the dried seed of *A. melegueta* yielded 8.34 % w/w dry extract which was oily and dark-brown in colour.

The phytochemical spot test showed that the extract contained saponins, tannins, terpenes/sterols, glycosides, alkaloids, flavonoids.

The result of the effect of AME treatment on the body weight and percentage weight gain is presented in Table 1. The extract did not produce any significant ($p > 0.05$) difference in the body weight and percentage weight gain in the treated rats, when compared to the negative control group rats.

The result of the relative organ (kidney) weight of AME treated rats is presented in Figure 1. The extract did not produce any significant ($p > 0.05$) difference in the relative kidney weight of treated rats, when compared to the negative control group rats.

The effect of AME on the mean serum creatinine, urea and electrolytes of the treated rats is presented in Table 2 and 3. The extract did not produce any significant ($p > 0.05$) difference in the mean serum creatinine, urea and electrolytes of treated rats, when compared to the negative control group rats.

Table 1. Effect of AME on the relative kidney weight

Group	Mean body weight in gram $\pm$ SEM (percentage increase in weight compared to day 0)			
	Day 0	Day 7	Day 14	Day 21
5 % Tween 20, 10 ml/kg	137.14 $\pm$ 4.09	153.96 $\pm$ 1.41 (11.74)	169.12 $\pm$ 2.05 (15.32)	174.02 $\pm$ 2.80 (17.49)
AME 100 mg/kg	124.24 $\pm$ 11.37	140.13 $\pm$ 12.50 (12.14)	150.94 $\pm$ 12.16 (21.33)	159.54 $\pm$ 13.31 (28.63)
AME 200 mg/kg	128.38 $\pm$ 6.61	142.52 $\pm$ 5.54 (11.57)	143.94 $\pm$ 5.07 (11.71)	146.18 $\pm$ 4.66 (16.54)
AME 400 mg/kg	139.38 $\pm$ 10.14	151.15 $\pm$ 10.71 (8.92)	155.58 $\pm$ 9.14 (10.22)	160.55 $\pm$ 7.36 (15.98)

No statistical difference ($P > 0.05$) compared with control group

Table 2. The effect of AME on the kidney function test of the treated rats

Group	Urea (mg/dl)	Creatinine (mg/dl)
5 % Tween 20, 10 ml/kg	15.96 $\pm$ 3.72	0.56 $\pm$ 0.08
AME 100 mg/kg	19.38 $\pm$ 4.60	0.20 $\pm$ 0.08
AME 200 mg/kg	27.22 $\pm$ 2.73	0.92 $\pm$ 0.28
AME 400 mg/kg	21.50 $\pm$ 3.17	0.51 $\pm$ 0.05


No statistical difference ($P > 0.05$) compared with control group

Table 3. The effect of AME on the mean serum electrolyte of the treated rats

Group	Sodium (mEq/L)	Potassium (mEq/L)	Chloride (mEq/L)	Carbonate (mmol/l)
5 % Tween 20, 10 ml/kg	111.01 $\pm$ 28.64	9.89 $\pm$ 4.43	166.10 $\pm$ 6.85	36.73 $\pm$ 1.02
AME 100 mg/kg	98.56 $\pm$ 20.18	3.00 $\pm$ 2.04	167.23 $\pm$ 26.88	33.98 $\pm$ 3.19

AME 200 mg/kg	109.93 ± 28.65	8.78 ± 1.90	145.76 ± 27.56	33.19 ± 4.66
AME 400 mg/kg	114.44 ± 11.85	6.11 ± 2.30	162.71 ± 6.85	36.55 ± 2.00

127 No statistical difference ($P > 0.05$) compared with control group



128

129

130 Figure 1. Effect of AME on the relative kidney weight

131

132

133 3.2 Discussion

134 Kidney is the main organ by which drug and their metabolites are eliminated from the body. It is the target
135 for drug toxicity. The main function of the kidney is to maintain the constancy of the “interior environment”
136 by eliminating waste products and by regulating volume, electrolyte content and pH of the extracellular
137 fluid in the face of varying environmental demand [6].

138 The present work evaluated the subacute effect of AME on the serum electrolyte, creatinine and urea
139 level in treated rats. The doses used in this study were based on report of previous studies on
140 *Aframomum melegueta* [7-11]. The result of the experiment showed that the extract has no adverse effect
141 on kidney at the doses used and for the period of the experiment, when compared to the negative control.
142 Increasing the dose may elicit some adverse effects. Toxicity depends on the dose of the intoxicant [14].
143 The serum creatinine, blood urea nitrogen and electrolyte are used to evaluate the functional capacity of
144 the kidney [15]. The result of the effect of AME on serum blood urea nitrogen level of the treated rats may

suggest that the extract has no the kidney function and protein catabolism. Creatinine is metabolic waste product of creatine and creatine phosphate. Its endogenous production is constant and directly proportional muscle mass of the individual. Its serum level is an index of renal function and measures the glomerular filtration rate. Creatinine is readily filtered and does not undergo any significant tubular reabsorption. The serum level of creatinine increases when the formation or excretion of urine is impaired [16]. In this present study there was no significant ($p > 0.05$) difference in the body weight of all the AME treated groups when compared to the control group, which indicate that the muscle mass were not affected.

The result of the effect of AME on serum blood urea nitrogen level of the treated rats may suggest that the extract has no effect on the kidney function and protein catabolism. Urea is the major nitrogen-containing metabolic waste product of protein catabolism in mammals. It formed from exogenous or endogenous protein from the breakdown of cell in the body [16]. More than 90% of the urea is excreted by the kidneys, and the remainder is lost through the gastrointestinal tract and skin. The measurement of BUN is used to evaluate renal function [15].

In the present study the extract produced no significant ($p > 0.05$) difference in all the serum electrolyte levels measured when compared to the negative control. This may suggest that the extract may not have any electrolyte reabsorption, secretion and diuretic effect in the kidney tubule. Diuretic agent (furosemide) act by inhibiting sodium chloride and water reabsorption at the loop of Henle of the nephron, which may lead to hyponatraemia or hyperkalcaemia [17].

In conclusion, the extract did not exhibit any adverse effect on the kidney at the doses used and for the period of experiment, but may exhibit toxic effect at higher dose and/or more prolonged period treatment.

CONSENT

Not applicable.

Conflict of interest statement

We declare that we have no conflict of interest.

Reference

1. Adeneye AA, Benebo AS. Pharmacological evaluation of a Nigerian polyherbal health tonic tea in rat. African Journal of Biomedical Research. 2007;10(3):249-255.
2. Ogunka-Nnoka CU, Uwakwe AA, Nnabuike CJ. Effects of Ethanolic /Potash Extract of Sorghum Bicolor Leaf Sheath on Serum Electrolytes, Liver and Kidney Indicative on Albino Rats. Journal of Natural Sciences Research. 2012;2(4):66-70.
3. World Health Organization. Guidelines for the Assessment of Herbal medicines. WHO expert committee on specification for pharmaceutical preparations. Technical Report series. No. 863 Geneva, 1996.
4. Jarnovska D, Kubikova K, Kokoska L. Screening for Antimicrobial Activity of some Medicinal Plant Species of Traditional Chinese Medicine. Czechoslovakian Journal of food Science. 2003;21:107-111.
5. Parek J, Chanda S. In vitro Antimicrobial activities of Extract of Launarea procumbens Roxb (Labiatae), vitis vinifera L. (vitaceae) and cyprenus rotundus L. (cyperaceae). African Journal of Biomedical Research. 2006;9:89-93.
6. Rang HP, Dale MM, Ritter JM, Moore RK. Pharmacology 6th edition, Edinburgh: Churchill livingstone; 2007.
7. Umukoro S, Ashorobi RB. Further studies on the antinociceptive action of aqueous seed extract of *Aframomum melegueta*. Journal of Ethnopharmacology, 2007;109:501–504.
8. Ilic N, Schmidt BM, Poulev A, Raskin I. Toxicological evaluation of grain of paradise (*Aframomum melegueta*) [Roscoe] K. Schum. Journal of Ethnopharmacology. 2010;127:352-356.
9. Galal AM. Anti-microbial activity of 6-paradol and related compounds. International Journal of Pharmacognosy. 1996;31:64–69.
10. Kamtchouing P, Mbonge GYF, Dimo T, Watcho P, Jatsa HB, Sokeng SD. Effects of *Aframomum melegueta* and *Piper guineense* on sexual behaviour of male rats. Behavioural Pharmacology. 2002;13:243–247.

- 199 11. Onoja SO, Omeh YN, Ezeja MI, Chukwu MN. Evaluation of the In Vitro and In Vivo Antioxidant
200 Potentials of *Aframomum melegueta* Methanolic Seed Extract. Journal of Tropical Medicine. 2014;
201 Article ID 159343, 6 pages.
- 202 12. DHHS, Guide for the Care and Use of Laboratory Animals, Institute of Laboratory Animal Resources
203 Commission on life Sciences, National Research Council, National Academy Press Washington,
204 D.C., 1985.
- 205 13 Evans WC. Trease and Evans Pharmacognosy. 16th ed. Edinburgh: W. B. saunders Elsevier, 2009;
206 16:58-302.
- 207 14. Braide VB, Anika SM. Environmental toxicology. 1st ed. Enugu: SNAAP press Ltd; 2007.
- 208 15. Sunheimer RL, Graves L. Clinical Laboratory Chemistry. New York: Pearson Education. 2011;154.
- 209 16. Brar RS, Sandhu HS, Singh A. Veterinary Clinical Diagnosis by Laboratory Methods. New Delhi:
210 Kalyani Publishers.2011;93-97.
- 211 17. Asuzu IU, Anika SM. Textbook of Veterinary Pharmacology. Enugu: Fourth Dimension Publishers.
212 2012;178-189.
- 213
- 214
- 215
- 216
- 217
- 218
- 219
- 220
- 221
- 222
- 223
- 224