

Evaluation of the changes in some liver function and haematological parameters in MSG fed rats.

ABSTRACT

The effect of Monosodium glutamate (MSG) was evaluated on hepatic functions and haematological parameters in adult Wistar albino rats. The rats were randomly assigned into four (4) groups of five rats each. Group 1 served as the control while groups 2, 3 and 4 were fed diets supplemented with MSG at doses of 0.5, 1.0 and 5.0% per Kg of diet respectively. Following the 28 days of feeding, the rats were sacrificed and blood samples were collected for analyses. The haematological parameters except PCV were estimated using haematocytometer while the biochemical parameters were determined using Randox enzymatic kit. There was no significant difference ($p>0.05$) in PCV, WBC, RBC, total protein, albumin, direct and conjugated bilirubin in the rats fed with MSG at all levels of supplementation when compared with the control group. The Alkaline phosphatase (ALP) and Alanine aminotransferase (ALT) activities were increased with increase level of MSG and ranged from 35.00 ± 0.17 - 59.00 ± 1.23 U/L and 6.4 - 11.9 U/L respectively. There was no significant ($p>0.05$) difference in the levels of both serum ALP and ALT activities in the group fed 0.5% MSG when compared with the control group while there were significant ($p>0.05$) increase in the other treatment groups (1 and 5% MSG supplemented diets groups). The results therefore suggest that MSG at the levels of supplementation in the diets of the rats had no effect on the haematological indices but increased the liver function enzymes in the serum as the level of MSG increased in the diet.

Key word: Packed Cell Volume, Red Blood Cell, White Blood Cell, Monosodium Glutamate (MSG)

INTRODUCTION

It is a common practice worldwide to improve the taste and flavour of food by means of food additives. Monosodium glutamate is an additive commonly used in flavouring and enhancing the taste of foods (Oyetunji, 2013). Monosodium glutamate (MSG) is sodium salt of naturally occurring L-glutamate which contains about 78% glutamic acid and 22% sodium and water. Glutamate is thus found in a wide variety of foods, especially high protein foods such as dairy products, meat, fish and many vegetables. Foods often used for their flavouring properties, such as mushrooms and tomatoes, have high levels of naturally occurring glutamate (Alao *et al.*, 2010). MSG is produced commercially by fermentation of molasses and other substrates suitable for the growth of *Clostridium glutamicum*. Monosodium glutamate is marketed in Nigeria as the popular Ajinomoto. It is used as flavor enhancer at low concentration of (0.1% - 0.8%) because high dose render the taste of the food unpleasant (Manal and Nawal, 2012). The unique flavor and taste of this compound has been categorized and established as a separate taste sensation “unami” taste (Ikeda, 1909). Industrial food manufacturers make use of monosodium glutamate as a flavor enhancer because it balances, blends and rounds the total perception of other tastes (Loliger, 2000).

Monosodium glutamate can be used to reduce salt intake (sodium), which predisposes to hypertension, heart diseases and stroke (Legetic and Campbell, 2012). This is because of its flavour enhancing property which results from its interaction with sodium chloride salt, and other “unami”

substances such as nucleotides. Even with a 30% salt reduction, the taste of low-salt foods improves with MSG inclusion. The sodium content of MSG (12%) is roughly a third of the amount of that in sodium chloride (39%). Other salts of glutamate have been used in low-salt soups, but with a lower palatability than MSG (Ball *et al.*, 2002). However, when enhancing palatability it is important to ensure that the food additive will not cause harm to the consumer. In recent years there have been much concern about possible adverse effect of MSG on human and animals and MSG is believed to cause neurotoxicity (Singh *et al.*, 2003), obesity (Gobatto *et al.*, 2012), impaired vision and endocrinal disorder in human subjects. It has also been associated with some pathological conditions such as schizophrenia, anxiety, Parkinson's disease, epilepsy, stroke and Huntington's disease (Mozes and Sefcikova, 2004). The metabolism of MSG occurs basically in the liver (the largest organs in the body). The liver is an organ which plays an important metabolic role such as detoxification of xenobiotic, transamination, removal of ammonia in form of urea, biosynthesis and release of non essential amino acid in the body.

Enzymes such as Alanine aminotransaminase (ALT) and Aspartate aminotransaminase (AST), Lactate dehydrogenase (LDH), Alkaline phosphatase and Gamaglutamyl transpeptidase (GGT) have been evaluated to explore the functionality of the liver (Burtis and Ashwood, 1999) while the evaluation of haematological parameters helps to determine the degree of toxicity of a test compound as well as to establish the degree of damage to the host tissue. Literatures are full of contradictory information on the toxicity of MSG on human and animals. Considering the discrepancies in the literature and growing safety concern for the use of MSG, there is need for further study on this important food additive. This study was therefore aimed at evaluating the effects of MSG supplemented diets on some liver function and haematological parameters in rats.

Materials and Method

Test Material

Ajinomoto a brand of monosodium glutamate manufactured by Ajinomoto co., inc. Tokyo, Japan was obtained from Bosso market Minna, Niger State, Nigeria. It was stored and kept away from direct sunlight.

Experimental Animals

Healthy Wistar albino rats weighing between 120 – 160g were obtained from animal breeding unit of the Department of Biochemistry University of Ibadan, Oyo State, Nigeria. The animals were kept in clean cages and allowed access to clean water and commercial rat pellet *ad libitum*. The animals were acclimatized for 2 weeks before the commencement of the experiment.

Feed Preparation

Experimental diets, 0.5%, 1.0% and 5.0% of MSG diet were prepared by adding 5g, 10g and 50g of MSG to 995g, 990g and 950g of standard feed respectively. The feed were pelletized according to Larry, (1990).

Experimental Design

A total of twenty (20) rats were used for the experiment and were randomly divided into four groups of five (5) rats each. Rats in group 1(control) were fed with normal commercial feed pellet while those in groups 2, 3 and 4 were placed on 0.5, 1.0 and 1.5% MSG supplemented commercial diets respectively for four weeks.

Blood and Serum collection

After feeding the rats for 28-days, they were anaesthetized with diethyl ether and blood was collected by cervical decapitation into clean lithium heparinised bottles. Blood samples were centrifuged for 5 min and the plasma obtained was subsequently used for biochemical assays while whole blood was used for haematological determinations.

Haematological assay

Haematological assays were carried out on the whole blood samples using haemocytometer (XFA 6000 Intelligent Auto Hematology Analyzer; Germany). The parameters evaluated included (PCV), white blood cell count (WBC) and red blood cell (RBC) count. Packed cell volume was determined using the method of Zuckerman, (2007).

Biochemical Analysis

Randox enzymatic kit was used for the determination of serum aminotransferase enzymes activities (AST and ALT) according to Reitman and Frankel (1957), alkaline phosphatase (ALP) activity (Young *et al.*, 1975), total lipids (Knight *et al.*, 1972), triglycerides (TG) level (Fossati and Prencipe, 1982), total cholesterol concentration (Allain *et al.*, 1974), high density lipoprotein (HDL) level (Lopez- Virella *et al.*, 1977), low density lipoprotein (LDH) level (Wieland and Seidel, 1983).

Results

Table 1 shows the effect of monosodium glutamate on the packed cell volume (PCV), red blood cells (RBC) and white blood cells (WBC) of rats placed on different concentrations of MSG in the supplemented diets. The PCV was highest ($48.50 \pm 1.55\%$) in the group fed 0.5% MSG supplemented diet and lowest ($45.20 \pm 2.72\%$) in the group supplemented with 5% MSG. There was no significant ($p > 0.05$) change in the PCV of the treated groups when compared with the control group. The RBC count in the group fed 1% MSG supplemented diet had the highest ($9.60 \pm 1.01 \times 10^{12}/L$) value while those placed on

5% supplementation had the least value ($9.30 \pm 3.42 \times 10^{12}/L$). Rats fed with 1.0% and 0.5% MSG supplemented diets had the highest ($9.60 \pm 1.10 \times 10^{12}/L$) and least ($9.20 \pm 0.34 \times 10^{12}/L$) values of white blood cell count respectively. This is also the trend observed for RBC count. No significant ($p > 0.05$) difference was observed in the WBC and RBC counts of rats fed with the commercial feed and those in the treatment groups.

The effects of monosodium glutamate on some biochemical indices are shown in Table 2. The concentration of total protein ranged from 8.47 ± 0.88 mg/dl in the group placed on 1% MSG supplemented diet to 8.80 ± 3.55 mg/dl in the group fed 0.5% MSG supplemented diet. The serum albumin level decreased as the percentage of MSG supplementation increased but there was no significant ($p > 0.05$) difference between the control and treatment groups. The same was observed with the concentration of direct bilirubin (0.5% MSG supplemented diet having the least concentration (1.10 ± 0.83 mg/dl) while rats fed with 5% MSG supplemented diet had the highest concentration of 1.40 ± 1.07 mg/dl). There was no significant ($p > 0.05$) difference in the total bilirubin concentration between the control group and other treatment groups.

Figures 1 to 3 show the effect of MSG on some serum enzyme (ALP, AST and ALT) activities. The levels of ALP and ALT activities increased as the percentage of MSG supplementation increased ranging from 35.00 ± 0.17 - 59.00 ± 1.23 IU/L and 6.4 ± 0.32 - 11.9 ± 0.46 IU/L respectively. There was no significant ($p > 0.05$) difference in the ALP and ALT activities in rats fed 0.5% MSG supplemented diets when compared with the control group while the other treatment groups (1 and 5% MSG supplemented diets) groups had their ALP and ALT activities significantly ($p < 0.05$) higher than the control group. The AST activity in all the treatment groups were significantly ($p < 0.05$) higher than the control group (Figure 3).

Table 1: Effect of Monosodium Glutamate on some Haematological Indices in Rats

Groups	PCV (%)	RBC ($\times 10^{12}/L$)	WBC ($\times 10^{12}/L$)
A	44.00 ± 0.84	9.40 ± 3.51	9.43 ± 1.40
B	48.50 ± 1.55	9.40 ± 2.21	9.20 ± 0.34
C	47.00 ± 3.17	9.60 ± 1.01	9.60 ± 1.10
D	45.20 ± 2.72	9.30 ± 3.42	9.35 ± 3.01

Note: Values are mean $\pm$ standard error of mean (SEM) of five determinations. Values along a column are not significantly ($p > 0.05$) different. **A:** Fed commercial feed (Control Group), **B:** Fed 0.5% MSG supplemented diet, **C:** Fed 1% MSG supplemented diet and **D:** Fed 5% MSG supplemented diet

Table 2: Effect of Monosodium Glutamate on some Biochemical Parameters (mg/dl) in Rats

Groups	Total protein	Albumin	Total bilirubin	Direct bilirubin
A	8.58±1.02 ^{ab}	4.47±0.72 ^{ab}	1.31±0.42 ^a	1.16±0.42 ^a
B	8.80±3.55 ^{ab}	4.53±0.32 ^{ab}	1.23±0.57 ^a	1.10±0.83 ^a
C	8.47±0.88 ^a	4.30±2.52 ^a	1.36±0.45 ^a	1.30±1.15 ^a
D	8.49±2.14 ^a	4.16±1.10 ^a	1.00±0.47 ^a	1.40±1.07 ^a

Note: Values are mean±standard error of mean (SEM) of five determinations. Values along a column are not significantly ($p>0.05$) different. **A:** Fed commercial feed (Control Group), **B:** Fed 0.5% MSG supplemented diet, **C:** Fed 1% MSG supplemented diet and **D:** Fed 5% MSG supplemented diet




Figure 1: Effect of Monosodium Glutamate on Alkaline Phosphatase (ALP) in Rats. Values are Mean±SEM (n=5). Bars with * are significantly ($p<0.05$) different from Control group. **A:** Fed commercial feed (Control Group), **B:** Fed 0.5% MSG supplemented diet, **C:** Fed 1% MSG supplemented diet and **D:** Fed 5% MSG supplemented diet.




Figure 2: Effect of Monosodium Glutamate on Alanine Transaminase (ALT) in Rats. Values are Mean±SEM (n=5). Bars with * are significantly (p<0.05) different from Control group. A: Fed commercial feed (Control Group), B: Fed 0.5% MSG supplemented diet, C: Fed 1% MSG supplemented diet and D: Fed 5% MSG supplemented diet.




Figure 3: Effect of Monosodium Glutamate on Aspartate Amino Transferase (AST) in rats. Values are Mean±SEM (n=5). Bars with * are significantly (p<0.05) different from Control group. A: Fed commercial feed (Control Group), B: Fed 0.5% MSG supplemented diet, C: Fed 1% MSG supplemented diet and D: Fed 5% MSG supplemented diet.

Discussion

The monosodium glutamate at the doses administered (0.5, 1.0 and 5.0% per kg of diet) did not cause any significant change in PCV, WBC and RBC when compared with the control group. This is in contrast with the findings of Ashaolu *et al.*, (2011) and Merayebu *et al.*, (2012) who reported that MSG is toxic to the RBC and have deleterious changes in haematological parameters. The variation with other literatures as cited above may be due to some factors such as the route of administration, amount administered or age of experimental subjects.

The insignificant difference in the total protein and albumin values between the control group and MSG fed groups suggests that there was no much depression in hepatic synthesis and or degradation of protein. Total billirubin and direct bilirubin are produced through hepatocytes degradation of erythrocytes (Garba *et al.*, 2013) and are therefore used to access the extent of hepatocellular damage (Paliwal *et al.*, 2009). Thus the insignificant difference in the total and conjugated bilirubin in the MSG fed groups when compared with the control group is an indication that MSG at the dose administered produced no significant liver damage.

The mean serum activity of alkaline phosphatase was significantly higher in all the groups of rats that received MSG when compared with the control group and the activities increased with increase in MSG concentration. Alkaline Phosphatase (ALP) is a biomarker enzyme for assessing the integrity of plasma membrane (Akanji *et al.*, 1993). Increase in the activities of Alkaline phosphatase is an indication that there could be damage due to cytotoxic effect of MSG (Ortiz *et al.*, 2006) thereby resulting to leakage of this enzyme from the liver into the serum. Such increase in alkaline phosphatase activities can constitute threat to the life of cells that are dependent on a variety of phosphate esters for their vital processes since there may be indiscriminate hydrolysis of phosphate esters in the tissue (Yakubu *et al.*, 2006). Increased activity of ALP which occurs due to *de novo* synthesis by liver cells is a reliable marker of hepatobiliar dysfunction due to damage (Muriel and Escobar, 2003). The increased ALP activity may also be due to the increased synthesis in the presence of increasing biliary pressure (Moss and Butterworth, 1974). Again, Rocek *et al.*, (2001) demonstrated that MSG administration could alter the intestinal function thereby releasing intestinal ALP.

The serum levels of ALT and AST were also significantly higher at all doses of MSG when compared with the control group (Figures 2 and 3). Alanine aminotransferase (ALT) enzymes are sensitive biomarkers of hepatic status (Al-mammary *et al.*, 2002). Although ALT and AST are synthesised in the liver, they are also present in serum and in various tissues. In particular, ALT serum levels become elevated during liver diseases, and therefore, it is considered a more specific marker for liver injury than AST (Kunutsor *et al.*, 2013). The elevated level of ALT in the rats fed MSG is an indication of abnormal liver function that might have been induced by the MSG. Aspartate

aminotransferase (AST) is primarily found in the liver mitochondrial and cytoplasm, it is also found in heart, muscle, kidney and brain. Its serum level increases in hepatic necrosis, myocardial infarction and muscle injury (Srinivasan *et al.*, 1997). In the present work, the significant rise observed in the activities of serum AST may be attributed to some damage to the liver or other organs where the AST is located. The increase in the activities of AST and ALT in this work is in agreement with the report of other investigators who have reported increase in the activities of AST and ALT subsequent to MSG administration (Onyema *et al.*, 2006; Thomas *et al.*, 2009). René *et al.*, (2014) observed a gradual increase in AST level while a non significant change in ALT was observed. Contrary to the present findings, Egbuono *et al.*, (2009) reported that low oral dose administration of monosodium glutamate in male albino rats may be nephroprotective. Monosodium glutamate dissociates readily and releases ammonium ion (NH_4^+) which could be toxic unless de-toxified through the urea cycle. Increased glutamate resulting from excess administration of MSG could lead to elevated NH_4^+ concentration as a result of oxidative deamination (Stryer, 1988). Elevation of ammonium ion could result to hepatic lesions, consequently releasing ALT and other enzymes present in the liver into the serum.

Conclusion

This work has shown that consumption of MSG has no effect on haematological parameters, however at very high concentrations varying degrees of injury to the liver may result.

Acknowledgement

This research was assisted by the laboratory unit, Biochemistry department, Federal University of Technology Minna, Niger State Nigeria. We also thank our colleagues who have made useful comments on this manuscript.

Competing Interest

The authors declare that they have no competing interests.

Authors Contribution

This work was carried out in collaboration between all authors. Author AHO designed the study, wrote the protocol and supervised the work. Author PS carried out all laboratories assays and performed the statistical analysis. Author PS wrote the first draft of the manuscript. Authors OIF and AHO edited the manuscript. Authors AHY and OIF worked on the literature review and references in the manuscript.

References

Akanji MA, Olagoke OA Oloyede OB. Effect of repeated consumption of metabisulphite on the integrity of the rat kidney cellular system. *Toxicology*, 1993; 81, 173 - 179.

212 Alao OA, Ashaolu JO, Ghazal OK, Ukwenya VO. Histological and biochemical effects of monosodium
 213 glutamate on the frontal lobe of adult Wistar rats. *Int. J. Biomed. Health Sci.* 2010; 6 (4): 197-
 214 203.

215 Allain CC, Poon LS, Chan CS, Richmond W, Fu PC. Enzymatic determination of total serum cholesterol.
 216 *Clin Chem.* 1974; 20: 470-475.

217 Al-Mamary M, Al-Habori M, Al-Aghbari AM, Baker MM. Investigation into the toxicological effects of
 218 *Catha edulis* leaves: a short-term study in animals. *Phytother Res.* 2002; 16: 127-132.

219 Ashaolu JO, Ukwenya VO, Okonoboh AB, Ghazal OK, Jimoh AAG. Effect of monosodium glutamate on
 220 hematological parameters in Wistar rats. *Inter. J. Med. Med. Sci.* 2011; 3(6), 219-222.

221 Ball P, Woodeard D, Bread T, Shoobridge A, Ferrier M. Calcium disglutamate improves taste
 222 characteristic of lower-salt. *Europ. J. Clin. Nutr.* 2002; .56 (6): 519-523.

223 **Burtis AC, Ashwood T. Textbook of Clinical Chemistry.3rd Ed. 1999, AACCW.B. Sounders co. London**

224 Egbuonu ACC, Obidoa O, Ezeokonkwo CAL, Ezeanyika LUS, Ejikeme PM. Low dose oral
 225 administration of monosodium glutamate in male albino rats may be nephroprotective. *Bio-Res.*
 226 2009; (1): 470-473.

227 Fossati P, Prencipe L.. Serum triglycerides determined colorimetrically with an enzyme that produced
 228 hydrogen peroxide. *Clin Chem.* 1982; 28(10): 2077-2080.

229 Garba U, Aminu N, Musa H, Musa AB, Timothy S. Biochemical and histogathologic changes in liver of
 230 Albino rat expose to 1% dichlorvos at sub acute period. *J. Pharmbiomed. sci.* 2013; 3(2): 1-6.

231 Gobatto CA, Mello MA, Souza CT, Ribeiro IA. The monosodium glutamate (MSG) obese rat as a model
 232 for the study of exercise in obesity. *Res Commun. Mol. Pathol. Pharmacol.* 2002; .3 (4): 89-307.

233 Ikeda K. On the taste of the salt of glutamic acid. *Journal of Tokyo Chemical Society.*1909; .30: 820-836.

234 Knight TA, Anderson S, James MR. Chemical basis of the sulphophospho – vanilline reaction for
 235 estimating total serum lipids. *Clin Chem.* 1972; 18(3): 199-202.

236 Kunutsor SK, Apekey TA, Walley J. Liver aminotransferases and risk of incident type 2 diabetes: A
 237 systematic review and meta-analysis. *Am. J. Epidemiol.* 2013; 178: 159–171.

238 Larry, HP Pelleting’s history and development. *Feed manag.* 1990; 41(10): 51-72.

239 Legetic B, Campbell N. Reducing salt intake in the Americans: pan American health organization action.
 240 *J. Health Comm.* 2012; 2, 37-48.

241 Loliger J. Function and important of glutamate of savoury foods. *J. Nutr.* 2000; 130 (4), 915-920.

242 Lopez – Virella MF, Stone P, Ellis S, Colwell JA. Cholesterol determination in high density lipoproteins
 243 separated by three different methods. *Clin Chem.* 1977; 23(5): 882-884.

244 Manal ST, Nawal A. Adverse effect of MSG on liver and kidney function and potential protective effect
245 of vitamin C and E. Food Nutr. Sci. 2012; 3: 651-659.

246 Meraiyebu A, Akintayo CO, Uzoechi AC, Okere S. The effects of orally administered monosodium
247 glutamate (MSG) on blood thrombocyte, blood coagulation and bleeding in rats. J. Pharm. Bio.
248 Sci. 2012; 4(1): 4-8.

249 Moss D.W, Butterworth, PJ. Enzymology and Medicine. Pitman Medical, London, 1974; 139.

250 Mozes S, Sefcikova Z. Obesity and changes of Alkaline Phosphatase Activity in the Small Intestine of 40
251 and 80-day old rats subjected to early postnatal overfeeding of monosodium glutamate. Physiol.
252 Res. 2004; 53, 177-186.

253 Muriel P, Escobar Y. Kupffer cells are responsible for liver cirrhosis induced by carbon tetrachloride. J.
254 Appl. Toxicol. 2003; 23(2): 103-108.

255 Onyema OO, Farombi EO, Emerole GO, Ukoha AI, Onyeze GO. Effect of vitamin E on monosodium
256 glutamate induced hepatotoxicity and oxidative stress in rats. Indian J. Biochem. Biophy. 2006;
257 43: 20-24.

258 Ortiz GG, Bitzer- Quintero OK., Zarate CB, Rodriguez- Reynoso S, Larios- Arceo F. Monosodium
259 glutamate- induced damage in liver and kidney: a morphological and biochemical approach.
260 Biomed. Pharmacol. 2006; 60, 86-91.

261 Oyetunji TK. Assessment of the effect of monosodium glutamate on some biochemical and hematological
262 parameters in adult Wistar rats. Am. J. Biosci. 2013; 1(1): 11-15.

263 Paliwal A, Gurjar RK, Sharma HM. Analysis of liver enzymes in albino rat under stress of 1-
264 cyhalothrin and nuvan toxicity. 2009; (cited on 4th December 2011). Available from:
265 <http://www.biolmedonline.com/article/view/4317>

266 Reitman S, Frankel S. A colorimetric method for the determination of serum glutamic oxaloacetic and
267 glutamic pyruvic transaminases. Am. J. Clin. Pathol. 1957; 2: 56-60.

268 René JH, Francisco JA, María Del CE, Julio CA, Héctor M, Mina KF, Norma EL. Biochemical
269 Alterations during the Obese-Aging Process in Female and Male Monosodium Glutamate
270 (MSG)-Treated Mice. Int. J. Mol. Sci. 2014; 15: 11473-11494

271 Rocek L, Lenharalt L, Mozes S. Effect of feeding and refeeding on duodenal alkaline phosphatase activity
272 in monosodium glutamate obese rats. Physiol. Res. 2001; 50: 365-372.

273 Singh P, Mann K.A., Mangat K., Kaur G. Prolonged glutamate excitotoxicity: effects on mitochondrial
274 antioxidants and antioxidant enzymes. Mol. cell biochem. 2003; 243: 139-145.

275 Srinivasan K., Radhakrishnamurthy R.. Effects of β and γ isomers of hexachlorocyclohexane on some
276 liver and kidney enzymes in albino rats. Current Sci. 1977; 46 (17): 598-600.

277 Stryer L. Amino acid degradation and the urea cycle. In *Biochemistry* edition (3rd). W.H Freeman and
278 company, New York. 1988; Chapter 21: 495-516

- 279 Thomas M, Sujatha KS, George S. Protective effect of *Piper longum* Linn. on monosodium glutamate
280 induced oxidative stress in rats. Indian J. Exper. Biol. 2009; 47(3): 186-192.
- 281 Wieland H, Seidel D. A simple specific method for precipitation of low density lipoproteins. J. lipid
282 Res.1983; 24(7): 904-909.
- 283 Yakubu MT, Adesokan AA, Akanji MA. Biochemical change in the liver kidney and serum of rat
284 following chronic administration of cimetidine. Afri J. Biomed Res. 2006; 9: 213-218.
- 285 Young DS, Pestner LC, Gibberman V. Effects of drugs on clinical laboratory tests. Clin. Chem. 1975; 21,
286 246.
- 287 Zuckerman K. *Approach to the anaemia* In: Goldman L, Ausiello D. (Editors). Cecil Medicine, 23rd
288 edition, Pa: Saunders Elsevier, Philadelphia, 2007; Chapter 162.