

Original Research Article

ANTAGONISTIC COMPATIBILITY OF *Streptomyces griseorubens*, *Gliocladium virens*, and *Trichoderma harzianum* AGAINST *Fusarium oxysporum* CAUSE OF TOMATO WILT DISEASES.

ABSTRACT

This research was intended to discover the compatibility of biological control agents (BCAs) *Streptomyces griseorubens*, *Gliocladium virens*, and *Trichoderma harzianum* against *Fusarium oxysporum* f.sp. *capsici* *in vitro* and *in vivo*. Study was done in rainy season 2012-2013 at East Java-Indonesia. Study was a true experiment designed, using a completely randomized design (CRD), consist of three stages research. These stages were the compatibility of three biological agents in Potato Dextrose Agar (PDA) medium, antagonistic test of biological agents combinations of *S. griseorubens*, *G. virens*, and *T. harzianum* capable inhibited microbial pathogens *F. oxysporum* f.sp. *lycopersici* in the laboratory and at the screen house. Mixed of BCAs *S. griseorubens*, *G. virens*, and *T. harzianum* were compatible and effectively against *F. oxysporum* in Petri dishes and at screen house. Clear zona average of antibiosis of BCAs filtrat in Potato Glucose extract shown that the antibiosis from mixed of *S. griseorubens* and *T. harzianum* was higher than the antibiosis of other BCAs treatment. Moreover plant infested with mixed of BCAs significantly protected plant tomato from *F. oxysporum* compared to the untreated control plants. Plant protection by BCAs mixed of *T. harzianum* with *S. griseorubens* was more pronounced than plant protection by mixed of *S. griseorubens* with *G. virens* and single BCAs.

Keywords: Wilt diseases, Antagonistic, mixed of BCAs

1. Introduction

Fusarium oxysporum f.sp. *lycopersici* (FOL) is a highly destructive pathogen of both greenhouse and field grown tomatoes in warm vegetable production areas. The disease caused by this fungus is characterized by wilted plants, yellowed leaves and minimal or absent crop yield. Based on our surveys, at East Java (Malang, Pare, Kediri) there was a 10 to 12 % yield loss. (10,14).

F. oxysporum is soil-borne plant pathogens and the most difficult to control. Chemical control effect negative to environment, there is renewed interest in biological control based on application of populations of antagonistic micro-organisms. (2,15). Soil microorganism challenge to be BCAs, the advance of technology these days has come to the application on field, by hoping to be able to make efficient of natural resources, conservation and the everlasting environment, also to produce cheaper and healthier agriculture products. Some saprophyte soil microorganism have been used as commercial biological agents, It can be single antagonistic or multi antagonistic (7). The research and the usage of *S. griseorubens* as a biological agent is rarely conducted in agriculture field especially in Indonesia, meanwhile the bacteria, fungus and virus had been through many researches. *T. harzianum* fungus is a saprophyte soil fungus able to become hyperparasite to several species of fungus pathogen. The growth of *T. harzianum* is very rapid and unpathogen for high level plants. The hypha threads of pathogenic fungus will be cut to pieces

because it wound by *Trichoderma* hypha (as antagonist fungus). *Trichoderma* eventually release antibiotics to pathogens fungus which is gliotoxin (9,13,23,25). *G. virens* control plant pathogen by several mechanisms such as parasitism, antibiosis, competition and cell destruction. *Gliocladium* will grow around the pathogen and release enzyme that can destroy cuticle of pathogen. *G. virens* also producing gliotoxin antibiotics (11). *S. griseorubens*, *G. virens* and *T. harzianum* as the single antagonist is able to control fusarium wilt disease on tomato and melon in the greenhouse scale (10,22). This research was intended to discover the biological agents *S. griseorubens*, *G. virens*, and *T. harzianum* capable inhibited microbial pathogens *F. oxysporum* f.sp. *lycopersici* in the laboratory and in the screen house..

2. Method and Material

2.1 Isolation of Biological agents

Isolation of biological agents used soil plating method by Dhingra and Sinclair (21): 1 gram of Pare-Kediri chilli and tomato soil was made suspension by dilution 10^4 . Subsequently 1 mL of soil suspension was spreaded on Glucose Nutrient Agar (GNA) to get *S. griseorubens* and 1 mL of *T. harzianum* and *G. virens* suspension (BPTPH Pandaan) also spreaded on Potato Dextrose Agar (PDA). Biological agents obtained was purified and propagated on PDA in Petri dishes.

2.2 Isolation of *F. oxysporum* f.sp. *lycopersici*

Cut into the base of the stem of a diseased plant lengthwise to reveal the xylem just below the epidermis. Turn off all the leaves and secondary roots, leaving only the main stem and the hypocotyls and main root. Sterilize the surface of stem by soaking in 10% bleach solution for 5 minutes. Dry the stem on paper towels. Using sterile technique, cut thin (2-4 mm thick) wedges out of one side of the stem near the root/stem junction making sure to include xylem tissue with each wedge. Place 5-6 wedges on PDA plates. Incubate the plates under fluorescent lights. Once the fungus has grown sufficiently from the pieces, transfer isolates onto fresh PDA plates. Incubate the plates for 10-14 days. Colonies of *F. oxysporum* are pigmented with a reddish purple color and surmounted by a pinkish white aerial mycelium (18).

2.3 Compatibility test

Compatibility test was done in Microbiology laboratory of Agriculture Faculty by watching the type of biological agents growth in descriptive and diameter colony. Every treatment was repeated five times. Diameter colony average were analysis by t-test (12)

Preparing 0.5 cm diameter colony of 10th day biological agents, *S. griseorubens*, *T. harzianum*, *G. virens*. Compatibility tests on the PDA media plating in 20 diameter cm Petri dishes were three types of biological agents colony *S. griseorubens*., *G. virens*., and *T. harzianum* (SGT) were placed on PDA medium, each with the same distance of 5 cm, and were incubated for 14 days. Along with the treatment of compatibility, also it was prepared control treatment as a comparison.

2.3 Antagonism compatibility *in vitro* test

Antagonism compatibility *in vitro* was done by completely randomized design (CRD) with seven treatment and every treatment was repeated three times. Data of percentage inhibition was analyzed by Duncan test

Biological agents suspension treatment 6 mL was mixed up in 44 mL of sterile water, 0.33 cc of this each suspension biological agents treatment was taken into holes

(wells). Then put the 0.5 cm colony of *F. oxysporum* 7 days age, in the presence of biological agents suspension that has been inoculated in the hole at a distance of 5 cm. Each treatment was randomly stored at room temperature for 8 days. The inhibition was calculated by the formula (8):

$$DI = \frac{Dc - Dt}{Dt} \times 100$$

DI is the percentage inhibition;

Dc is the diameter colony of the control treatment;

Dt is the diameter colony of BCAs treatment.

2.4 Antibiosis in *Potato Glucose Extract* medium

Antibiosis test consist of 7 treatment placed in randomly, every treatment was done in three repeated.

Six mili liter of BCAs (every BCA 38-42 spore/cc) treatment entered into 44 mL Potato Glucose Extract solid medium in Enlenmeyer tubes. BCAs suspension was shaken. Inhibition ability of antibiosis filtrat of BCAs againts *F. oxysporum* was done by Steinkelner method. 0.55 cc antibiosis was given into 0.5 cm Whatman paper disc, then air drying them. These paper disc was inoculated on PDA medium in Petri dish contains *F. oxysporum* suspension. Inhibition ability of antibiosis was done by counting of clear zone diameter (27).

2.5 Antagonism *in vivo* test

This research was done by Completely Randomized Design (CRD) with seven treatment and every treatment was repeated three times. Data of percentage inhibition was analyzed by Duncan test.

Seedling was inoculated by soaking a solution of inoculum combination of biological agents that have been prepared before. Furthermore, the seed was planted in the soil that had been inoculated *F. oxysporum* suspension (10^9 spore/mL) for 14 days. It was prepared by filled 3 liters of sterile soil in polybags and inoculated with spore inoculum mass suspension of *F. oxysporum* has been sprayed with a hand sprayer at ground level (21). Stored for 14 days in the screen house and watering with sterile water every day. Data were collected for : (1) Period of incubation, performed daily until symptoms of disease were yellowing of the leaves from the bottom, where the control healthy plants not showing symptoms. (2) Severity of the disease, was conducted with most of the leaves are yellow, wilt and dry every 7 days until harvest (3).

$$KP = \frac{\text{count of yellowing leaf}}{\text{count of all leaf of plant}} \times 100\%$$

KP is the percentage of disease severity per plant(3)

3. RESULTS AND DISCUSSION

3.1 Compatibility test

Performance of three biological agents shown no antagonistic of them. *S. griseorubens* (S) *G. virens* (G) and *T. harzianum* (T) grew more thick than single biological agent (Fig. 1)





Figure 1. *T. harzianum* colony(T), *S. griseorubens* colony (S), *G. virens* colony (G) on PDA medium, 14th days after inoculations BCAs.

Growth average of diameter colony *S. griseorubens* (S) *G.virens* (G) and *T. harzianum* (T) in compatibility test (SGT) on 2nd days after inoculation shown no significantly different when compared with controls (SSS, GGG, TTT). However, on 4th and 6th days after inoculations, diameter growth of biological agents such colony in compatibility testing (SGT) was larger and significantly different than the control. On 8th days, the growth of biological agents on a compatibility test was not different from the control of biological agents (Fig.2)






Figure 1. Graphic of Biological agents colony diameter on compatibility test. SGT/GTS/TGS are the same treatment, consist of *S. griseorubens*, *G.viren*, *T. harzianum*, Control consist of three colony of GGG (*G.viren*), TTT (*T. harzianum*), SSS (*S. griseorubens*.)

S. griseorubens, *G. virens* and *T. harzianum* as biological agents could grow on PDA media and formed an association that does not harm each other or not produced secondary metabolites that could inhibit the growth of biological agents each other. The three biological agents are saprophyte soil microbes, and produce antibiosis to mycoparasite only affect microbial pathogens. This opinion is supported by the results of some other researchs, plant pathogenic fungi cell walls composed of chitinase which is a key enzyme and responsible for the lysis of the cell wall. *S. griseus* degrade fungal cell walls by lytic enzymes (8). *Trichoderma* sp. produced lytic enzyme that degraded chitin, mycoparasite process and can improve the cell wall itself on the division process (16). *Gliocladium* sp. and *Trichoderma* sp. produce chitinase enzymes that can cause parasitic on plant pathogens, whereas less effective antibiosis produced degrades *F. oxysporum* (11). The existence of microbes of different biological agents also induce the microbes to grow faster. As noted by some researchers that *Trichoderma* sp. and *Gliocladium* sp. produce fungal pathogen that work in synergy with the intracellular enzyme produced by *G.virens*. Both of these biological agents were synergy in controlling pathogenic tomatoes until 57% (23; 24).

3.2 Antagonism compatibility *in vitro*

Giving a single biological agents *S. griseorubens* (S), *G.virens* (G), *T. harzianum* (T), a mixed of two biological agents (SG, ST, GT) and a mixed of three biological agents (SGT) to tomato seed, significantly inhibited the development of the colony diameter of *F. oxysporum*. Giving only *G.virens* to tomato seed shown smaller and significantly different average inhibition than the other single of biological agents *S. griseorubens* and *T. harzianum* and the combination treatment (Table 1).

243 Tabel 1. Inhibitors average of biological agents *S. griseorubens* (S), *G. virens* (G) and
 244 *T. harzianum* (T) against *F. oxysporum*

Treatment of BCAs	Inhibitor average \pm SE(%)
<i>S. griseorubens</i> , <i>T. harzianum</i> (ST)	63.97 \pm 7.27 ^a
<i>G. virens</i> , <i>T. harzianum</i> (GT)	59.29 \pm 3.87 ^{ab}
<i>S. griseorubens</i> , <i>G. virens</i> , <i>T. harzianum</i> (SGT)	59.07 \pm 1.82 ^{ab}
<i>S. griseorubens</i> , <i>G. virens</i> (SG)	58.41 \pm 1.93 ^{ab}
<i>T. harzianum</i> (T)	49.05 \pm 3.41 ^{ab}
<i>S. griseorubens</i> (S)	45.11 \pm 5.37 ^b
<i>G. virens</i> (G)	40.24 \pm 7.27 ^c

245 Explanation : The same letters in addition to numbers showed no significant
 246 difference in the Duncan level test ($p < 0.05$).
 247

248 Single biological agent *G. virens* (G), have inhibition ability of microbial pathogens
 249 *F. oxysporum* (F) lower than the inhibition ability of single biological agents *T. harzianum*
 250 and *S. griseorubens* (T, S), mixed of biological agents (TS, GS, TG and TGS). *G. virens*
 251 serves only as competitors and parasite, it did not produce antibiosis on PDA media (*in*
 252 *vitro*) (11,15,21). Antibiosis of biological agents *G. virens*, *T. harzianum* and *S. griseorubens*
 253 on Potato Glucose Extract (PGE) also proved that *G. virens* not produced antibiosis on
 254 observations 2nd, 4th, 6th day after inoculation of biological agents. Single BCA *T. harzianum*
 255 and *S. griseorubens* and combination of BCAs on observation produced antibiosis (Table 2)
 256

257 Table 2. Avarage of Inhibitor zone of biological agents crude extract to
 258 *F. oxysporum*

No	Biological agents giving	Clear zone (halo) ... mm		
		2 dai	4 dai	6 dai
1	<i>T. harzianum</i>	0,2	0,2	0,3
2	<i>G. virens</i>	0	0	0
3	<i>S. griseorubens</i>	1,3	0,4	0
4	<i>G. virens</i> , <i>T. harzianum</i> .	0	0	0
5	<i>T. harzianum</i> , <i>S. griseorubens</i>	1,2	0,7	0,2
6	<i>G. virens</i> , <i>S. griseorubens</i>	0,3	0,3	0
7	<i>G. virens</i> , <i>S. griseorubens</i> and <i>T. harzianum</i>	0	0,4	0,2

259 Notes : Dai is days after inoculations
 260

This suggests that a **mixed** of all three biological agents produced antibiosis derived from *T. harzianum* and *S. griseorubens f.sp. capsicum*. Single biological agents *T. harzianum* (T) and a **mixed** of biological agents (TG, TS, GS and SGT) can result inhibition ability higher than inhibition ability of single biological agents *S. griseorubens* (S) and *G.virens* (G). Several studies have shown that the fungus *Trichoderma* sp. is a saprophyte **fungi** that lives in the soil and becomes hyper parasite on some pathogenic fungi. *T. harzianum* also inhibit the growth of *F. oxysporum* colonies growing very rapidly, and producing antifungal namely Glicotoxin. *Gliocladium* sp., *Streptomyces* sp. *Trichoderma* sp. clasified as soil saprophyte fungi and **used as biological agents have multi antagonistic mechanism and compatible againts *F. oxysporum*** (27).

3.4 Antagonistic compatibility **in vivo**

Tomato plants that were not given biological agents showed **Fusarium** wilt symptoms on 39-40th days after planting tomato seedlings. Meanwhile, the plants treated with biological agents, appeared symptoms at 46-50th days after planting. Longer incubation occured because of biological agents mixture could inhibit the development of *F. oxysporum*. Giving of biological agents with the pathogen in the soil for one week, the incubation period occurred after 45 days (20). **S**ingle biological agents *S. griseorubens* (S) and a **mixed** of biological agents, *S. griseorubens* with *G.virens* (SG), *S. griseorubens*, with *T. harzianum* (ST), and a mixed of *S. griseorubens*, *G.virens*, *T. harzianum* (SGT) can inhibit disease severity (Table 3).

Table 3 Avarage of wilt deseases severity of tomato plant at 41th, 48th, 55th and 62th days after planting (dap).

No	Giving of biological agents	Avarage of deseases severity (%)			
		41 dap	48 dap	56 dap	62 dap
1	Control (withaot bcas) (K)	3.80±2.19 ^a	7.60±0 ^a	17.86±1.57 ^a	44.64±3.30 ^a
2	<i>S. griseorubens</i> , <i>T. harzianum</i> . (ST)	0.00±0 ^a	5.70± 1.9 ^a	7.14± 5.91 ^a	19.65±3.41 ^b
3	<i>S. griseorubens</i> (S)	0.00± 0 ^a	1.90±1.9 ^a	15.42±2.90. ^a	16.07±7.46 ^b
4	<i>S. griseorubens</i> , <i>G. virens</i> (SG)	1.90± 1.9 ^a	3.80±2.19 ^a	7.15± 4.13 ^a	16.07±7.36 ^b
5.	<i>S. griseorubens</i> , <i>G. virens</i> , <i>T. harzianum</i> (SGT)	1.52±1.9 ^a	5.77±5.77 ^a	5.77±3.77 ^a	14.29±5.63 ^b

Explanation: The same alphabet beside the number in the same coloum shown insignificant Duncan test (p < **0.05**)

Each of these biological agents inhibit the development of pathogens with multiple mechanisms and can develop optimally in soils containing organic matter, so that they can be complement each other. Based on several studies, the *Trichoderma* sp., *Streptomyces* sp., and *Gliocladium* sp., were antagonistic to fungal and bacterial diseases of plant roots (**3**, **20**, **27**). Three biological agents can flourish together on compost, manure and garden soil . In synthetic medium, *S. griseorubens* did not develop **optimal**, **but** developed optimal in field conditions. The study also found that for 4 weeks, the average population of biological agents *S. griseorubens* more higher than the avarage populations of biological agents *T. harzianum* (25).

On 55th and 62nd days after planting, combination of biological agents *T. harzianum* and *S. griseorubens* demonstrated lower power resistor ability of diseases severity than power resistor ability of other biological agents combinations. Several studies in the screenhouse and in the field proved that inhibit ability of *T. harzianum* to pathogens was lower than inhibit ability of *G.virens* in soil. Otherwise, *Gliocladium virens* grew very fast and produced antibiotics gliovirin. Antibiotics worked in synergy with intracellular enzymes to inhibit the development of fungal plant pathogens (3,4). Providing a mixed of biological agents *T. harzianum* (T), *G.virens* (G), *S. griseorubens* f.sp. *capsicum* (S) can also enhance plant growth and provide organic material for plants as planting soil decomposition. Biological agents can also grow pheriphere new roots more and replaces the root of suffering discoloration and can repair the affected plant roots by *F. oxysporum* (17,26).

4. Conclusion

S. griseorubens, *G. virens* and *T. harzianum* as biological agents were compatible grow on PDA media and formed an association that does not harm each other or not produced secondary metabolites that could inhibit the growth of biological agents each other. A single biological agents *S. griseorubens* (S), *T. harzianum* (T), a mixed of two biological agents (SG, ST, GT) and a mixed of three biological agents (SGT) more inhibited the development of the colony diameter of *F. oxysporum* than a single biological agent *G. virens* *in vivo*. Giving mixed of two biological agents *S. griseorubens* and *G. virens* as well as *S. griseorubens* and *T. harzianum* as well as three biological agents *S. griseorubens*, *G. virens* and *T. harzianum* to inhibit disease severity of tomato fusarium wilt caused by *T. oxysporum* f.sp. *lycopersici*

CONSENT

All authors declare that written informed consent was obtain from the approved of our research parties

REFERENCES

1. Abeysinghe, S. 2007. Biological control of *Fusarium solani* f.spp. *phaseoli* the causal agents of root rot of bean using *Bacillus subtilis* CA 32 and *Trichoderma harzianum* RU01. *Ruhuna Journal of Science*. 2: 62-88.
2. Alabouvette, C., C. Olivain, Q. Migheli, C. Steinberg. 2009. Microbiological control of soil-borne phytopathogenic fungi with special emphasis on wilt-inducing *Fusarium oxysporum*. *On Line Journal* DOI: 10.1111/j.1469-8137.2009.03014.x
3. Anitha, A. and M. Rabeeth. 2010a. Control of Fusarium wilt tomato by bioformulation of *Streptomyces griseus* in green house condition. *African Journal of Basic & Applied Sciences* 1 (1-2): 9 – 14.
4. Anitha, A. and M. Rabeeth. 2010^b. Degradation of fungal cell walls of phytopatogenic fungi by lytic enzyme of *Streptomyces griseus*. *African Journal of Plant Science* . 4 (3): 061-066.
5. Asha, B., C. Nayaka, U. Shankar, Srinivas, Nirjana. 2011. Biological control of *F. oxysporum* f. sp. *lycopersici* causing wilt of tomato by *Pseudomonas fluorescens*. *International Journal of Microbiology Research*. 3(2): 79-84

- 345 6. Bollen, G.J. 1974. Fungal recolonization of heat-treated glasshouse soils. *Agro*
346 *Ecosystems* 1: 139-155
- 347 7. Cook J. R. and K. F. Baker. 1996. The nature and practice of biological control of
348 plant pathogens. APS PRESS. *The American Phytopathological Society*. St.
349 Paul, Minnesota.
- 350 8. Fahri, Y. and M. Dikilita. 2007. Control of fusarium wilt of tomato by combination of
351 *Pseudomonas fluorescent*, non *patogen Fusarium* and *Trichoderma*
352 *harzianum* T-22 in greenhouse conditions. *Plant Pathology Journal* 6(2) :
353 159-163.
- 354 9. Gruber, S. and V. Seiboth. 2012. Self versus non-self: fungal cell wall degradation In
355 *Trichoderma. Microbiology* 158:26-34.
- 356 10. Kaewchai, S., Soyong, K., and Hyde, K.D. 2009. Mycofungicides and fungal
357 biofertilizers. *Fungal Diversity*. 38: 25-50.
- 358 11. Kyeong, S. J., Hong, M. K., and Bong, K. C., 2000. Purification and antifungal activities
359 of an antibiotic produced by *Gliocladium virens* G1 against plant pathogen.
360 *Plant Pathology Journal*, J.17(1) : 53-56.
- 361 12. Kusningrum, R.S. 2008. *Perancangan Percobaan*. Airlangga University Press.
- 362 13. Larkin, R.P. and D.R. Fravel. 1993. Biocontrol of Fusarium Wilt of tomato. Biocontrol of
363 Plant Diseases Laboratory. Bestville
- 364 14. Menzies JG, Koch C, Seywerd F. 1990. Additions to the host range of *Fusarium*
365 *oxysporum* f. sp. *radicis-lycopersici*. *Plant Diseases*. 74: 569-572.
- 366 15. Morid, B., S. Hajmansoor, N. Kakvan, 2012. Screening of resistance genes to fusarium
367 root rot and fusarium wilt diseases in tomato (*Lycopersicon esculentum*)
368 cultivars using RAPD and CAPs markers. *European Journal of*
369 *Experimental Biology*. 2 (4):931-939
- 370 16. Nourozian J., H. R. Etebarian, and G. Khodakaramian, 2006. Biological control of
371 *Fusarium grameniarum* on wheat by antagonistic bacteria. *Songklanakarin*
372 *Journal, Sci Technol* 28 :29-38.
- 373 17a. Olivain C., C. Humbert, J. Nahalkova, J. Fatehi, FL. Haridon, and C. Alobouvette. 2006.
374 Colonization of tomato root by pathogenic and non pathogenic *Fusarium*
375 *oxysporum* strains inoculated together and separately into the soil.
376 *Aplaid and Enviromental Microbiology*. 72 (2): 1523-1531.
- 377 18. Reis A, Costa H, Boiteux LS, Lopes CA. 2005. First Report of *Fusarium oxysporum* f. sp.
378 *lycopersici* Race 3 on Tomato in Brazil. *Fitopatologi. Bras*. 30(4): 426-428.
- 379 19. Siddiquee, Shafiquzzaman, Soon G. T., and Y. U. Kalsum. 2010. Isozyme Analisis and
380 Relationships Among Three Species in Malaysia *Trichoderma* Isolates.
381 *Mycrobial Biotechnol*. 20(9): 1266 – 1275
- 382 20. Singh R., B.K. Singh, R.S. Upadhyay, B. Rai and Y. S. Lee. 2002 Biological control of
383 fusarium wilt disease of pigeonpea. *Plant Pathology Journal* 18(3) : 279-
384 283.
- 385 21. Singleton, J.D. Mihail and C.M. Rush. 1993. Methods for research on soilborne
386 phytopathogenic fungi . APS Press. *The American Phytopathological Society*.
387 St. Paul Minesota.
- 388 22. Staniazsek M, Kozik EU, Marczewski W . 2007. A CAPS marker TAO1902 diagnostic
389 for the I-2 gene conferring resistance to *Fusarium oxysporum* f. sp.
390 *lycopersici* race 2 in tomato. *Plant Breeding*. 126(3): 331-333.
- 391 23. Stipanovic, R.D. and C.R. Howell. 1982. The Structure of *Gliovirin*, A new Antibiotic
392 from *Gliocladium virens*. *The Journal of Antibiotics* cotoxin.
- 393 24. Suharjono, Tri Handayani, Soejono, Susanti Dewi. 2008. Antagonis test of *Trichoderma*
394 sp. dan *Gliocladium* sp. againts *Fusarium oxysporum* cause of wilt
395 diseases of some variety of Purwodadi field banana in Vitro (*In*
396 *Indinesian*). Biologi Study, Mathematic and Scient Faculty, Unibraw
397 Malang,

- 398 25. Suryaminarsih dan Mujoko. 2012. Growth population of multiantagonis *Streptomyces*
399 sp. *Gliocladium* sp and *Trichoderma harzianum* as biological agents of
400 fusarium wilt disease in natural and semi natural package pellet formula
401 (In Indonesian). *Plumula*, 1 (2): 202-210.
- 402 26. Suryaminarsih, Kusningrum, Ni'matuzahroh, Surtiningsih, 2014. Plant Resistance with
403 pheriphere new roots by BCAs *Gliocladium* sp and *T. harzianum* againts *F.*
404 *oxysporum* on sprout of tomato. *Prossiding of Plant Protection national*
405 *Seminar (In Indonesian)*.
- 406 27. Steinkellner S., R. Mammerder, and H. Vierhellig. 2008. Germination of *Fusarium*
407 *oxysporums* in root exudates from tomato plants callenged with diffrent
408 *Fusarium oxysporums* strains. *Plant pathology*. 122: 395-401
- 409 28. Titus, A. and G.N. Pereira. 2008. The role of Actinomycetes in coffee plantation
410 ecology. Ineedcoffee.com