

**The effect of soil use and management on the chemical properties of an Ultisol**

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

---

**ABSTRACT**

The use of vegetation cover to protect the soil and increase the soil organic matter content available to plants is a good practice to minimize loss of soil quality. The effect of soil management practices on the objective of this study was to evaluate the soil quality under different types of management. A completely randomised experimental design was used with three treatments as; area of natural vegetation; Area of perennial crop cultivation and degraded pasture area; with five replications. Chemical soil properties evaluated were: phosphorus, potassium, magnesium and calcium contents by extraction with ion exchanger resin. The content of organic matter was determined by the colorimetric method and the pH, in calcium chloride, as well as the acidity potential of the pH 7.0. The sums of bases, cation exchange capacity and base saturation were calculated. Under the treatment of natural vegetation the soil depths: 0.00-0.10; 0.10-0.20 and 0.20-0.40m. Data were analyzed statistically using the Tukey test at 5% for comparison of means. In the treatment, there was an accumulation of exchangeable bases, organic matter and phosphorus, as well as reducing potential acidity. The results show that soil use in spite of its management interferes with the chemical quality of the soil.

*Keywords:* organic matter, soil acidity, phosphorus.

**1. INTRODUCTION**

Sustainable use of natural resources, especially soil and water, has become a topic that is increasingly relevant, due to the increase of anthropogenic activities. Consequently there is, increases the concern with the quality and sustainable use of these resources (Araújo et al., 2007).

The introduction of agricultural systems to replace the forests causes an imbalance in the ecosystem in which the removal of the original vegetation and the deployment of crops, combined with inadequate management practices, lead to the disruption of the balance between the soil and the environment. In this way, the modification of soil chemical, physical and biological properties, limits the soils their agricultural use (Richart, et al., 2005).

The quality to attributes provides suitable conditions for the growth and development of plants and the maintenance of the diversity of organisms that inhabit the soil (Doran & Parkin, 1994). The current research challenge is the assessment of soil quality of reliable According to Doran & Parkin (1994)., it can be measured by quantifying to attributes, i.e., physical, chemical and biological properties have been used to monitor medium and long term changes, in the quality of the soil. Soil quality must be monitored to detect changes that may be mediated at a certain time and can be done on the farm or on a larger scale, such as watersheds and regions.

36

37 According to Araujo et al.,(2007), management practices and soil conservation and water  
38 must be planned and executed attempting to maintain or even improve their attributes, in  
39 order to increase the soil's ability to sustain a competitive biological productivity. The  
40 conventional management systems (plowing and harrowing), results in intensified processes  
41 of erosion and soil compaction which in the medium and long term, may lead to physical,  
42 chemical and biological degradation of the soil (Leite et al., 2003; Araújo et al., 2004). In  
43 these conditions, the absorption of nutrients available to the root system is hampered and  
44 the amount of oxygen in the rhizosphere can become limiting in several metabolic processes  
45 (Beulter & Centurion, 2004).

46

47 Islam & Weil (2000) proposed the use of data of physical, chemical and biological  
48 characteristics of soil collected in an area of undisturbed natural forest, as a reference for  
49 assembling a general index of soil quality. According to Brookes (1995), the relationship  
50 between the physical, chemical and biological properties of the soil vary in time and space.  
51 And these changes can be beneficial or harmful to production systems. Proper management  
52 promotes gain in soil quality and crop productivity. Consequently, the variation of these  
53 attributes, determined by management and soil use, and its evaluation are important for  
54 better management aiming at sustainability of the system (Carneiro et al., 2009).

55

56 Works performed by Souza & Alves (2003) in dystrophic Oxisol, show that soil management  
57 systems with less or no soil disturbance led to significant accumulation of organic matter,  
58 phosphorus, potassium, magnesium and calcium and decrease of exchangeable aluminum  
59 and potential acidity. With soil disturbance as a result of cropping, the effects was a  
60 reduction of the levels of organic matter, phosphorus, potassium, calcium and magnesium  
61 and increased potential acidity and exchangeable aluminum.

62

63 Studies conducted in Oxisol, in the region north of the state of Pará, found that the sum of  
64 bases and cation exchange capacity (CEC) were the most sensitive variables to changes in  
65 the use and management (primary forest, explored forest, grassland, scrub, jiquira of typical  
66 Brazilian vegetation), and the pH and potential acidity and soil pH were influenced by the  
67 replacement of forest by pasture (Ferreira et al., 2011).

68

69 The aim of this work was to evaluate the chemical quality of the soil of an Ultisol under  
70 different uses and management.

71

## 72 2. MATERIALS AND METHODS

73

74 The experiment was conducted at the experimental area of farm Teaching and Research at  
75 the University Camilo Castelo Branco - UNICASTELO, Fernandópolis city, São Paulo (SP)  
76 (Santa Rita farm), located between coordinates 20°16'50" south latitude and 50°17'43" west  
77 longitude and 20°18'05" south latitude and 50°16'26" west longitude (Figure 1).

78


79 The climate of the region, according to the Koppen classification, is humid subtropical, Aw,  
80 with winter (dry and mild) and summer (hot and rainy) (Rolim et al., 2007). According to the  
81 Embrapa (2007) region is characterized by a period of 6 months of the year with water deficit  
82 and average temperature of 23.5°C or 74,3° F (Table 1).

83

84



85



86

87

**Figure 1: Location of the Santa Rita farm, Fernandópolis city, SP, Brazil**

88

89 According to Soil Survey Staff (1999) soils of Santa Rita farm consist of Ultisol. The  
 90 experimental design **used was a** completely randomized, with three treatments and five  
 91 **replications**, totaling 25 experimental plots. The treatments **were:** Natural vegetation area  
 92 (Figure 2); Area of cultivation of perennial crops (implanted 15 years ago, spacing 1 x 2 m,

93 area without management of manuring) (Figure 2) and Degraded pasture area (3 ha) (Figure  
94 2).

95  
96

**Table 1: Water balance of the Fernandópolis city, SP.**

Month	AT (°C)	AT (°F)	P (mm)	PE (mm)	WS (mm)	AE (mm)	DE (mm)	EXC (mm)
January	25.2	77.4	242	130	100	130	0	84
February	25.3	77.5	193	120	100	120	0	73
March	25.0	77.0	131	124	100	124	0	7
April	23.4	74.1	65	94	74	91	4	0
May	21.2	70.2	47	70	59	62	8	0
June	20.2	68.4	37	57	48	48	9	0
July	20.1	68.2	13	58	31	30	27	0
August	22.3	72.1	20	80	17	34	46	0
September	24.1	75.4	35	101	9	43	58	0
October	24.8	76.6	126	119	16	119	0	0
November	24.8	76.6	120	120	16	120	0	0
December	25.0	77.0	187	131	72	131	0	0
Totals	-	-	1.216	1.205	642	1.053	152	163
Average	23,5	74.2	-	-	-	-	-	-

97 NOTE: AT (average temperature); P (precipitation); PE (potential evapotranspiration); WS (water  
98 storage in the soil); AE (actual evapotranspiration); DE (disability); EXC (excess).  
99

100

101 The native vegetation is composed mainly by the respective species: *Stylosantes capitata*,  
102 *Talisia esculenta*, *Hymenaea stilbocarpa*, *Matayba elaeagnoides*, *Anadenanthera falcata*.  
103 *Peltophorum dubium*, *Tabebuia serratifolia*, *Mimosa caesalpiniaefolia*. As a floristic  
104 composition heterogeneous and rich in species.

105

106 In the area with perennial crop is basically composed by species of *Eucaliptus grandis*,  
107 without ground vegetation, i.e., as the area is already being cultivated for 15 years, the  
108 canopy of trees shades the whole area, hindering the onset of ground vegetation on site.  
109 The planting was done with cuttings. The sampling in this treatment was performed at 0.50  
110 m from the plant row. And the degraded area is cultivated by the grasses *Panicum*  
111 *maximum*.

112

113 For each treatment soil samples were obtained in duplicate for better data reliability.

114

115 Chemical soil properties were evaluated according to the methodology described by Raij &  
116 Quaggio (1983) and the composition of phosphorus, potassium, magnesium and calcium by  
117 extraction with an ion exchange resin were evaluated. The organic matter (OM) was  
118 determined by the colorimetric method and the pH, in calcium chloride, as well as the  
119 potential acidity (hydrogen + aluminum) at pH 7.0. Were calculated sums of bases (SB = Ca  
120 + Mg + K), cation exchange capacity (CEC = SB + (H + AL)) and base saturation (V% = (100  
121 x SB) / CEC). The samples were collected with the aid of exchange of mug, in three soil  
122 depths: 0.00-0.10; 0.10-0.20 and 0.20-0.40 m (Figure 3).

123



124

125

126

Natural vegetation.



Perennial crop – *Eucalyptus grandis*



127

128

Degraded pasture.

129

**Figure 2: View of the experimental area**

130

131

132

133

For data analysis, was performed the analysis of variance (F test) and applied the Tukey test (5%), for comparison of means, using SISVAR program (Ferreira, 2008).



Soil collected



Trench to collect soil

134 **Figure 3: Composite sample of soil was used for the determination of soil fertility and**  
 135 **collecting.**  
 136

137 **3. RESULTS AND DISCUSSION**

138

139 The results for the chemical properties of the soil are presented in Table 1. For all properties  
 140 and soil depths studied, the F test was significant at 5% of probability.

141

142 **Table 1: Effect of soil use and management on the chemical properties of an Ultisol.**

Soil property	Soil use and management and soil depth (A: 0.00-0.10; 0.10-0.20 and 0.20-0.40m)								
	Natural Vegetation			Perennial Crops			Degraded Pasture		
	A	B	C	A	B	C	A	B	C
OM (g dm <sup>-3</sup> )	18 a	18 a	16 a	14 a	13 c	13 b	16 ab	15 b	15 ab
P (mg dm <sup>-3</sup> )	18 a	15 a	10 a	6 c	4 c	3 b	14 b	12 b	11 a
pH	5,6 a	5,2 a	5,1 a	4,8 a	4,7 b	4,5 b	5,4a	5,4 a	5,3 a
K (mmolc dm <sup>-3</sup> )	1,6 a	1,5 a	1,3 a	1,1 b	1,1 b	0,8 b	1,3 b	1,2 b	1,3 a
Ca (mmolc dm <sup>-3</sup> )	17 a	16 a	14 a	6 c	5 c	5 c	12 b	11 b	10 b
Mg (mmolc dm <sup>-3</sup> )	6 a	6 a	5 a	3 b	2 c	2 b	5 a	4 b	4 a
H+Al (mmolc dm <sup>-3</sup> )	23 c	24 c	26 c	34 a	35 a	36 a	28 b	30 b	31 b
SB (mmolc dm <sup>-3</sup> )	24,6 a	23,4 a	20,6 a	10,1 c	8,1 c	7,9 c	18,3 b	15,9 b	15,3 b
CEC (mmolc dm <sup>-3</sup> )	47,6 a	47,5 a	45,7ab	44,1 b	44,1 c	43,9 b	47,3 a	46,2 b	46,6 a
V%	52 a	49 a	44 a	23 c	18 c	18 c	39 b	35 b	33 b

Soil property	F Test, SDM - significant difference minimum 5%, VC- variation coefficient (%) (A: 0.00-0.10; 0.10-0.20 and 0.20-0.40m)								
	F			SDM(5%)			VC(%)		
	A	B	C	A	B	C	A	B	C
OM (g dm <sup>-3</sup> )	5.471*	57.00*	7.63*	3.943	1.445	2.397	11.30	4.35	7.54
P (mg dm <sup>-3</sup> )	384.00*	134.08*	66.07*	1.352	2.184	2.314	4.92	9.67	13.34
pH	384.00*	13.00*	14.182*	31.912	0.433	0.479	15.71	3.92	4.45
K (mmolc dm <sup>-3</sup> )	16.28*	15.97*	35.48*	0.270	0.237	0.194	9.35	8.69	7.92
Ca (mmolc dm <sup>-3</sup> )	78.00*	166.50*	25.11*	2.704	1.770	1.198	10.69	7.78	5.82
Mg (mmolc dm <sup>-3</sup> )	12.92*	72.00*	11.19*	1.843	1.022	2.068	18.21	11.79	26.31
H+Al (mmolc dm <sup>-3</sup> )	111.76*	164.58*	18.00*	2.257	1.946	1.616	3.66	3.03	2.40
SB (mmolc dm <sup>-3</sup> )	668.38*	732.43*	35.29*	1.222	1.224	1.467	3.19	3.57	4.64
CEC (mmolc dm <sup>-3</sup> )	165.22*	225.44*	9.288*	0.654	0.495	1.976	0.65	0.50	2.01
V%	585.3*	24.21*	98.14*	0.817	1.371	1.777	1.00	1.84	2.58

143 abc Values with different superscripts are significantly different at (P< .05). \* Significant at 5%, ns no  
 144 significant.

145

146 For native vegetation, the surface layer (0-0,10m) was found the highest content of organic  
 147 material, and exchangeable ions (cations: Ca, K and Mg and anions: P). This behavior was  
 148 observed due to the presence of an equilibrium soil-plant-water. The nutrient input has as  
 149 main source, the mineralization of nutrients coming from the roots and leaf litter. This effect  
 150 is the result of greater accumulation due to greater species richness of this treatment. The  
 151 CEC of the soil in the same layer (0-0,10m) is almost entirely occupied by SB (Ca, Mg and  
 152 K) and at lower extent by the potential acidity. As the increases the sampling depth (0.10-  
 153 0.20 and 0.20-0.40m), the contents of cations and anions decreases, this behavior is  
 154 explained by the best influence of leaf litter and of the roots. With the smaller influence of  
 155 OM there is an increased potential acidity.

156

157 According to Souza & Alves (2003) who observed similar behavior that found in this work,  
 158 claim that management systems with little or no soil disturbance, tends to accumulate  
 159 exchangeable cations, phosphorus and organic matter. Was also observed the same  
 160 behavior in research conducted by Ferreira et al (2011). This accumulation of exchangeable  
 161 bases in the soil surface depth agrees with a lower potential acidity found in treatments  
 162 native vegetation and grassland. Regarding the treatment eucalyptus there is a greater  
 163 potential acidity and lower content of exchangeable bases, phosphorus and soil organic  
 164 matter. Identical behavior was verified by Souza & Alves (2003) and Ferreira et al. (2011).  
 165

166 For the cultivation of eucalyptus, there was lower contents of exchangeable cations and  
167 anions in relation to the native vegetation treatment of cerrado. In the surface layer (0-  
168 0.10m) the contents were higher than in the other layers (0,10-0,20 and 0.20-0.40m) for the  
169 treatment eucalyptus.

170  
171 The lowest concentration of Ca and Mg in K in eucalyptus treatment, is result of lower  
172 influence of roots and ground cover, due to the taproot system. Eucalyptus the growth of the  
173 root system of plants resulting seedlings or cuttings are distinct According Gaspar et al  
174 (2005) cuttings produce fewer primary roots, often with no tap root and have a shallower root  
175 system. And Fabião et al (1985) the turnover rate of fine roots in Eucalyptus stands is  
176 otherwise fairly high.

177  
178 In grasses (treatment degraded pasture) is the fibrous root system, influence in greater  
179 proportion. This effect of the grasses roots was verified by Bonini (2012). The effect of roots  
180 was also verified by Beutler & Centurion (2004) and Richart et al (2005).

181  
182 Soil management by man leads to soil depletion and degradation. The sustainable  
183 management (with low soil disturbance and vegetation cover) is recommending in Brazil,  
184 due to the hot and humid climate that leads the organic matter to highest mineralization  
185 rates.

186  
187 The highest levels of exchangeable bases in the treatment degraded pasture in relation to  
188 eucalyptus treatment is verified by the greatest influence of vegetation cover provided by  
189 *Brachiaria decumbens*. But these treatments obtained lower beneficial characteristics for the  
190 development of plants (low CEC, SB, pH, V% and high potential acidity) in relation to native  
191 vegetation.

192  
193 To soil organic matter, the highest contents found were to treatment native vegetation in any  
194 soil depth. The same behavior was observed for phosphorus, calcium, potassium and  
195 magnesium (table 1). This higher content of nutrients that are beneficial and essential for  
196 plant development was recorded in the treatment where there was no human intervention. In  
197 grasslands and eucalyptus treatments there was a gradual reduction, this due to greater soil  
198 disturbance and lower vegetation cover.

199  
200 Regarding the sum of bases, cation exchange capacity and base saturation, values found in  
201 this research, are considered low compared to the values considered average by Rajj et al.  
202 (1996). But even with values below of the optimal considered, was portrayed very well the  
203 difference between the treatments. The optimal saturation bases should be around 70% for  
204 most of the crops. The pH was an index with little variation among the treatments studied,  
205 this is due to no soil correction of cultivated areas and the native vegetation be slightly acidic  
206 (inherent soil characteristic). Bonini (2012) verified in Oxisol corrected higher pH than the  
207 native vegetation. Similar results were also found by Islam et al (2000), Carneiro et al (2009)  
208 and Souza & Alves (2003).

209

#### 210 **4. CONCLUSION**

211

212 The chemical properties of the soil were good indicators of soil quality, except for the pH  
213 index. Treatment with higher accumulation of organic matter, phosphorus, exchangeable  
214 cations was the native vegetation. The worst treatment was eucalyptus. The analyzes  
215 performed indicate that the management interfere with the chemical quality of the soil.

216

217

218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269

## REFERENCES

- Araújo, M.A.; Tormena, C.A.; Silva, A.P. Physical properties of a dystrophic Red Latosol (Oxisol) under crop cultivation and native forest. *R. Bras. Ci. Solo*, 2004; 28(2): 337-346. Portuguese.
- Araújo, R.; Goedert, W. J. & Marilusa, Lacerda, P.C. Soil Quality Under Different Uses And Native Cerrado. *R. Bras. Ci. Solo*, 2007; 31(1): 1099-1108. Portuguese.
- Beutler, A. N.; Centurion, J. F. Effect of soil compaction in root development and in soybean yield. *Pesq. Agropec. Bras.*, 2004; 39(6): 581-588. Portuguese.
- Bonini, C. S. B. Ecological restoration of soil beheaded under human intervention 17 years ago, Thesis (Graduate Production Systems) - Faculty of Engineering, São Paulo State University, Ilha Solteira, 2012. Portuguese.
- Brookes, P.C. The use of microbial parameters in monitoring soil pollution by heavy metals. *Biol. Fert. Soils*, 1995; 19(1): 269-279.
- Carneiro, M.A.C.; Souza, E.D.; Reis, E.F.; Pereira, H.S. & Azevedo, W.R. physical, chemical and biological properties of cerrado soil under different land use and tillage systems. *R. Bras. Ci. Solo*, 2009; 33(1): 147-157. Portuguese.
- Doran, J.W. & Parkin, T.B. Defining and assessing soil quality. In: Doran, J.W. & Coeman, D.C.; Bezdicsek, D.F & Stewart, B.A. Defining soil quality for sustainable environment. Madison, Soil Science Society of America, 1994. 3-21. (SSSA Special Publication, 35)
- EMBRAPA. Bank of climatic data from Brazil. Brasília: Embrapa Satellite Monitoring, 2007. Available: <http://www.bdclima.cnpm.embrapa.br/>. Accessed 23 november 2008. Portuguese.
- Ferreira, D. F.; SISVAR: a program for analysis and teaching of statistics. Symposium, 2008, 6(1): 36-41. Portuguese.
- Gaspar, M. J.; Borralho, N; Gomes, A. L. Comparison between field performance of cuttings and seedlings of *Eucalyptus globulus*. *Ann. For. Sci*, 2005, 62 (1): 837-841.
- Fabiao, A; Persson, H. A.; Steen, E.; Growth dynamics of superficial roots in Portuguese plantations of *Eucalyptus globulus* Labill studied with a mesh bag technique. *Plant and Soil*, 1985, 83 (2): 233-242.
- Ferreira, A N.K.F.; Souza, C.M.C.; Bastos, L.F.; Silva Junior, M.L.; Melo, V.S. Soil chemical properties under different land use systems and management in Pacajá (PA). In: Proceedings of the 9th Annual Scientific Seminar of Initiation. 2011, 1(1): 1-4. Portuguese.
- Islam, K.R. & Weil, R.R. Land use effects on soil quality in a tropical forest ecosystem of Bangladesh. *Agric. Ecosys. Environ.*, 2000, 79(1): 9-16.
- Leite, L. F. C; Mendonça, E. S., Machado, P. L. O. A., Matos, E. S. Total C and N storage and organic C pools of a Red-Yellow Podzolic under conventional and no tillage at the Atlantic Forest Zone, Southeastern Brazil. *Australian Journal Soil Research*, 2003; 41(1) 717-730.



270 Soil Survey Staff. Soil taxonomy: A basic system of soil classification for making and  
271 interpreting soil surveys. 2nd ed. Washington, Government Printing Office, 1999. (USDA,  
272 Handbook, 436).  
273  
274 Raj, B. V., Quaggio J. A. Methods of soil analysis for the purpose of fertility. Agronomic  
275 Institute of Campinas,1983. Portuguese.  
276  
277 Raj, B. V.; Cantarella, H.; Quaggio, J. A.; Furlani, A. M. C. Adubação orgânica. In: Raj, B.  
278 V.; Cantarella, H.; Quaggio, J. A.; Furlani, A. M. C. Liming and fertilizer recommendations for  
279 the State of São Paulo. 2nd ed. Campinas Agronomic Institute: IAC Foundation, 1996.  
280 Portuguese.  
281  
282 Richart, A.; Tavares Filho, J.; Brito, O. R.; Llanillo, R. F.; Ferreira, R. Soil compacting:  
283 causes and effects. Semina: Ciências Agrárias, 2005; 26(3): 321-344. Portuguese.  
284  
285 Rolim, G.S.; Camargo, M.B.P.; Lania, D.G.; Moraes, J.F.L. Climatic classification of Köppen  
286 and Thornthwaite systems and their applicability in the determination of agroclimatic zoning  
287 for the state of São Paulo, Brazil. Bragantia, 2007; 66(4): 711-720. Portuguese.  
288  
289 Souza, Z.M. & Alves, M.C. Chemical properties of a red cerrado latosol under different use  
290 and management conditions. Rev. Bras. Ci. Solo, 2003; 27(1):133-139. Portuguese.  
291