

The effect of soil use and management on the chemical properties of an Ultisol

12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

ABSTRACT

The use of vegetation cover to protect the soil and increase the soil organic matter content available to plants is a good practice to minimize loss of soil quality. The effect of soil management practices on the objective of this study was to evaluate the soil quality under different types of management. A completely randomised experimental design was used with three treatments as; area of natural vegetation; Area of perennial crop cultivation and degraded pasture area; with five replications. Chemical soil properties evaluated were: phosphorus, potassium, magnesium and calcium contents by extraction with ion exchanger resin. The content of organic matter was determined by the colorimetric method and the pH, in calcium chloride, as well as the acidity potential of the pH 7.0. The sums of bases, cation exchange capacity and base saturation were calculated. Under the treatment of natural vegetation the soil depths: 0.00-0.10; 0.10-0.20 and 0.20-0.40m. Data were analyzed statistically using the Tukey test at 5% for comparison of means. In the treatment, there was an accumulation of exchangeable bases, organic matter and phosphorus, as well as reducing potential acidity. The results show that soil use in spite of its management interferes with the chemical quality of the soil.

Keywords: organic matter, soil acidity, phosphorus.

1. INTRODUCTION

Sustainable use of natural resources, especially soil and water, has become a topic that is increasingly relevant, due to the increase of anthropogenic activities. Consequently there is, increases the concern with the quality and sustainable use of these resources (Araújo et al., 2007).

The introduction of agricultural systems to replace the forests causes an imbalance in the ecosystem in which the removal of the original vegetation and the deployment of crops, combined with inadequate management practices, lead to the disruption of the balance between the soil and the environment. In this way, the modification of soil chemical, physical and biological properties, limits the soils their agricultural use (Richart, et al., 2005).

The quality to attributes provides suitable conditions for the growth and development of plants and the maintenance of the diversity of organisms that inhabit the soil (Doran & Parkin, 1994). The current research challenge is the assessment of quality soil restored. According to Doran & Parkin (1994), it can be measured by quantifying to attributes, i.e., physical, chemical and biological properties have been used to monitor medium and long term changes, in the quality of the soil. Soil quality must be monitored to detect changes that may be mediated at a certain time and can be done on the farm or on a larger scale, such as watersheds and regions.

36

37 According to Araujo et al.,(2007), management practices and soil conservation and water
38 must be planned and executed attempting to maintain or even improve their attributes, in
39 order to increase the soil's ability to sustain a competitive biological productivity. The
40 conventional management systems (plowing and harrowing), results in intensified processes
41 of erosion and soil compaction which in the medium and long term, may lead to physical,
42 chemical and biological degradation of the soil (Leite et al., 2003; Araújo et al., 2004). In
43 these conditions, the absorption of nutrients available to the root system is hampered and
44 the amount of oxygen in the rhizosphere can become limiting in several metabolic processes
45 (Beulter & Centurion, 2004).

46

47 Islam & Weil (2000) proposed the use of data of physical, chemical and biological
48 characteristics of soil collected in an area of undisturbed natural forest, as a reference for
49 assembling a general index of soil quality. According to Brookes (1995), the relationship
50 between the physical, chemical and biological properties of the soil vary in time and space.
51 And these changes can be beneficial or harmful to production systems. Proper management
52 promotes gain in soil quality and crop productivity. Consequently, the variation of these
53 attributes, determined by management and soil use, and its evaluation are important for
54 better management aiming at sustainability of the system (Carneiro et al., 2009).

55

56 Works performed by Souza & Alves (2003) in dystrophic Oxisol, show that soil management
57 systems with less or no soil disturbance led to significant accumulation of organic matter,
58 phosphorus, potassium, magnesium and calcium and decrease of exchangeable aluminum
59 and potential acidity. With soil disturbance as a result of cropping, the effects were a
60 reduction of the levels of organic matter, phosphorus, potassium, calcium and magnesium
61 and increased potential acidity and exchangeable aluminum.

62

63 Studies conducted in Oxisol, in the region north of the state of Pará, found that the sum of
64 bases and cation exchange capacity (CEC) were the most sensitive variables to changes in
65 the use and management (primary forest, explored forest, grassland, scrub, jiquira of typical
66 Brazilian vegetation), and the pH and potential acidity and soil pH were influenced by the
67 replacement of forest by pasture (Ferreira et al., 2011).

68

69 The aim of this work was to evaluate the chemical quality of the soil of an Ultisol under
70 different uses and management.

71

72 2. MATERIALS AND METHODS

73

74 The experiment was conducted at the experimental area of farm Teaching and Research at
75 the University Camilo Castelo Branco - UNICASTELO, Fernandópolis city, São Paulo (SP)
76 (Santa Rita farm), located between coordinates 20°16'50" south latitude and 50°17'43" west
77 longitude and 20°18'05" south latitude and 50°16'26" west longitude (Figure 1).

78


79 The climate of the region, according to the Koppen classification, is humid subtropical, Aw,
80 with winter (dry and mild) and summer (hot and rainy) (Rolim et al., 2007). According to the
81 Embrapa (2007) region is characterized by a period of 6 months of the year with water deficit
82 and average temperature of 23.5°C or 74,3° F (Table 1).

83

84



85



86

87

Figure 1: Location of the Santa Rita farm, Fernandópolis city, SP, Brazil

88

89 According to Soil Survey Staff (1999) soils of Santa Rita farm consist of Ultisol. The
 90 experimental design **used was a** completely randomized, with three treatments and five
 91 **replications**, totaling 25 experimental plots. The treatments **were:** Natural vegetation area
 92 (Figure 2); Area of cultivation of perennial crops (implanted 15 years ago, spacing 1 x 2 m,

93 area without management of manuring) (Figure 2) and Degraded pasture area (3 ha) (Figure
94 2).

95
96

Table 1: Water balance of the Fernandópolis city, SP.

Month	AT (°C)	AT (°F)	P (mm)	PE (mm)	WS (mm)	AE (mm)	DE (mm)	EXC (mm)
January	25.2	77.4	242	130	100	130	0	84
February	25.3	77.5	193	120	100	120	0	73
March	25.0	77.0	131	124	100	124	0	7
April	23.4	74.1	65	94	74	91	4	0
May	21,2	70.2	47	70	59	62	8	0
June	20.2	68.4	37	57	48	48	9	0
July	20.1	68.2	13	58	31	30	27	0
August	22.3	72.1	20	80	17	34	46	0
September	24.1	75.4	35	101	9	43	58	0
October	24.8	76.6	126	119	16	119	0	0
November	24.8	76.6	120	120	16	120	0	0
December	25.0	77.0	187	131	72	131	0	0
Totals	-	-	1.216	1.205	642	1.053	152	163
Average	23,5	74.2	-	-	-	-	-	-

97 NOTE: AT (average temperature); P (precipitation); PE (potential evapotranspiration); WS (water
98 storage in the soil); AE (actual evapotranspiration); DE (disability); EXC (excess).

99

100

101 The native vegetation is composed mainly by the respective species: *Stylosantes capitata*,
102 *Talisia esculenta*, *Hymenaea stilbocarpa*, *Matayba elaeagnoides*, *Anadenanthera falcata*.
103 *Peltophorum dubium*, *Tabebuia serratifolia*, *Mimosa caesalpiniaefolia*. As a floristic
104 composition heterogeneous and rich in species.

105

106 In the area with perennial crop is basically composed by species of *Eucaliptus grandis*,
107 without ground vegetation, i.e., as the area is already being cultivated for 15 years, the
108 canopy of trees shades the whole area, hindering the onset of ground vegetation on site.
109 The planting was done with cuttings. The sampling in this treatment was performed at 0.50
110 m from the plant row. The degraded area is cultivated by the grasses *Panicum maximum*.

111

112 The samples were collected with the aid of exchange of mug, in three soil depths: 0.00-0.10;
113 0.10-0.20 and 0.20-0.40 m (Figure 3). For each treatment soil samples were obtained in
114 duplicate for better data reliability, in may to 2012.

115

116 Chemical soil properties were evaluated according to the methodology described by Raij &
117 Quaggio (1983) and the composition of phosphorus, potassium, magnesium and calcium by
118 extraction with an ion exchange resin were evaluated. The organic matter (OM) was
119 determined by the colorimetric method and the pH, in calcium chloride, as well as the
120 potential acidity (hydrogen + aluminum) at pH 7.0. Were calculated sums of bases (SB = Ca
121 + Mg + K), cation exchange capacity (CEC = SB + (H + AL)) and base saturation (V% = (100
122 x SB) / CEC).

123



124
125
126

Natural vegetation.



Perennial crop – *Eucalyptus grandis*



127
128

Degraded pasture.

129

Figure 2: View of the experimental area

130
131
132
133

For data analysis, was performed the analysis of variance (F test) and applied the Tukey test (5%), for comparison of means, using SISVAR program (Ferreira, 2008).



Soil collected



Trench to collect soil

134 **Figure 3: Composite sample of soil was used for the determination of soil fertility and**
 135 **collecting.**
 136

137 **3. RESULTS AND DISCUSSION**

138
 139 The results for the chemical properties of the soil are presented in Table 1. For all properties
 140 and soil depths studied, the F test was significant at 5% of probability.
 141

142 **Table 1: Effect of soil use and management on the chemical properties of an Ultisol.**

Soil property	Soil use and management and soil depth (A: 0.00-0.10; 0.10-0.20 and 0.20-0.40m)								
	Natural Vegetation			Perennial Crops			Degraded Pasture		
	A	B	C	A	B	C	A	B	C
OM (g dm ⁻³)	18 a	18 a	16 a	14 a	13 c	13 b	16 ab	15 b	15 ab
P (mg dm ⁻³)	18 a	15 a	10 a	6 c	4 c	3 b	14 b	12 b	11 a
pH	5,6 a	5,2 a	5,1 a	4,8 a	4,7 b	4,5 b	5,4a	5,4 a	5,3 a
K (mmolc dm ⁻³)	1,6 a	1,5 a	1,3 a	1,1 b	1,1 b	0,8 b	1,3 b	1,2 b	1,3 a
Ca (mmolc dm ⁻³)	17 a	16 a	14 a	6 c	5 c	5 c	12 b	11 b	10 b
Mg (mmolc dm ⁻³)	6 a	6 a	5 a	3 b	2 c	2 b	5 a	4 b	4 a
H+Al (mmolc dm ⁻³)	23 c	24 c	26 c	34 a	35 a	36 a	28 b	30 b	31 b
SB (mmolc dm ⁻³)	24,6 a	23,4 a	20,6 a	10,1 c	8,1 c	7,9 c	18,3 b	15,9 b	15,3 b
CEC (mmolc dm ⁻³)	47,6 a	47,5 a	45,7ab	44,1 b	44,1 c	43,9 b	47,3 a	46,2 b	46,6 a
V%	52 a	49 a	44 a	23 c	18 c	18 c	39 b	35 b	33 b

F Test, SDM - significant difference minimum 5%, VC- variation coefficient (%) (A: 0.00-0.10; 0.10-0.20 and 0.20-0.40m)

Soil property	F			SDM(5%)			VC(%)		
	A	B	C	A	B	C	A	B	C
OM (g dm ⁻³)	5.471*	57.00*	7.63*	3.943	1.445	2.397	11.30	4.35	7.54
P (mg dm ⁻³)	384.00*	134.08*	66.07*	1.352	2.184	2.314	4.92	9.67	13.34
pH	384.00*	13.00*	14.182*	31.912	0.433	0.479	15.71	3.92	4.45
K (mmolc dm ⁻³)	16.28*	15.97*	35.48*	0.270	0.237	0.194	9.35	8.69	7.92
Ca (mmolc dm ⁻³)	78.00*	166.50*	25.11*	2.704	1.770	1.198	10.69	7.78	5.82
Mg (mmolc dm ⁻³)	12.92*	72.00*	11.19*	1.843	1.022	2.068	18.21	11.79	26.31
H+Al (mmolc dm ⁻³)	111.76*	164.58*	18.00*	2.257	1.946	1.616	3.66	3.03	2.40
SB (mmolc dm ⁻³)	668.38*	732.43*	35.29*	1.222	1.224	1.467	3.19	3.57	4.64
CEC (mmolc dm ⁻³)	165.22*	225.44*	9.288*	0.654	0.495	1.976	0.65	0.50	2.01
V%	585.3*	24.21*	98.14*	0.817	1.371	1.777	1.00	1.84	2.58

143 abc Values with different superscripts are significantly different at (P< .05). * Significant at 5%, ns no
 144 significant.

145

146 For native vegetation, the surface layer (0-0,10m) was found the highest content of organic
 147 material, and exchangeable ions (cations: Ca, K and Mg and anions: P). This behavior was
 148 observed due to the presence of an equilibrium soil-plant-water. The nutrient input has as
 149 main source, the mineralization of nutrients coming from the roots and leaf litter. This effect
 150 is the result of greater accumulation due to greater species richness of this treatment. The
 151 CEC of the soil in the same layer (0-0,10m) is almost entirely occupied by SB (Ca, Mg and
 152 K) and at lower extent by the potential acidity. As the increases the sampling depth (0.10-
 153 0.20 and 0.20-0.40m), the contents of cations and anions decreases, this behavior is
 154 explained by the best influence of leaf litter and of the roots. With the smaller influence of
 155 OM there is an increased potential acidity.

156

157 According to Souza & Alves (2003) who observed similar behavior that found in this work,
 158 claim that management systems with little or no soil disturbance, tends to accumulate
 159 exchangeable cations, phosphorus and organic matter. The same behaviour was also
 160 observed in research by Ferreira et al (2011). This accumulation of exchangeable bases in
 161 the soil surface depth agrees with a lower potential acidity found in treatments native
 162 vegetation and grassland. Regarding the treatment eucalyptus there is a greater potential
 163 acidity and lower content of exchangeable bases, phosphorus and soil organic matter.
 164 Identical behavior was verified by Souza & Alves (2003) and Ferreira et al. (2011).
 165

166 For the cultivation of eucalyptus, there was lower contents of exchangeable cations and
167 anions in relation to the native vegetation treatment of cerrado. In the surface layer (0-
168 0.10m) the contents were higher than in the other layers (0,10-0,20 and 0.20-0.40m) for the
169 treatment eucalyptus.

170

171 The lowest concentration of Ca and Mg in K in eucalyptus treatment, is result of lower
172 influence of roots and ground cover, due to the taproot system. In Eucalyptus, the growth of
173 the root system of plants resulting of seedlings or cuttings are distinct. According Gaspar et
174 al (2005) cuttings produces less, primary roots, frequently without main root and have a
175 more shallow root system. Fabião et al (1985) states that the amount of fine roots in
176 Eucalyptus is very low.

177

178 In grasses (treatment degraded pasture), the fibrous root system, influence in greater
179 proportion the soil quality in the surface depth. This effect of the grasses roots was verified
180 by Bonini (2012). The effect of roots was also verified by Beutler & Centurion (2004) and
181 Richart et al (2005).

182

183 Soil management by man leads to soil depletion and degradation. The sustainable
184 management (with low soil disturbance and vegetation cover) is recommending in Brazil,
185 due to the hot and humid climate that leads the organic matter to highest mineralization
186 rates.

187

188 The highest levels of exchangeable bases in the treatment degraded pasture in relation to
189 eucalyptus treatment is verified by the greatest influence of vegetation cover provided by
190 *Brachiaria decumbens*. But these treatments have characteristics fewer beneficial for the
191 development of plants (low CEC, SB, pH, V% and high potential acidity) in relation to native
192 vegetation.

193

194 To soil organic matter, the highest contents found were to treatment native vegetation in any
195 soil depth. The same behavior was observed for phosphorus, calcium, potassium and
196 magnesium (table 1). This higher content of nutrients that are beneficial and essential for
197 plant development was recorded in the treatment where there was no human intervention. In
198 grasslands and eucalyptus treatments there was a gradual reduction, this due to greater soil
199 disturbance and lower vegetation cover.

200

201 Regarding the sum of bases, cation exchange capacity and base saturation, values found in
202 this research, are considered low compared to the values considered average by Rajj et al.
203 (1996). But even with values below of the optimal considered, was portrayed very well the
204 difference between the treatments. The optimal saturation bases should be around 70% for
205 most of the crops. The pH was an index with little variation among the treatments studied,
206 this is due to no soil correction of cultivated areas and the native vegetation be slightly acidic
207 (inherent soil characteristic). Bonini (2012) verified in Oxisol corrected higher pH than the
208 native vegetation. Similar results were also found by Islam et al (2000), Carneiro et al (2009)
209 and Souza & Alves (2003).

210

211 4. CONCLUSION

212

213 The chemical properties of the soil were good indicators of soil quality, except for the pH
214 index. Treatment with higher accumulation of organic matter, phosphorus, exchangeable
215 cations was the native vegetation. The worst treatment was eucalyptus. The analyzes
216 performed indicate that the management interfere with the chemical quality of the soil.

217

218

219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270

REFERENCES

- Araújo, M.A.; Tormena, C.A.; Silva, A.P. Physical properties of a dystrophic Red Latosol (Oxisol) under crop cultivation and native forest. *R. Bras. Ci. Solo*, 2004; 28(2): 337-346. Portuguese.
- Araújo,R.; Goedert, W. J. & Marilusa, Lacerda, P.C. Soil Quality Under Different Uses And Native Cerrado. *R. Bras. Ci. Solo*, 2007; 31(1): 1099-1108. Portuguese.
- Beutler, A. N.; Centurion, J. F. Effect of soil compaction in root development and in soybean yield. *Pesq. Agropec. Bras.*, 2004; 39(6): 581-588. Portuguese.
- Bonini, C. S. B. Ecological restoration of soil beheaded under human intervention 17 years ago, Thesis (Graduate Production Systems) - Faculty of Engineering, São Paulo State University, Ilha Solteira, 2012. Portuguese.
- Brookes, P.C. The use of microbial parameters in monitoring soil pollution by heavy metals. *Biol. Fert. Soils*, 1995; 19(1): 269-279.
- Carneiro, M.A.C.; Souza, E.D.; Reis, E.F.; Pereira, H.S. & Azevedo, W.R. physical, chemical and biological properties of cerrado soil under different land use and tillage systems. *R. Bras. Ci. Solo*, 2009; 33(1): 147-157. Portuguese.
- Doran, J.W. & Parkin, T.B. Defining and assessing soil quality. In: Doran, J.W. & Coeman, D.C.; Bezdicsek, D.F & Stewart, B.A. Defining soil quality for sustainable environment. Madison, Soil Science Society of America, 1994. 3-21. (SSSA Special Publication, 35)
- EMBRAPA. Bank of climatic data from Brazil. Brasília: Embrapa Satellite Monitoring, 2007. Available: <http://www.bdclima.cnpm.embrapa.br/>. Accessed 23 november 2008. Portuguese.
- Ferreira, D. F.; SISVAR: a program for analysis and teaching of statistics. Symposium, 2008, 6(1): 36-41. Portuguese.
- Gaspar, M. J.; Borralho, N; Gomes, A. L. Comparison between field performance of cuttings and seedlings of *Eucalyptus globulus*. *Ann. For. Sci*, 2005, 62 (1): 837-841.
- Fabiao, A; Persson, H. A.; Steen, E.; Growth dynamics of superficial roots in Portuguese plantations of *Eucalyptus globulus* Labill studied with a mesh bag technique. *Plant and Soil*, 1985, 83 (2): 233-242.
- Ferreira, A N.K.F.; Souza, C.M.C.; Bastos, L.F.; Silva Juniiior, M.L.; Melo, V.S. Soil chemical properties under different land use systems and management in Pacajá (PA). In: Proceedings of the 9th Annual Scientific Seminar of Initiation. 2011, 1(1): 1-4. Portuguese.
- Islam, K.R. & Weil, R.R. Land use effects on soil quality in a tropical forest ecosystem of Bangladesh. *Agric. Ecosys. Environ.*, 2000, 79(1): 9-16.
- Leite, L. F. C; Mendonça, E. S., Machado, P. L. O. A., Matos, E. S. Total C and N storage and organic C pools of a Red-Yellow Podzolic under conventional and no tillage at the Atlantic Forest Zone, Southeastern Brazil. *Australian Journal Soil Research*, 2003; 41(1) 717-730.

271 Soil Survey Staff. Soil taxonomy: A basic system of soil classification for making and
272 interpreting soil surveys. 2nd ed. Washington, Government Printing Office, 1999. (USDA,
273 Handbook, 436).
274
275 Raj, B. V., Quaggio J. A. Methods of soil analysis for the purpose of fertility. Agronomic
276 Institute of Campinas,1983. Portuguese.
277
278 Raj, B. V.; Cantarella, H.; Quaggio, J. A.; Furlani, A. M. C. Adubação orgânica. In: Raj, B.
279 V.; Cantarella, H.; Quaggio, J. A.; Furlani, A. M. C. Liming and fertilizer recommendations for
280 the State of São Paulo. 2nd ed. Campinas Agronomic Institute: IAC Foundation, 1996.
281 Portuguese.
282
283 Richart, A.; Tavares Filho, J.; Brito, O. R.; Llanillo, R. F.; Ferreira, R. Soil compacting:
284 causes and effects. Semina: Ciências Agrárias, 2005; 26(3): 321-344. Portuguese.
285
286 Rolim, G.S.; Camargo, M.B.P.; Lania, D.G.; Moraes, J.F.L. Climatic classification of Köppen
287 and Thornthwaite systems and their applicability in the determination of agroclimatic zoning
288 for the state of São Paulo, Brazil. Bragantia, 2007; 66(4): 711-720. Portuguese.
289
290 Souza, Z.M. & Alves, M.C. Chemical properties of a red cerrado latosol under different use
291 and management conditions. Rev. Bras. Ci. Solo, 2003; 27(1):133-139. Portuguese.
292