

Original Research Article

Title- Effect of age at harvest and leaf position on the yield and nutritional composition of *Celosia argentea* L.

Abstract

A pot experiment was carried out at the nursery site of the Department of Crop Production, Federal University of Technology, Minna, Niger state, Nigeria. The study aimed at determining the effect of age of celosia plant at harvest on the yield and nutritional composition of the plant as well as the concentration of nutrients at different leaf positions. The experiment was a 3x3 factorial combination of three harvest periods (5, 7 and 9 weeks after sowing) and three leaf positions on the mother plant (upper, middle and basal) arranged in a completely randomized design. Harvested leaves were analyzed for the nutritional composition. The results showed that the whole plant fresh weight, varied significantly ($p < 0.05$) with the age of plant at harvest, having the maximum and the minimum values at 9 WAS (266.19 g/pot) and 5 WAS (96.12g/pot) respectively. The leaf fresh weight and leaf dry weight followed the same trend with the whole plant fresh weight. Ca, K, Fe, and Vit C content were significantly higher at 9WAS. Crude protein and Na reduced significantly ($p < 0.05$) with the age of the plant with the highest values recorded at 5WAS. Zn was highest at 7WAS. The Mg content was not significantly affected by the age at harvest. The middle leaves had significant higher content of Mg and Vit. C when compared to the basal leaves but there was no significant difference between the values obtained in upper and middle leaves. Significant ($p < 0.05$) higher values of Ca, Fe, and crude protein were recorded in the basal leaves. There was no significant difference in the values of K, P and Na obtained at the different leaf positions.

Key words: *Celosia argentea*, leaf positions, age at harvest, nutrients, yield

1.0 Introduction

Celosia argentea is a tropical annual leafy vegetable, and a leading leafy vegetable in SouthWestern Nigeria where it is known as 'Sokoyokoto'. The leaves and tender stems are cooked into soups. It is rich in protein, minerals and vitamins. It had been documented that the nutritional composition of *Celosia argentea* per 100 g edible portion is 83.8 g water; 185kj energy; 4.7 g protein; 0.7 g fat; 7.3 g carbohydrate; 1.8 g fibre ; 260 mg Ca; 43 mg P and 7.8 mg Fe, respectively (2). However, the composition of *Celosia argentea* is strongly influenced by environmental factors such as soil fertility, fertilizer application and age of the plant at harvest (2). Various methods had been used to harvest vegetables; they could be uprooted or ratooned in such a way that the lower leaves are left unharvested. Even when the whole plant parts are harvested, some people do not consume the lower leaves (older leaves) believing that it is too fibrous and less nutritious when compared to the upper leaves (younger leaves). The mineral content in the different plant tissues is related to their mobility in the plant. In conditions of mineral deficiency, some nutrients may be translocated from the mature leaves and fruits to the younger

leaves(3).Some nutrients are relatively immobile in plants and cannot be easily redistributed to younger leaves or other parts(4, 5, 6) thus: making the concentration of such nutrients higher in some plant part than the other.In view of the above, this research was carried out with the aim of determining the best age to harvest the plant to get the highest yield and optimum nutrients as well as the leaf position in which the derivable nutritional potential is highest.

2.0 Materials and Methods

The pot experiment was carried out at the Horticulture Nursery of Federal University of Technology Minna, Niger state in the raining season of 2013. It was a 3x3 factorial experiment arranged in completely randomized design. The treatments were 3 harvest periods: 5, 7 and 9 weeks after sowing and 3 leaf positions (upper, middle and basal leaves). The treatments were replicated three times. Each pot was filled with 8kg top soil. Four seeds of TLV8 variety were sown per pot and at two weeks after planting, the seedlings were thinned to two per pot.NPK 20:10:10 fertilizer was applied at the rate of 80kg N ha⁻¹, 40kgP₂O₅ ha⁻¹ and 40kg K₂O ha⁻¹ at two weeks after planting. Weeds werehand-picked whenever noticed. The plants were harvested at the sampling period stated above and fresh weights were taken after which they are separated into the upper, middle and the basal leaves.The leaves were dried inan oven at 65⁰C till constant weight was obtained to get the leaf dry weight and were subsequentlyanalyzed for protein, fat, carbohydrate, crude fibre, Vit. C and mineral elements(Fe, Mg, Zn, Ca, P, Na and K).

The mineral elements (Fe, Mg, Ca, Na and K)in the test samples were determined by digesting sample in mixture of concentrated HNO₃ and perchloric acid and read using atomic absorption spectrophotometer. Flame photometer was used for Na and K only. The P was determined using the molybdate method and quantified using a spectrophotometer. The ascorbic acid concentration in thesamples was determined by 2, 6-dichlorophenol indophenol titrimetric method. The crude protein was determined based on total N content by Kjeldahl method (1).All the datacollected were subjected toanalysis of variance (ANOVA) using version 9.0 of SAS (GLM procedure).Treatment means were separated using the least significant difference where significant differencesoccurred at 5% level of probability.

3.0 Result and Discussion

Table 1 reveals that the yield obtained (whole plant fresh weight, leaf fresh weight and leaf dry weight) increased with the plantage and the highest value was recorded at 9 weeks. This could be attributed to dry matter accumulation with increase in age.Several author have reported that there is increase in dry matter yield as plant age. (7, 8). However, the difference between the yield values obtained at 5 and 7 WASand between 7 and 9 WAS were not statistically different.

Table 1. Yield values of *Celosia argentea* at different harvesting period

Harvesting periods	Whole plant fresh weight	Leaf fresh weight	Leaf dry weight
(WAS)	(g/pot)	(g/pot)	(g/pot)
5	96.12	36.18	3.03
7	173.94	56.94	3.93
9	266.19	58.83	5.71
LSD	169.65	22.53	1.61

The result of the effect of the age at harvest and leaf position on the nutritional content of celosia are presented in Table 2. The result shows that there were significant differences in calcium content with respect to the plantage. The highest value of calcium was recorded at 9WAP. This is in agreement with the result obtained for *Amaranthus* by (9) who recorded the highest amount of calcium at the highest sampling period (60 DAP). Calcium content was significantly higher in basal leaves than the other two leaf positions. This value (146.60mg/100g) obtained in *Celosia argentea* doubled the amount (42-62mg/100g) recorded for different *Amaranthus species* reported by (9). This confirms the fact that *Celosia argentea* is rich in calcium(10). The value obtained in this study is still far below the recommended dietary allowance of 1000-1200mg/ day (11).(12) also observed that the highest amount of calcium was recorded in the basal leaves. This could be because calcium is immobile (non-translocatable) within plants and remains in the older tissue throughout the growing season. This is why the deficiency symptoms of Ca appears first in the young growing part of the plant (6).There was no significant difference between the amount of calciumrecordedin upper and middleleaves.

The age of the plant at harvest did not contribute significantly to the variation in Magnesium content recorded in the leaves. The magnesium value of the upper and the middle leaves were at par and were both significantly higher than the value for the lower leaves. This confirms the fact that Mg is withdrawn from ageing leaves due to its highly mobile nature (5). (13), observed no significant difference between the values of Magnesium obtained at the basal, middle and upper leaf position of *Hibiscus sabdariffa* plant.

The amount of K recorded in plant harvested at 9WAS (94.94mg/100g)was significantly higher than those obtained at 5 and 7WASwhich were at par.(14), recorded the highest amount of K at

112 6WAS in *Amaranthus*. There was no significant difference between the values of K recorded at
113 the different leaf positions.

114 The Fe content increased with the age of the plant. This is in agreement with the report of (15).
115 The highest value of Fe (38.98mg/100g) was obtained at 9WAS but was statistically similar to
116 the value obtained at 7WAS. The basal leaves contained significantly more Fe than other leaf
117 positions. This may be because Fe is relatively immobile in plant (5). (16) also recorded the
118 highest Fe content (27.53mg/kg) in the basal leaves of *Amaranthus cruentus* and with no
119 significant difference between the values recorded in the upper and middle leaves.

120 Phosphorus value significantly decreased with the age of the plant with 30.17mg/100g,
121 19.87mg/100g and 17.36 mg/100g recorded at 5, 7 and 9WAS respectively. (14) recorded the
122 highest P content at 4WAS (160mg/100g) beyond which the values declined
123 in *Amaranthus cruentus*. There was no significant difference between the amounts of P recorded at
124 the different leaf positions. This may be attributed to the fact that phosphate is easily
125 redistributed in most plants from one organ to another (4).

126 Significantly higher amount of Na was recorded at 5WAS (18.90mg/100g) compared to the
127 values obtained at 7 (16.74mg/100g) and 9 WAS (17.33mg/100g). There was no significant
128 difference between the values obtained at 7 and 9 WAS. There was no significant difference in
129 the amount of Na recorded in the three leaf positions. (12) also reported similar findings
130 in *Telfaria occidentalis*. The basal leaves of *Hibiscus sabdariffa* were however reported by (13) to
131 contain significantly higher value (3.38mg/kg) of the mineral than the middle and the upper
132 leaves. Though the values obtained in this study is low when compared with the recommended
133 dietary allowance of 2300mg/day (11) but table salt is the primary source of this mineral. Intake
134 of a teaspoon of salt per day is capable of supplying the recommended rate of Na.

135

136 **Table 2.** Effect of the age at harvest and positions of leaf on the nutritional content of *celosia argentea*

	Ca	Mg	K	Fe	PO ₄ Na	C.P	Fat	Vit.C	Zn	
Treatments	Mg/100g	Mg/100g	Mg/100g	Mg/100g	Mg/100g	Mg/100g	Mg/100g	Mg/100g	Mg/100g	Mg/100g
Age (WAS)										
5	134.52	21.12	79.74	26.68	30.17	18.90	3.20	2.70	27.92	3.13
7		115.51	23.17	74.84	35.41	19.89	16.74	2.90	3.02	35.00
9		144.97	24.43	94.94	38.98	17.36	17.33	2.64	2.61	38.10

LSD	11.77	NS	10.19	4.05	2.37	1.10	0.30	NS	2.60	0.51	
Position											
Upper leaves	132.04		24.33	90.65	30.78	23.76	17.94	2.66	2.61	34.44	3.72
Middle leaves	123.31		24.41	80.52	29.20	21.77	18.32	2.81	3.06	35.89	3.63
Basal leaves	146.60		20.69	81.44	38.38	23.18	17.38	3.34	3.06	29.67	3.81
LSD	13.59		3.50	NS	4.67	NS	NS	0.34	NS	3.01	NS
Interaction											
(Age x position)	NSNSNSNSNSNSNSNSNSNSNS										

C.P- Crude protein

NS- Not significant

WAS- Weeks after sowing

137

138 Crude protein reduced with the age of the plant. The value obtained (3.2g/100g) at 5WAS was
139 significantly higher than at 7 and 9 WAS. There was no significant difference between the values
140 obtained at 7 and 9 WAS. This concurs with the reports of (7, 14, 17). (18) observed that crude
141 protein content increased from 4WAS till 7 WAS in *Sesamum radiatum* leaves after which there
142 was a decline in amount till 10 WAS. The basal leaves had the highest crude protein content
143 (3.34g/100g) which was significantly higher than the other positions which were at par. The
144 highest value of crude protein obtained in the basal leaves (3.34g/100g) in this study is lower than
145 the values obtained in *Amaranthus cruentus* leaves (23%) as reported by (19). This confirms the
146 assertion of (9) that *Amaranthus* is higher in protein than *Celosia*. (2) reported that the amount of
147 protein found in *Celosia* was 4.7g/100g. Varietal factors and the environment could also
148 contribute to the variation in the value of crude protein obtained.

149 There was no significant difference between the fats amount obtained at the different harvesting
150 periods and the different leaf positions. This is in line with the report of (18) who reported that
151 the age of plant did not have any effect on the fat content of *Sesamum radiatum* leaves.

152 Vitamin C (Ascorbic acid) content increased progressively and significantly with age. The values
153 recorded at 5, 7 and 9WAS were 27.92mg/100g, 35.00mg/100g and 38.10mg/100g
154 respectively. The values obtained for both upper and middle leaves were statistically similar but

significantly higher than the value for basal leaves. Musa (2012) recorded the highest Vit. C content in the middle leaves of *Hibiscus sabdariffa*. The value of Vit. C obtained implies that if 200g of *Celosia* is eaten, it could supply the daily recommended daily allowance of 75mg/day (11) if minimally processed. This confirms the assertion of (10 and 20) that *Celosia* is a good source of Vit. C.

The value of zinc (4.03mg/100g) recorded at 7WAS was significantly higher than those at 5 and 9 WAS which were similar statistically. There was no significant difference between the values of zinc obtained at the different leaf position. This could be as a result of the fact that the mineral is highly mobile and is found in every part of the plant (4). (13) also observed that leaf position had no significant effect on the zinc content of *Hibiscus sabdariffa*. However, in *Amaranthus cruentus*, (16) recorded the highest value (0.11mg/kg) in the middle leaves. This value obtained in *Amaranthus cruentus* is low compared to the value obtained in *Celosia argentea*. This suggests that *Celosia argentea* is a moderately rich source of zinc. Deficiency of this mineral could cause growth retardation and poor sexual development (11)

4.0 Conclusion

Harvesting *Celosia* at 9 weeks after sowing seems better with respect to the yield obtained and the nutrient obtainable from the leaves at this age. However, P, Na and Crude Protein were significantly higher in leaves harvested at 5WAS. The lower leaves have significant higher levels of Ca, Fe and crude protein.

5.0 References

(1) Association of Analytical Chemists. 2000. *Official Methods of Analysis*. 17th ed. Gaithersburg, Maryland, USA, AOAC International.

(2) Denton, O.A, 2004. *Celosia argentea* L. In: Grubben G.J.H and Denton, O.A. Plant Resources of Tropical Africa 2. Vegetables. PROTA Foundation, Wageningen, Netherlands/ Backhuys. Leiden, Netherlands/CTA, Wageningen, Netherlands. Pp 167-171.

(3) Giuffrida F., Martorana M. and Leonardi C. 2009. How sodium chloride concentration in the nutrient solution influences the mineral composition of tomato leaves and fruits. *HortScience*. 44 (3):707-711.

(4) Taiz L. and Zeiger E. 2002. Plant physiology. Sinauer Associates, Inc. Sunderland. Massachusetts. 3rd edition. Pp 370-372.

- 189 (5) Hocmuth G., Maynard D., Vavrina C., Hanlon E., Simonne E. 2004. Plant tissue analysis and
190 interpretation for vegetable crops in Florida: University of Florida's Institute of Food and
191 Agricultural Sciences (UF/IFAS). Pp. 1-79.
192
- 193 (6) Fageria N.K. 2009. The use of nutrients in crop plants. CRC Press, Taylor and Francis group.
194 USA. Pp. 165-173.
195
- 196 (7) Collar C., Wright S., Robinson P. and Putnam D. Effect of harvest timing on yield and quality
197 of small grain forage. In: Proceedings, National Alfalfa Symposium: 13-15 December, 2004, San
198 Diego, CA, UC Cooperative Extension University of California. Available: <http://alfalfa.ucdavis.edu>
- 199 (8) Mahala A.G., Amasiab S.O., Yousif A.M. and Elsadig A. 2012. Effect of Plant age on DM
200 yield and nutritive value of some leguminous plants (*Cyamopsis tetragonoloba*, *Lablab*
201 *purpureus* and *Clitoria* (*Clitoria ternatea*)). International Research Journal of Agricultural Science and
202 Soil Science. 2(12): 502-508. Available: <http://www.interesjournals.org/IRJAS>
203
- 204 (9) Albert T. Modi . 2007. Growth temperature and plant age influence on nutritional quality of
205 *Amaranthus* leaves and seed germination capacity. Available: <http://www.wrc.org.za>
- 206 (10) Sato. 2002. Evaluation of antioxidant activity of indigenous vegetable from South and South
207 East Asia. In: JIRCAS research highlights, 10-11. Ohwash, tsukuba, Ibaraba, Japan. Available
208 : <http://www.jircas.affrc.go.jp/English/publication/highlights/index.html>
- 209 (11) Wardlaw G.M. and Smith A.M. 2011. Contemporary nutrition. Eight edition. Mc Graw-hill,
210 New York. Pp. 288-366.
211
- 212 (12) Musa Amanabo and Ogbadoyi Emmanuel O. 2012. Effect of Plant Leaf Positions on
213 Some Micronutrients, Anti-nutrients and Toxic Substances in *Telfairia occidentalis* at the
214 vegetative phase. *American Journal of Experimental Agriculture* 2(2): 219-232.
215
- 216 (13) Musa Amanabo 2012. Effect of plant leaf positions on the concentration of some
217 micronutrients, antinutrients and toxic substances in *Hibiscus Sabdarifa* at vegetative phase.
218 *Asian Journal of Plant Science and Research*. 2(3): 342-349. Available:
219 www.pelagiaresearchlibrary.com
- 220 (14) Makobo N.D., Shoko M.D. and Mtaita T.A. 2010. Nutrient content of vegetable Amaranth
221 (*Amaranthus cruentus* L.) at different harvesting stages. *World Journal of Agricultural Sciences*
222 6(3): 285-289.
223
- 224 (15) Oduntan A. O. Akinwande B. A. and Olaleye O. 2011. Effect of plant maturity on the
225 antioxidant properties, total phenolic and mineral contents of *Sesamum radiatum* leaves. *African*
226 *Journal of Food Science*. 5(17): 914-920.
227

- (16) Musa A., Oladiran J.A., Ezenwa M.I.S, Akanya H.O. and Ogbadoyi E.O. 2011. The effects of applied nitrogen fertilizer and leaf positions on levels of micronutrients, antinutrients and toxic substances in *Amaranthuscreuntus*. *African Journal of Biotechnology*. 10(48): 9857-9863.
- (17) SmartA.,Jeranyama P. and OwensV. 2004. The Use of turnips for extending the grazing season. Cooperative Extension Service. ExEx 2043. 166.
- (18) Oduntan A.O. and Olaleye O. 2012. Effect of plant maturity on the proximate composition of *Sesamumradiatum* Shum leaves. *Journal of Food Studies*.1 (1): 69-76. Available: www.macrothink.org/jfs
- (19) Fasuyi A.O., DairoF.A.S., and AdenijiA.O. 2008. Tropical vegetable (*Amaranthuscruentus*)leaf meal as alternativeprotein supplement in broiler starter diets: bionutritionalevaluation.*Journal of Central European Agriculture*. 9(1):23-34.
- (20) Sheela 2004. Proximate composition of underutilized green vegetables in southern Karnataka. *Journal of Human Ecology*. 15(3):229.