

Original Research Article

Some Nutritive and Antifungal Properties of *Citrus sinensis* (Sweet Orange) Peels and Seeds

ABSTRACT

Introduction: *Citrus sinensis* (sweet orange) fruit is widely consumed the world over for its sweet juice. The peels and seeds may have some nutrients and antifungal properties but are essentially discarded with attendant waste generation.

Aim: The study evaluated some nutritive and antifungal properties of the peels and seeds of *Linnaeus Osbeck* variety of *Citrus sinensis* (sweet orange) fruits purchased from Eke-Okigwe market in Imo state, Nigeria.

Study Design: The peels and seeds were respectively investigated for some minerals and vitamins content and for activity against some fungi.

Place and Duration of Study: The study was conducted at the Department of Biochemistry, Michael Okpara University of Agriculture Umudike, Nigeria between May and August, 2015.

Methodology: The minerals and vitamins content in the respective sample flour and the antifungal activity of the respective sample crude ethanol (95 %) and water extracts were determined by standard methods. Each extract (100 mg/ml) was tested against two fungi – *Candida albicans* and *Aspergillus flavus*.

Results: Results showed that the vitamins content of the orange peels and seeds respectively for vitamin A (IU) (85.71 ± 0.63 , 22.51 ± 18.04) was highest followed by ascorbic acid (mg/100g) (12.91 ± 1.02 , 7.04 ± 1.76). The content of the other determined vitamins in the samples was low (0.09 ± 0.00 to 0.81 ± 0.01). The minerals (mg/100g) in the peels were higher than that in the seeds. The calcium content in the peels (49.05 ± 26.24) was highest followed by magnesium (41.83 ± 5.59), sodium (19.44 ± 1.58) and potassium (14.90 ± 32.94). The ethanol extracts of the peels and seeds showed activity (mm) against the tested fungi (*Candida albicans* and *Aspergillus flavus*). However, the activity of the aqueous extract of the peels against *Candida albicans* and *Aspergillus flavus* respectively (12.33 ± 1.15 , 6.67 ± 0.58) was higher ($p < 0.05$) than that of the seeds (7.00 ± 1.00 , 0.00 ± 0.00). The observations, aside the difference in the riboflavin and thiamine content and the activity elicited by the ethanol extracts of the peels and seeds, were significant ($p < 0.05$).

Conclusion: The sweet orange peels could be a better source than the seeds for these nutrients, hence may offer higher nutrient benefits while ethanol may be preferred solvent to water for extracting the active principles in the samples. The higher nutrient mix in the peels probably accounted for the higher antifungal activity of the peels extracts against the tested fungi. The study provides basis for exploiting these sweet orange fruit wastes in diets and drugs, warranting further studies to harness the present findings.

Keywords: Minerals, vitamins, antifungal, *Candida albicans*, *Aspergillus flavus*

1. INTRODUCTION

The practice of consuming the juice of most fruits while discarding the peel and seed could contribute to increasing solid food wastes with potential adverse environmental and public health implications [1,2]. Orange pulp is an excellent source of vitamin C [3] and it is interesting to find out if the peels and seeds contained vitamin C and other nutrients to serve as alternative sources. Furthermore, the antifungal properties in orange peel and seed are of interest because drugs resistance is becoming a world-wide public health concern. Important bioactive compounds and activities could be inherent in these seeming food wastes. For instance, the seeds and peels of grapes and pomegranates have rich natural antioxidant [4] that could reduce oxidative stress in animals while the seed and peel of watermelon have nutritive and antimicrobial properties [5,6,1]. Possible roles of bioactive compounds in improving and managing even metabolic diseases have been suggested [7,8].

Sweet orange fruits contain vitamin C, fiber, as well as other bioactive components, including carotenoids and phenolic compounds [9]. The edible orange fruits juice have antioxidant property [10] attributable to the rich vitamin C, flavonoids and phenolic compounds content. Generally, antioxidants reduce oxidative stress which is a common feature in health dysfunctions in apparent support of the suggestion by Crowell [11] that these bioactive compounds, including the antioxidant compounds, in sweet orange juice could reduce the risk for cancers and many chronic diseases. *Citrus limonum* (lemon) contain esculetin, a bioactive compound that improved markers of health functions in rats [12]

while the sweet orange peel essences had antiseptic, analgesic and anti inflammatory values [13,14]. Human health challenges are seemingly ever growing and could result from adverse effects of foods and food condiments [15,16] warranting constant search for scientific basis to utilize plants/plant parts, fruits and fruit wastes.

Studies on orange fruits were reported [17], but not essentially on the mineral and vitamin compositions or the antifungal properties of the peels and seeds. The food value and pharma-food potentials of a food source could be further assessed through the mineral and vitamin compositions of the food source while the possible pharmacologic properties could be assessed through the antifungal activities, warranting the present study of the milled peels and seeds from *Linnaeus osbeck* variety of sweet orange (*Citrus sinensis*). The aim of this study was to determine some nutritive and antifungal properties of ~~aqueous and ethanolic~~ extracts of sweet orange (*Citrus sinensis*) peels and seeds extracts. The objectives of this work were to determine some minerals and vitamins in the sweet orange peels and seeds flour and the activity of the ethanol and aqueous extracts of the sweet orange peels and seeds against some fungi (*Candida albicans* and *Aspergillus flavus*).

2. MATERIALS AND METHODS

2.1 Chemicals and Reagents

The solvent, ethanol and other chemicals used, including those used in the preparation of reagents, were of analytical grade and products of reputable companies. This study was conducted between May and August, 2015 at the Department of Biochemistry, Michael Okpara University of Agriculture Umudike, Nigeria.

2.2. Collection, Identification and Preparation of Samples

The sweet orange fruits were purchased from a particular supplier at Eke-Okigwe market, a weekly market in Okigwe Imo state, Nigeria. The fruits were identified as *Linnaeus osbeck* variety of the *Citrus sinensis* (sweet orange) by a taxonomist in the Central Laboratory Unit of National Root Crop Research Institute Umudike, Nigeria. As described in a similar study [1], the orange fruits were thoroughly washed to remove sand particles and unwanted particles. The sweet orange fruits were thoroughly washed with clean water to remove dust particles and sliced longitudinally into four equal parts, using a home choice European knife. The juicy flesh or pulp containing the seeds was carefully removed from the peel. The seeds were carefully picked from the pulp and washed off the orange juice using clean water while the peel was chopped into bits. The samples were separately placed on a foil and weighed with a Satorious Digital Weighing Balance, Model BP210S, Germany before and after sun drying for seven days to obtain the respective wet weight (seeds = 86.05 g, peels = 169.74 g), dry weight (seeds = 46.89 g, peels = 153.12 g) and percentage yield (seeds = 54.49 %, peels = 90.21 %). The respective dry weight sample was separately blended into powder using Arthur Thomas Laboratory Mill Crypto model, USA, covered separately in a labeled white nylon and kept in the desiccator until used.

2.3. Samples Extraction

The aqueous and ethanol extracts respectively of the samples (peels and seeds) were separately obtained as described previously [18].

2.4. Nutritive (Minerals and Vitamins) Content Determination

Vitamins A, B₁ (thiamine), B₂, B₃ (niacin), and B₆ were variously determined by the spectrophotometric methods [19,20] whereas vitamin C (ascorbic acid) was determined by the method described by Okwu and Josiah [21]. Mineral content viz: potassium, sodium, calcium and magnesium were determined by the spectrophotometric method described by James [22], using Jenway Digital Spectrophotometer, Model 6320D, manufactured by Jenway Equipment Company, France. Potassium and sodium concentration were determined by the flame photometric method, using Jaway Digital Flame Photometer.

2.5. Tested Fungal Species

The fungal species used for the antifungal test, *Candida albicans* and *Aspergillus flavus*, were clinical isolates provided by the Central Laboratory of National Root Crop Research Institute Umudike Abia state, Nigeria.

2.6. Antifungal Activity Test

The disc agar diffusion method was used to determine the antifungal activity of the extracts as reported earlier [18]. Incubation was at 37°C for 24 hours under aerobic condition. The antifungal activity was determined by measuring the diameter (in millimetres, mm) of the zone of inhibition formed around the discs. The antifungal activity test was performed in triplicate and the mean zone of inhibition calculated.

2.7. Data Analysis

The data obtained by triplicate determinations were subjected to analysis of variance (ANOVA) using SPSS 16.0 for Windows. Comparison of difference in means was based on Students t-test. Difference in mean at a p value < 0.05 was regarded as statistically significant. Results were expressed as mean± standard deviation (SD).

3. RESULTS AND DISCUSSION

Results as shown in Table 1, showed that the vitamins content of the orange peels and seeds respectively for vitamin A (IU) (85.71±0.63, 22.51±18.04) was highest followed by ascorbic acid (mg/100g) (12.91±1.02, 7.04±1.76). The content of the other determined vitamins in the samples was low (0.09±0.00 to 0.81±0.01).

Generally, the minerals (mg/100g) in the peels were higher than that in the seeds. In particular, the calcium content in the peels (49.05±26.24) was highest followed by magnesium (41.83±5.59), sodium (19.44±1.58) and potassium (14.90±32.94) (Table.2).

As shown in Tables 3 and 4, the ethanol extracts of the peels and seeds showed activity (mm) against the tested fungi (*Candida albicans* and *Aspergillus flavus*). However, the activity of the aqueous extract of the peels against *Candida albicans* and *Aspergillus flavus* respectively (12.33±1.15, 6.67±0.58) was higher (p<0.05) than that of the seeds (7.00±1.00, 0.00±0.00).

The observations, aside the difference in the riboflavin and thiamine content and the activity elicited by the ethanol extracts of the peels and seeds, were significant (p<0.05).

Table 1: Some vitamin composition of *Citrus sinensis* (sweet orange) peels and seeds

Vitamins	Peels	Seeds	Difference
Retinol (vitamin A) IU	85.71±0.63 ^a	22.51±18.04 ^b	±63.20*
Ascorbic acid (vitamin C) (mg/100 g)	12.91±1.02 ^a	7.04±1.76 ^b	± 5.87*
Niacin (vitamin B ₃) (mg/100 g)	0.81±0.01 ^a	0.23±0.01 ^b	±0.58*
Riboflavin (vitamin B ₂) (mg/100 g)	0.15±0.20 ^a	0.06±0.01 ^a	±0.09 ^{ns}
Thiamine (vitamin B ₁) (mg/100 g)	0.11±0.00 ^a	0.09±0.00 ^a	±0.02 ^{ns}

Result = Value ± SD of duplicate determinations. Different superscript in a row or column means that the difference is significant (p<0.05), ns = difference is not significant (p > 0.05). * = difference is significant (p < 0.05)

Table 2: Some mineral composition of *Citrus sinensis* (sweet orange) peels and seeds

Minerals	Peels (mg/100 g)	Seed (mg/100 g)	Difference (mg/100 g)
Calcium	49.05±26.24 ^a	2.00±0.10 ^b	±47.05*
Magnesium	41.83±5.59 ^a	0.33±0.30 ^b	±41.50*
Sodium	19.44±1.58 ^a	0.33±0.30 ^b	±19.11*
Potassium	14.90±32.94 ^a	0.82±0.10 ^b	±14.08*

Result = Value ± SD of duplicate determinations. Different superscript in a row or column means that the difference is significant (p<0.05), ns = difference is not significant (p > 0.05). * = difference is significant (p < 0.05)

Table 3: Anti-fungal activity (inhibition zone diameter, IZD (mm)) at a concentration of 100 mg/ml of water extract of *Citrus sinensis* (sweet orange) peels and seeds

Fungal species	Water extract (100mg/ml)		Difference (mm)
	Sweet orange peels	Sweet orange seeds	
<i>C. albicans</i> (IZD, mm)	12.33±1.15 ^b	7.00±1.00 ^a	±5.33*
<i>A. flavus</i> (IZD, mm)	6.67±0.58 ^b	0.00±0.00 ^a	±6.67*

Result = Value ± SD of duplicate determinations. Different superscript in a row or column means that the difference is significant ($p < 0.05$), ns = difference is not significant ($p > 0.05$). * = difference is significant ($p < 0.05$)

Table 4: Anti-fungal activity (inhibition zone diameter, IZD (mm)) at a concentration of 100 mg/ml of ethanol extract of *Citrus sinensis* (sweet orange) peels and seeds

Fungal species	Ethanol extract (100mg/ml)		Difference (mm)
	Sweet orange peels	Sweet orange seeds	
<i>C. albicans</i> (IZD, mm)	14.67±1.15 ^c	14.00±1.00 ^c	±0.67 ^{ns}
<i>A. flavus</i> (IZD, mm)	11.67±1.15 ^d	10.00±1.00 ^d	±1.67 ^{ns}

Result = Value ± SD of duplicate determinations. Different superscript in a row or column means that the difference is significant ($p < 0.05$), ns = difference is not significant ($p > 0.05$). * = difference is significant ($p < 0.05$)

The sweet orange (*Citrus sinensis*) peels and seeds are usually discarded as food wastes. This could adversely affect the environment and public health. Thus, the present study investigated some nutritive and antifungal properties of the *Citrus sinensis* (sweet orange) peels and seeds. The result could provide scientific basis for enhanced possible beneficial use of these hitherto food wastes in animal diets and drugs ultimately reducing the accumulation of the wastes in, and the attendant waste burden on, the environment.

Compared to the other determined vitamins, vitamin A (IU) content in the peels and seeds was highest followed by vitamin C (mg/100g) while the content of the B vitamins in the samples was comparatively low (Table 1). Generally, vitamins play important roles in the regulation of normal metabolism and as an antioxidant [23] but may be required in small amounts. The vitamin C composition in the peels and in the seeds compared with the value reported for *Teculia africana* [24], but lower than the values reported for *Aspilia africana* and *Bryophyllum pinnatum* [21] and for *Citrus lemon* [25]. The result implies appreciable quantity of vitamin A (retinol) and vitamin C (ascorbic acid) in the sweet orange peels and seeds, though higher in the peels than in the seeds. The comparatively lower content of the B vitamins (thiamine, riboflavin and niacin) in the samples compared fairly with the values reported for *Aspilia africana* and *Bryophyllum pinnatum* [21]. Thus, the vitamins as in the samples could be adequate the vitamin functions warranting further studies to support the use of these samples as dietary vitamins supplement.

The minerals (mg/100g) in the peels were higher than that in the seeds while the calcium content in the peels was highest followed by magnesium, sodium and potassium (Table 2). This supports the presence of these minerals in food and food wastes [26]. However, the lower value of these minerals in the seeds than in the peels of the sweet orange contradicts the lower minerals content in the *Citrullus lanatus* (watermelon) rind/peels than in the seed [6], though not distinctive as suggested by the ash content of the watermelon rind and seed [27]. The sodium content in the sweet orange seeds did compare with the value for *Vitex mombassae* (96.08 ± 0.28) and *Maenua angolensis* (96.11 ± 0.76) as reported by Emmanuel *et al.* [24]. The composition of sodium, potassium, calcium and magnesium in the peels was higher when compared with the respective value for *Aspilia africana* and *Bryophyllum pinnatum* [21]. Manganese and magnesium contents value in some plant species reported by Emmanuel *et al.* [24] compared with the value in the present samples. Minerals in adequate amount ensure the normal physiological functions [28,29]. Hence the abundance of these minerals in the samples, especially in that of the peels, is nutritionally and physiologically noteworthy. Prior study by Nicolosi *et al.* [30] and Barros *et al.* [31] confirmed the presence of vitamins, minerals and other phytochemicals in the fruits of *Citrus sinensis* and in the fruit peels and seeds [31].

The ethanol extracts of the peels and seeds showed activity (mm) against the tested fungi (*Candida albicans* and *Aspergillus flavus*). However, the activity of the aqueous extract of the peels against

Candida albicans and *Aspergillus flavus* respectively was higher ($p<0.05$) than that of the seeds (Table 3). The result indicated that the peel and seed extracts showed antifungal activity against the tested fungi.

The ethanol extracts showed broader activity against the tested fungi, hence ethanol may be preferred solvent to water for extracting the active principles in the samples. The higher nutrient mix in the peels probably accounted for the higher antifungal activity of the peels extracts against the tested fungi. Egbunu et al. [2] reported the properties of watermelon seed oil whereas Ejikeme et al. [32] reviewed the melon seed oil potential for diesel fuel application. Thus, further studies in these directions are needed to improve the possible uses of the sweet orange seeds.

4. CONCLUSION

Thus, the sweet orange peels could be a better source than the seeds for these nutrients, hence may offer higher nutrient benefits while ethanol may be preferred solvent to water for extracting the active principles in the samples. The higher nutrient mix in the peels probably accounted for the higher antifungal activity of the peels extracts against the tested fungi. The study provides basis for exploiting these sweet orange fruit wastes in diets and drugs, warranting further studies to harness the present findings in that direction.

CONSENT (WHEREEVER APPLICABLE)

Not applicable

ETHICAL APPROVAL (WHEREEVER APPLICABLE)

Not applicable

REFERENCES

- Egbunu ACC, Osuji CA. Proximate compositions and antibacterial activity of *Citrus sinensis* (sweet orange) peel and seed extracts. *European Journal of Medicinal Plants*, 2016; 12(3): 1-7. DOI:10.9734/EJMP/2016/24122.
- Egbunu ACC, Aguguesi RG, Samuel R, Ojunkwu O, Onyenmeri F, Uzuegbu U. Some physicochemical properties of the petroleum ether-extracted watermelon (*Citrullus lanatus*) seed oil. *Asian Journal of Scientific Research*, 2015; 8(4): 519-525. DOI: 10.3923/ajsr.2015.519-525.
- ~~Aschoff JK, Kaufmann S, Kalkan O, Neidhart S, Carlo R, Schweiggert RM. "In Vitro Bioaccessibility of Carotenoids, Flavonoids and Vitamin C from differently processed Oranges and orange Juices (Citrus Sinensis (L) Osbeck)" *Journal Agric Food Chem.*, 2015; 63(2): 578-587.~~
- Jayaprakasha GK, Singh RP, Sakariah KK. Antioxidant activity of grape seed *Vitis vinifera* extracts on peroxidation models in-vitro: *Fd. Chem.*, 2001, 73: 285-290.
- Egbunu ACC. Assessment of some antinutrient properties of watermelon (*Citrullus lanatus*) rind and seed. *Res. J. Environ. Sci.*, 2015a; 9(5): 225-232.
- Egbunu ACC. Comparative assessment of some mineral, amino acid and vitamin compositions of watermelon (*Citrullus lanatus*) rind and seed. *Asian Journal of Biochemistry*, 2015b; 10(5): 230-236. DOI: 10.3923/ajb.2015.230.236
- Ezeanyika LUS, Egbunu ACC. Impact of nitric oxide and insulin resistance on the pathophysiology of the metabolic syndrome: Possible role of L-arginine and glutamate. *J. Med. and Med. Sci.*, 2011; 2(20): 657-662.
- Egbunu ACC, Ezeanyika LUS. L-arginine exposure improves renal function markers of metabolic syndrome in female rats. *Am. J. Biochem. and Mol. Biol.*, 2013; 3(1) 50-60.
- Atolani O, Omere J, Otuechere CA, Adewuyi A. Antioxidant and cytotoxicity effects of seed oils from edible fruits. *Journal of Acute Disease*, 2012; 130-134 (doi: 10.1016/S22216189(13)60073-6).
- Tripoli E, La Guardia M, Giammanco S, Di Majo D, Giammanco M. Citrus flavonoids: Molecular structure, biological activity and nutritional properties: A review. *Food Chemistry*, 2007; 104: 466-479.
- Crowell PL. Prevention and therapy of cancer by dietary monoterpenes. *The Journal of Nutrition*, 2009; 129 (3): 775S-778S, PMID 10082788.
- Egbunu ACC, Ogbu AE, Ezeanyika LUS. Dose-related influence of esculetin (6,7-dihydroxycoumarin) on some liver and prostate function markers of male Wistar rats. *Journal of Biological Sciences*, 2012; 12(4): 253-257.

- 273 13. Muhammad G, Sajid M, Nasir M, Nyla J, Rehana K, Kokab J, Gulshan A. Composition and
274 antimicrobial properties of essential oil of foeniculum vulgare. *African Journal of*
275 *Biotechnology*, 2008; 7(24): 4364-4368.
- 276 14. Akin M, Aktumsek A, Nostro A. Antibacterial activity and composition of the essential oils of
277 eucalyptus camaldulensis dehn and myrtus communis L. growing in northern cyprus. *African*
278 *Journal of Biotechnology*, 2010; 9(4): 531-535.
- 279 15. Egbuonu ACC, Osakwe ON. Effects of high monosodium glutamate on some serum markers
280 of lipid status in male Wistar rats. *Journal of Medicine and Medical Sciences*, 2011; 2(1):653-
281 656.
- 282 16. Egbuonu ACC, Obidoa O, Ezeokonkwo CA, Ezaeanyika LUS, Ejikeme PM. Low dose oral
283 administration of monosodium glutamate in male albino rats may be nephroprotective. *Bio-*
284 *Research*, 2009; 7(1): 470-473.
- 285 17. Omodamiro OD Umekwe JC. Evaluation of anti-inflammatory, antibacterial and antioxidant
286 properties of ethanolic extracts of *Citrus sinensis* peel and leaves. *Journal of Chemical and*
287 *Pharmaceutical Research*, 2013; 5(5):56-66.
- 288 18. Egbuonu ACC. Comparative investigation of the antibacterial and antifungal potentials of the
289 extracts of watermelon (*Citrullus lanatus*) rind and seed. *European Journal of Medicinal*
290 *Plants*, 2015c; 9(4): 1-7. DOI:10.9734/EJMP/2015.
- 291 19. Onwuka GI. Food analysis and instrumentation: Theory and Practice. 1st edn., 2005; Naphtali
292 Prints, Surulere, Lagos-Nigeria.
- 293 20. Okwu DE. Spectrophotometric method for vitamin analysis: Phytochemicals, vitamins and
294 minerals contents of two Nigerian medicinal plants. *International J. Molecular Medicine and*
295 *Advance Sciences*, 2004; 1: 375-381.
- 296 21. Okwu DE, Josiah C. Evaluation of the chemical composition of two Nigerian medicinal plants.
297 *Afr. J. of Biotechnol.*, 2006; 5(4): 357-361.
- 298 22. James CS. Analytical Chemistry of Foods. Chapman and Hall, New York, p. 178.
- 299 23. Barminas JT, Charles M, Emmanuel D. Mineral composition of non-conventional leafy
300 vegetables. *Plant Fds. Hum. Nutr.*, 1998; 53: 29-36.
- 301 24. Emmanuel VE, Njoka JT, Catherine LW, Lyaruu HVM. Nutritive and anti-nutritive qualities of
302 mostly preferred edible woody plants in selected drylands of Iringa district, Tanzania. *Pak. J.*
303 *of Nutr.*, 2011; 10 (8): 786-791.
- 304 25. Nangbes JG, Lawam DT, Nvau JB, Zukdimma NA, Dawam NN. Titrimetric determination of
305 ascorbic acid levels in some citrus fruits of Kurgwi, Plateau State Nigeria. *IOSR J. of Appl.*
306 *Chem. (IOSR-JAC)*. 2014; 7(9): 01-03.
- 307 26. Hafiza MA, Parveen B, Ahmad F, Hamid K. Phytochemical and antifungal screening of
308 *Medicago sativa* and *Zinnia elegans*. *Online J. Biol. Sci.*, 2002; 2 (2): 130-132.
- 309 27. Egbuonu ACC. Comparative investigation of the proximate and functional properties of
310 watermelon (*Citrullus lanatus*) rind and seed. *Res. J. of Environ. Toxicol.*, 2015d; 9(3): 160-
311 167.
- 312 28. Adeyeye EI. Bio-concentration of macro and trace mineral in prawns living in Lagos Lagoon.
313 *Pak. J. Sci. Ind. Res.*, 2000; 43: 367-373.
- 314 29. Umeta M, West CE, Habtamu F. Content of zinc, iron, calcium and their absorption inhibitors
315 in foods commonly consumed in Ethiopia. *J. of Fd. Comp. and Anal.*, 2005; 18(8): 803-817.
- ~~316 30. Nicolosi E, Deng ZN, Gentile A, La Malfa S, Continella G, Tribulato E. "Citrus phylogeny and~~
~~317 genetic origin of important species as investigated by molecular markers" TAG Theoretical~~
~~318 and Applied Genetics, 2000; 100(8):1155-1166~~
- 319 31. Barros HR, Ferreira TA, Genovese MI. "Antioxidant Capacity and Mineral Content of pulp and
320 peel from Commercial Cltivars of Citrus from Brazil". *Food Chem.*, 2012; 134(4):1892-1898
- 321 32. Ejikeme PM, Egbuonu ACC, Anyaogu ID, Eze VC. Fatty acid methyl esters of melon seed oil:
322 Characterization for Potential Diesel Fuel Application. *Leonardo Journal of Sciences*, 2011;
323 18: 75-84.