

Modeling of advanced photo oxidation of Alizarin Red-S dye using TiO₂ as photo catalyst

Tariq H. Talib¹, Murad A. AlDamen², Ramzie R. Alani^{1 *}.

¹Department of chemistry, Al-Mustansiriya University, Baghdad, Iraq

²Department of Chemistry, University of Jordan, Amman 11942, Jordan

Abstract: The objective of this study was to investigate the experimental study and theoretical study of advanced photo oxidation using TiO₂ as photo catalyst and to compare between these two studies. This work contains two parts. The first part deals with removing efficiency of an industrially important dye Alizarin Red-S (ARS) from aqueous media using TiO₂ anatase as photo catalyst. Various parameters, which can influence the degradation rate like, initial concentration, temperature, pH, photo catalyst dosage were investigated.

The second part deals with the theoretical study of the TiO₂ surface and the pollutant substance (ARS) using the program (Spartan '08 V1.2.0). we used two kinds of calculations, the first was the Equilibrium Geometry optimization at ground state by Density Functional Theory/B3LYP method and Semi-empirical/PM3 method to predict the most stable geometrical formula for TiO₂ anatase (001),(101) and (ARS) pollutant in different medium (neutral, acidic, basic). The program showed that the method DFT/B3LYP give the best results compared with DFT/PM3 and that the basic medium was better than acidic and neutral medium which means that the highest rate of (ARS) degradation was in basic medium. The second calculation was the energy geometry at ground state by density functional theory/EDF2 method to calculate standard thermodynamic quantities for (ARS) pollutant and TiO₂ anatase (001),(101) surface that showed the highest stability for them in the basic medium.

The results from theoretical study agreed with the results obtained in the experimental study. Results from the experimental study showed Thermodynamic properties was calculated for the photodegradation of (ARS) as (ΔH , ΔG , ΔS) which is showed an exothermic heterogeneous reaction with pseudo first order reaction and optimum parameters for removal of this dye from aqueous medium were (0.16 g/l, 20ppm, temp.30°C, pH9). Theoretical study DFT/B3LYP method showed better results than DFT/PM3 to predict the most stable geometrical formula, and TiO₂ anatase (001) surface was more active than TiO₂ anatase (101) surface.

Keywords: [Molecular modeling, Alizarin Red-S Degradation, TiO₂ anatase (001)(101) surface, DFT Methods, Equilibrium Geometry optimization, energy geometry]

1. Introduction

Textile dyeing process is a significant source of contamination responsible for the environmental pollution. To meet the demand of textile industry, a large variety of dyes are annually produced in tremendous quantities (7×10^5 Tones) in the world. Thus the volume of wastewater containing processed textile dyes is on steady increase⁽¹⁾.

Most of the dyes used in the textiles industries are stable to light and non biodegradable⁽²⁾. In order to reduce the risk of environmental pollution from such wastewater, it is necessary to treat them before discharging into the environment⁽³⁾. Today more than 10,000 dyes have been incorporated in colour index⁽⁴⁾, as being the derivatives of aromatic compounds, with

* Tel.: +9647902986137.

E-mail address: ramziealani@yahoo.com.

halo, nitro, and sulpho grouping in their structures most of these are toxic and carcinogenic in nature and pose serious environment threats to human beings as well as aquatic life when present in industrial effluents. Many physical, chemical, and biological methods are employed to eliminate dyes from the wastewater. Physical methods are preferred over chemical and biological methods, because firstly they are simple and inexpensive and secondly do not produce byproducts in water bodies^(5,6), Alizarin red-S is an anionic dye which are widely used in woven fabrics, wool, cotton textiles^(7,8).




Figure .1 : Alizarin Red-S

Alizarin Red-S (1,2-dihydroxy-9,10-anthra-quinone sulfonic acid sodium salt, Fig. 1) is a water-soluble, widely used anthraquinone dye. It is synthesized by sulfonation of alizarin which is a neutral dye obtained from madder (*Rubia tinctorum* L. Rubiaceae). Anthraquinone dyes like ARS are durable pollutants, released especially by textile industries in the aquatic ecosystems.

Removal of these dyes from industrial wastewaters is a crucial process, from both economical and environmental points of view⁽⁹⁾.

In this study TiO_2 is used as a photocatalyst. K.M.Joshi et al⁽¹⁰⁾ indicated that TiO_2 is a good photocatalyst for the removal of Alizarin Red-S from aqueous medium at pH 8. The aqueous phase photochemical degradation of dyes like Alizarin red-S has been studied in the presence of semiconducting oxide like TiO_2 and ZnO ^(11,12). In recent years, photocatalytic oxidation technology has been found capable of effectively removing pollutants⁽¹³⁾, Anatase TiO_2 is in the focus of photocatalytic research because of its strong oxidizing property, high catalytic ability, etc. The anatase TiO_2 surface has an important impact on photocatalytic activity, photovoltaic properties, and gas sensor characteristics. Hence, an intensive study of the anatase TiO_2 surface is very significant. The most stable surface of anatase TiO_2 is the (101), followed (001) surfaces^(14,15).

The experimental part of our study aims to evaluate the photodegradation of Alizarin Red-S (ARS) as pollutant substance using TiO_2 anatase as an absorbance surface and photocatalyst with (UV) Light. The effect of initial concentration of (ARS), the PH, and temperature was studied on the rate of photodecomposition of (ARS).

The theoretical part included two calculations, the first was the Equilibrium Geometry optimization at ground state using Density Functional Theory/B3LYP method and Semi-empirical/PM3 method to predict the most stable geometrical formula, the second calculation of the energy at ground state by using Density Functional Theory/EDF2 method to calculate Standard Thermodynamic quantities for pollutant and TiO_2 surface, and to compare between these two studies.

Quantum chemical calculations employing DFT⁽¹⁶⁾ are conducted to investigate the energetic of TiO_2 (101) and (001) anatase surfaces and ARS surface in different mediums (basic, natural, acidic). (101) face represents the most probable anatase surface⁽¹⁷⁾ and (001) surface is considered to be the most stable surface of TiO_2 anatase⁽¹⁸⁾. The anatase TiO_2 cluster contains two Ti atoms and nine O atoms. This cluster approach is a well known and successful approach applied in quantum chemical calculations^(19,20). However, since there may be limitations to the use of small cluster models to represent oxide surfaces where the peripheral oxygen atoms are saturated with hydrogen atoms, the effect of cluster size is

also tested by use of semi empirical PM3 calculations. The basis set employed in the DFT calculations was 6-31G** provided in SPARTAN' 08 (Wave function Inc.). SPARTAN'08 was used for all the semi empirical and DFT calculations reported, Neutral cluster models of (101) and (001) titanium dioxide anatase surfaces represented by $Ti_2O_9H_{10}$ but basic cluster models was represented by $Ti_2O_9H_9$ and acidic cluster models represented by $Ti_2O_9H_{12}$ for (101) surface whereas (001) surface acidic cluster models was represented by $Ti_2O_9H_{11}$ ⁽²¹⁾ These clusters model geometries are then energetically optimized. A. Cordoba⁽²²⁾ investigate the possibility of use of enzymatic treatment by experimental and theoretical study involve molecular modeling through MM2–PM6 by used MOPAC.2009, Vittadini et al⁽²³⁾ indicate that the adsorption of formic acid and sodium format on the dry and hydrated TiO_2 anatase (101) surfaces a molecular monodentate species hydrogen bonded to a surface two-fold coordinated oxygen is preferred on the dry surface. But on the hydrated surface, HCOOH and HCOONa preferentially form an inner-sphere adsorption complex. Opera et al⁽²⁴⁾. published that The Ti–O bond lengths for NKX-2398 dye are shorter from (NKX-2311 and C343) dyes, suggesting a strong binding, . Selloni et al⁽²⁵⁾ indicated by Molecular Dynamics (MD) calculations adsorption of water be more favorable on the surface TiO_2 (101) Anatase these results are agree with the findings of the RedFevn⁽²⁶⁾ But using methods Ab-initio Molecular Orbital Theory Density Functional Theory, Gong et al⁽²⁷⁾. explained TiO_2 (001) anatase surface which contains a lot of Ti5c atoms that give high efficiency on the adsorption molecules such and formic acid, that is agree with M. Vittoria⁽²⁸⁾ which indicate the high activity for TiO_2 (001) anatase surface from Ti5c atoms and large value of (Ti-O-Ti) bond.

2. Experimental details

Chemicals

All chemicals used during experimental work were of analytical grade and were used as such without purification. Alizarin Red-S (Riedel de Haën) and HCl (EDH 37%). NaOH (Fluka 99%) were used. Double distilled water was used for the preparation of all types of solution and dilution when required.

Instrumentation

Balance METTLER AE240 and Uv-vis spectrophotometer (PG Instruments Limited U.V- T80) were used in this study.

Standard Solutions

The stock solution of the dye was prepared by dissolving 0.1g/l of dye in water and made up a stock solution in volumetric flask. The concentration of the dye solution was determined spectrophotometrically. Standard solutions of dye were prepared by successive dilution of stock solution.

Experimental Procedure

The photo catalytic degradation of Alizarin red-S for both initial concentration and irradiation sample was determined by Uv-Visible Spectrophotometer (PG Instruments Limited U.V- T80). The calibration curve was obtained at $\lambda_{max} = 260 \text{ nm}$. The reaction mixture was irradiated with light source Uv lamp (PHILIPS-125Watt) at a distance of 15cm from the reaction vessel. Double distilled water was used throughout the experiment. Kinetics of adsorption was determined by analyzing adsorptive uptake of dye from aqueous solution at different time intervals for 15 minutes. In each experiment an accurately weighted amount of TiO_2 was taken in flask in 25 ml dye solution by adjusting pH of the Alizarin red-S dye solution.

* Tel.: +9647902986137 .

E-mail address: ramziealani@yahoo.com.

3. Results and Discussion

In order to find the appropriate conditions of photo catalyst dose , concentration of dye , pH , and temperature for the adsorption of Alizarin Red-S on TiO_2 anatase , various experiments were conducted .The results of these experiments were as follow.

Effect of Photo catalyst dose

The effect of variation in the photo catalyst amount on the process adsorption of Alizarin Red-S was studied, with different photo catalyst amount in the range of (0.1, 0.3, 0.4) g . The results are shown in Fig. 1.




Fig.2 Effect of photo catalyst dose on ARS degradation rate. experimental parameters: contact time, t (15 min), initial concentration of dye (20 ppm), temp. (30 $^{\circ}\text{C}$) and solution volume 25 ml .

From Fig.2, an increase in photo catalyst dose leads to increase in the percentage of the removed of alizarin. This is caused by an increased in number of available active points with increase in photo catalyst dose which mean increase of ($\text{h}^+ \text{e}^-$) pair number which caused increase in degradation rate⁽¹²⁾. On the other side when photo catalyst dose increase the absorbate photons number well increase that caused increase in degradation rate⁽²⁹⁾.

Effect of Initial Dye Concentration

Initial dye concentration was one of the effective factors on the rate of degradation reaction. The rate of Alizarin Red-S degradation was studied as a function initial dye concentration of in the range of 10-30 ppm. The results obtained are present in Fig.2. The rate of (ARS) degradation increases with increasing the initial concentration of the dye TiO_2 . It was observed that adsorption yield increased with increase in initial concentration of the dye. Minimum rate was 1×10^{-7} mol/L.min for 10ppm concentration to maximum rate value 1.3×10^{-7} mol/L.min for 30 ppm concentration of dye solution. This may be due to available active sites and increase in the adsorbed molecules which mean molecules involved in the interaction , as an increase in the high initial concentration of the dye.⁽¹⁰⁾




Fig. 3 : Effect of initial concentration of dye on (ARS) degradation rate. experimental parameters: t (15 min) ,temperature (30 $^{\circ}\text{C}$) and solu tion volume (V) 25 ml.

Effect of pH

The pH of the aqueous solution was clearly an important parameter that controls the degradation process. The rate of Alizarin Red-S degradation was studied as a function of pH in three mediums. The results obtained were shown in Fig. 3. The minimum degradation rate was 0.3×10^{-7} at pH 5.0 and maximum degradation rate was 0.9×10^{-7} at pH 9 was followed the consequences basic > neutral > acidic .

Because ARS as pollutants, which are hydrolyzed under alkaline conditions may show an increase of reaction rate with increase of pH⁽³⁰⁾. Finally, the presence of ionic species could affect the degradation process via adsorption of the pollutants, absorption of UV light and reaction with hydroxyl radicals that is leads to increase in degradation rate^(31,32).




Fig. 4 : Effect of pH on (ARS) degradation rate. experimental parameters: t(15 min), initial concentration of dye (20ppm), temperature (30 °C) and solution volume (V) 25 ml.

Effect of Temperature

Temperature has an important effect on the rate of adsorption. The rate of Alizarin Red-S degradation was studied as a function of temperature in the range of 25-40 °C. The results obtained were present in Fig.3. It was observed that the ARS degradation rate decreases with increase in temperature. The minimum degradation rate was at 40 °C and maximum degradation was at 30°C for 20 ppm initial concentration of dye solution. The decrease in adsorption was obtained at high temperature because molecules move with great speed and less time of interaction was available for dye anions with photo catalyst material^(33,34).




Fig. 5: Effect of temperature on (ARS) degradation rate of Alizarin Red-S. experimental parameters: t (15 min), initial concentration of dye (20ppm), and solution volume (V) 25 ml .

Thermodynamic parameters

Thermodynamic parameters such as standard Gibbs free energy (ΔG), Enthalpy (ΔH) and entropy (ΔS) were also calculated using equations 1,2 and 3 and the results obtained are illustrated in table-1 .

$$\Delta G = -RT \ln K \quad \text{-----(1)}$$

$$\frac{\Delta H^*}{R} \left[\frac{T_2 - T_1}{T_2 T_1} \right] = \ln \frac{K_2}{K_1} \quad \text{-----(2)} \quad 207$$

$$\Delta G = \Delta H - T \Delta S \quad \text{-----(3)}$$

T is the absolute temperature in Kelvin .

210

Table:1

Adsorbent	Temperature K	ΔG kJ.mol ⁻¹	ΔH kJ.mol ⁻¹	ΔS kJ.mol ⁻¹ . K ⁻¹
TiO ₂	303	-12.250	-20.9	0.028

211

The negative value of the ΔG at the studied temperature range indicated that the sorption of alizarin Red S on sorbent was thermodynamically feasible and spontaneous. The negative value of ΔH showed that the sorption was exothermic. The positive value of ΔS showed an increased randomness at the solid alizarin Red S solution interface during the adsorption of alizarin Red S.⁽¹²⁾

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489


490

491

234 **Fig. 6 : (001)surface Optimized geometrical formula in neutral medium .**
 235

236 **2. basic medium**

238




Heat of Formation: -798.443293 au

241 **Fig. 7 : (001)surface Optimized geometrical formula in basic medium .**

242 **3. Acidic medium**

244



Heat of Formation: -800.013728 au

246

247 **Fig. 8 : (001)surface Optimized geometrical formula in acidic medium .**

248

249

Table.2 bonds length and angle value for (001) surface.

	Medium	Method	Ti _{5C} -O _{2C} (a°)	θ
Calculated	Basic	B3LYP	1.831	131.73°
		PM3	1.949	80.70°
Calculated	Natural	B3LYP	1.738	129.73°
		PM3	1.950	81.22°
Calculated	Acidic	B3LYP	2.138	128.28°
		PM3	2.010	76.78
Ref.35		B3LYP	1.857	-
Ref.37		PBE	-	146°

250 **θ** : value of (Ti-O-Ti) bond

Ti_{5C}: fivefold Titanium atom

O_{2C}: twofold oxygen atom

253

* Tel.: +9647902986137 .

E-mail address: ramziealani@yahoo.com.

From table (2) value of (θ) angle showed that the (001) surface has most value from others that is mean highest reactivity in basic medium and showed that the method DFT/B3LYP give the best results and confirmed with The practical methods that the basic medium give the highest rate of decomposition of (ARS) followed the consequences:

basic > neutral > acidic

whereas ($Ti_{5c} - O_{2c}$) bond lengths in acidic medium has most value from others that is mean non favorite medium for TiO_2 anatase (001) surface.

TiO_2 anatase (101) Surface Geometrical Optimization

1. neutral surface




Fig. 9 : (101) surface Optimized geometrical formula in neutral medium .

2. basic medium




Fig. 10 : (101) surface Optimized geometrical formula in basic medium .

3. acidic medium




Fig. 11 : (101) surface Optimized geometrical formula in acidic medium.

Table.3 bonds length and angle value for (101) surface

	Medium	Method	Ti _{5C} -O _{2C}	Ti _{6C} -O _{2C}	θ
calculated	Basic	B3LYP	1.953	1.831	99.13
		PM3	1.896	1.981	83.09
calculated	Natural	B3LYP	1.873	1.823	96.71
		PM3	1.917	1.938	80.36
calculated	Acidic	B3LYP	2.004	1.979	109.38°
		PM3	2.014	2.027	75.70°
Ref.35		B3LYP	1.966	1.937	-
Ref.36		B3LYP	1.830	1.850	-
Ref.38		GGA	1.836	1.849	-
Ref.37		PBE	1.942	-	102

θ : value of (Ti-O-Ti) bond

Ti_{5C}: fivefold Titanium atom

Ti_{6C}: sixfold Titanium atom

O_{2C}: twofold oxygen atom

Both methods PM3 and B3LYP showed that the θ value for TiO₂ anatase (101) surface in the basic medium is more than that's in neutral medium which refers to high reactivity for (101) surface in basic medium but in acidic medium values varied where (B3LYP) method give large value of θ which refers to high reactivity from the basic and neutral mediums that doesn't agree with the experimental results while PM3 method showed small value of θ which refers to low reactivity less than in basic and neutral mediums which agree with the experimental results.

Comparing between TiO₂ anatase (001) and (101) surfaces by bonds length and angle value, we find that the (001) surface was more reactive than (101) surface. Our results agree with the previous findings reported in different studies^(39,40,27).

Table.4 : heat of formation for (001),(101) surface in different medium.

	Method	Basic	Neutral	Acidic	Energy Unit
001	PM3	-5452.21	-5535.16	-5089.44	kJ/mol au
	B3LYP	-798.443	-799.639	-800.013	
101	PM3	4798.750-	-5441.139	-4066.36	kJ/mol au
	B3LYP	-797.6	-799.646	-800.256	

* Tel.: +9647902986137 .

E-mail address: ramziealani@yahoo.com.

PM3 methods in table (4) showed that both TiO₂ anatase (001) and (101) surfaces has less heat of formation in neutral medium which means each of them had highest stability and lowest reactivity in neutral medium, whereas in acidic medium both surfaces have high heat of formation which means lowest stability and highest reactivity in acidic medium.

B3LYP methods showed that both TiO₂ anatase (001) and (101) surfaces have high heat of formation in basic medium which means each of them had lowest stability and highest reactivity in basic medium, whereas in acidic medium both surface have lowest heat of formation which means highest stability and lowest reactivity ,that's agree with the experimental results which indicate ARS degradation rate is the most in basic medium and this was very clear in 101 surface where the highest heat of formation was observed.

On the other hand comparing between calculated heat of formation by B3LYP method for both TiO₂ surfaces showed that 101 surface has highest heat of formation which means low stability and high reactivity than 001 surface. This result agree with previous studies^(25,26)

Geometrical Optimization) Alizarin Red-S (ARS)

1. Neutral medium




Fig. 12 :ARS pollutant Optimized structure in neutral medium by(PM3)(B3LYP) .

2. Basic medium




Fig. 13 :ARS pollutant Optimized structure in basic medium by(PM3)(B3LYP).

* Tel.: +9647902986137 .

E-mail address: ramziealani@yahoo.com.

3. Acidic medium




Fig. 14 :ARS pollutant Optimized structure in acidic medium by(PM3)(B3LYP).

Table.5 : heat of formation for ARS pollutant in different medium .

	Method	Basic	Natural	Acidic	Energy Unit
ARS	PM3	996.948-	-1081.805	-902.387	<i>kJ/mol</i>
	B3LYP	-1461.857	-1462.497	-1462.995	<i>au</i>

Heat of formation values in table (1-4) was calculated by B3LYP showed that's ARS pollutant has highest heat of formation in basic medium followed the consequences basic > neutral > acidic whereas PM3 method showed that's highest heat of formation in acidic medium followed the consequences acidic > basic > neutral that's indicate B3LYP methods give good results which is agree with the experimental results .

Theoretical Calculations for Surface and Pollutant

The thermodynamic calculation was performed depending on the energy calculations at ground state by using Density Functional Theory/EDF2 method to calculate Standard Thermodynamic quantities for pollutant and TiO₂ surface.

Tabel. 6: Standard Thermodynamic quantities for surface at 298.15 K and 1 atm.

surface	Medium	ZPE kJ/mol	Enthalpy kJ/mol	Entropy J/mol.K	Cv J/mol.K	Gibbs free energy G (kJ/mol)
001	Basic	254.6410	344.2600	528.5494	254.6410	186.916
	Natural	361.9670	409.9561	537.8440	266.3754	249.924
	Acidic	388.7331	434.5602	522.5073	257.6182	279.004
101	Basic	268.0738	313.1480	517.0250	250.3672	159.082
	Natural	361.6021	414.0866	574.2387	284.7062	243.028
	Acidic	412.2940	470.5403	606.5702	317.5846	289.952

C_v : heat capacity

ZPE: Zero point energy

* Tel.: +9647902986137 .

E-mail address: ramziealani@yahoo.com.

363 **Tabel. 7: Standard Thermodynamic quantities for ARS pollutant at 298.15 K and 1 atm.**

	system	ZPE kJ/mol	Enthalpy kJ/mol	Entropy J/mol.K	Cv J/mol.K	Gibbs free energy G kJ/mol
ARS/EDF2	Basic	467.4259	511.7977	527.8044	268.1554	354.4328
ARS/EDF2	Natural	502.9677	549.7038	546.4649	279.8043	386.7753
ARS/EDF2	Acidic	532.9485	577.9819	532.0865	274.4549	419.3403

364
365 The theoretical thermodynamic values for both (101),(001) surfaces and ARS pollutant in
366 different mediums in tables(1-4),(1-5) the increase of this values Means an increase of
367 stability and decrease of reactivity followed the consequences :

368 Increase reactivity ← basic > neutral > acidic → Increase stability

369 This results agreed with the experimental results that showed the highest reactivity and
370 lowest stability for each pollutant (ARS) and TiO₂ surface in the basic medium .

371

372 **Conclusion**

373 Results from the experimental study showed that the photodegradation reaction from
374 pseudo first order and exothermic reaction and optimum parameters for removal of this
375 dye from aqueous medium were (0.16 g/l, 20ppm, temp. 30 °C, pH9).

376 Theoretical study DFT/B3LYP method showed better results than DFT/PM3 to predict
377 the most stable geometrical formula, and TiO₂ anatase (001) surface was more active
378 than TiO₂ anatase (101) surface.

379

380 **4. ACKNOWLEDGEMENTS**

381 Authors would like to thank Al-Mustansiriya University for providing the financial support for
382 this work.

383

384

385

386

385 **References**

- 387 1. M. J. Iqbal, and M. N. Ashiq, *Journal of Hazardous Materials*, **139** (1), 57 (2007).
- 388 2. R. Dave et al.,(2010), "Photochemical and photocatalytic of cypermethrin under UV radiation"
389 *Der pharma chemica*, 2(1) pp 152-158.
- 390 3. Muhammad et al.(2007), "Thermodynamic adsorption of dyes from aqueous media on
391 activated charcoal", *Journal of Research SciencePakistan*, 18 (2) pp 91-99.
- 392 4. D. Jalajaa. et al.(2007), "Removal of acid dye from textile wastewater by adsorption using
393 Activated carbon prepared from pomegranaterind", *Pollution Research.*, 28(2), pp 287-290.
- 394 5. T. M. Elmorsi, Y. M. Riyad Zeinhom, H.Mohamed, and Hassan M. H. A. El-Bary,*Journal of*
395 *Hazardous Materials*, **174** (1-3), 352(2010).
- 396 6. S. Yahya. Al-Degs, I. Musa. El-Barghouthi, H.Amjad, El-Sheikh, and M. Gavin Walker, *Dyes,*
397 *and Pigments*, **77**, 16 (2008).
- 398 7. S. Parra, , Sarria V., Maloto S., Periner P., and pulgarin C.,(2004) "Photocatalytic Degradation
399 of Attrazine using suspended and supported TiO₂. *Journal of AppliedCatalyssis.B:*
400 *Environmental .*, 51, pp 107-116.
- 401 8. N. Kannan and Nagnathan G.G.(2007), "Photocatalytic degradation of Saferaninedye on
402 semiconductor by UV radiation" *International Journal of Environmental Protection*, 27 , pp
403 1103- 1108.
- 404 9. P.Zucca, C.Vinci, F.Sollai, A.Rescignoand E. Sanjust, *Journal of Molecular Catalysis*
405 *A:Chemical*, 288, 97(2008).
- 406 10. K.M. Joshi Degradation of Alizarine Red-S (A Textiles Dye) by Photocatalysis using ZnO and
407 TiO₂ as Photocatalyst. *International Journal of Environmental Sciences Volume 2 No.1,*
408 *2011,p10.*
- 409 11. A. C. Cakir, P. Guloglu, M. Bilgin, S. Erten-Ela, *Comparison of Randomly-oriented ZnO*
410 *Nanorods and Butterfly-type ZnO Nanorod Bundles for Toluidine Blue Photocatalytic*

- Degradation, *JOURNAL of OPTOELECTRONICS and ADVANCED MATERIALS*, Vol. 14, Issue 3-4, pp.230-237, 2012
12. J.solomon . Removal of Alizarin Red S (Dye) from Aqueous Media by using *Cynodon dactylon* as an Adsorbent, *J. Chem. Pharm. Res.*, 2011, 3(5):478-490
 13. Q. Tao, Wang J. Application of TiO₂ photocatalytic oxidation technology to gas-phase contaminants degradation. *Safety and Environmental Engineering*, 2010, 17(5): 26
 14. W. Hebenstreit, Ruzycski N, Herman G S, et al. Scanning tunneling microscopy investigation of the TiO₂ anatase (101) surfaces. *Phys Rev B*, 2000, 62: 16334
 15. R. Hengerer, Bolliger B, Erbudak M, et al. Structure and stability of the anatase TiO₂ (101) and (001) surfaces. *Surf Sci*, 2000, 460: 162.
 16. W. Kohn, L.J. Sham, *Phys. Rev.* 140 (1965) A1133.
 17. A. Vittadini, A. Selloni, F.P. Rotzinger, M. Gratzel, *Phys. Rev. Lett.* 81 (1998) 2954.
 18. A. Beltran, J.R. Sambrano, M. Calatayud, F.R. Sensato, J. Andres, *Surf. Sci.* 490 (2001) 116.
 19. M.T.M. Koper, R.A. van Santen, *J. Electroanal. Chem.* 472 (1999) 126.
 20. R.A. van Santen, M. Neurock, *Catal. Rev.Sci.*(1995), Eng. 37. (557)
 21. M.L.Machesky ,The protonation behavior of metal oxide surfaces to hydrothermal conditions, *ECS Transactions* ,11(27) 151-166(2008).
 22. A. Cordoba ,I. Magarinoa,1, M. L. Ferreir. Experimental design and MM2–PM6 molecular modelling of hematin as a peroxidase-like catalyst in Alizarin Red S degradation / *Journal of Molecular Catalysis A:Chemical* 355(2012)44-60.
 23. A. Vittadini, A. Selloni, F. P. Rotzinger, and M. Gratzel. Formic Acid Adsorption on Dry and Hydrated TiO₂ Anatase (101) Surfaces by DFT Calculations, *J. Phys. Chem. B* 2000, 104, 1300-1306.
 24. C. I. Oprea 1, P. Panait, F. Cimpoesu, M. Ferbinteanu and M. A. Gîrțu. Density Functional Theory (DFT) Study of Coumarin- based Dyes Adsorbed on TiO₂ Nanoclusters—Applications to Dye-Sensitized Solar Cells, *Materials* 2013, 6, 2372-2392.
 25. A. Selloni, A. Vittadini, M. Gratzel, *Surf. Sci.* 402–404 (1998) 219.
 26. C. Redfern, P. Zapol, L.A. Curtiss, T. Rajh, M.C. Thurnauer, *J. Phys. Chem. B* 107 (2003) 11419.
 27. X.Q. Gong,; Selloni, A. Reactivity of anatase TiO₂ nanoparticles. The role of the minority (001) surface. *J. Phys. Chem. B* 2005, 109, 19560–19562.
 28. M. Vittoria, E. Selli, Specific Facets-Dominated Anatase TiO₂: Fluorine-Mediated Synthesis and Photoactivity, *Catalysts* **2013**, 3, 455-485.3
 29. S. W. Radi, M.Sc. Thesis ,Salah Al-Den University, Iraq, 1989.
 30. W. Choi and Hoffmann M.R. (1997) Novel photocatalytic mechanism for CHCl₃, CHBr₃ and CClCO₂ - degradation and the fate of photogenerated trihalomethyl radicals on TiO₂, *Environ.Sci. Technol.*, 37, 89-95.
 31. P.R. Gogate and Pandit A.B. (2004a) A review of imperative technologies for wastewater treatment I: oxidation technologies at ambient conditions, *Adv. Environ. Res.*, 8, 501-551.
 32. S. A. Shapovalov, Ionic association with Inions of Alizarin Red S in Aqueous Solutions with Surfactants, *Russian Journal of physical Chemistry A*, 2011, Vol.85, No.8, pp, 1369-1375.
 33. T. MAHMUD , Removal of Alizarin Red S (Dye) from Aqueous Media by using Alumina as an Adsorbent , *J.Chem.Soc.Pak.*, Vol. 33, No. 2, 2011.P (230).
 34. H. V. Jadhava, Removal of alizarin red-S from aqueous solution by adsorption on nanocrystalline Cu_{0.5}Zn_{0.5}Ce₃O₅, *Der Chemica Sinica*, 2011, 2(6):p (72)
 35. I. Onal, S. Soyer , S. Senkan. Adsorption of water and ammonia on TiO₂-anatase cluster models, *Surface Science* 600 (2006) 2457–2469.
 36. H. Liu. Formaldehyde on TiO₂ anatase (101): A DFT study, *Computational Materials Science* 51 (2012) 389–395.
 37. LAZZERI et al. Structure and energetics of stoichiometric TiO₂ anatase surfaces, *PHYSICAL REVIEW B, VOLUME* 63, 155409(1-9).
 38. Y. Ying . First-principle study on the electronic and optical properties of the anatase TiO₂ (101) surface, *J. Semicond.* 2013, 34(7), 073004(1-5)
 39. W. Zeng, T. Liu, Z. Wang, S. Tsukimoto. Oxygen Adsorption on Anatase TiO₂ (101) and (001) Surfaces from First Principles, *Materials Transactions*, Vol. 51, No. 1 (2010) pp. 171 to 175.

* Tel.: +9647902986137 .

E-mail address: ramziealani@yahoo.com.

- 468 40. G.S. Herman,; Sievers, M.R.; Gao, Y. *Structure determination of the two-domain (1 × 4)*
469 *anatase TiO₂(001) surface. Phys. Rev. Lett. 2000, 84, 3354–3357.*