

RELATIONSHIP OF VISCERAL OBESITY AND BMI WITH BLOOD PRESSURE AMONG OSUN STATE COLLEGE OF EDUCATION STAFF, NIGERIA

ABSTRACT

Aims: To determine prevalence of visceral obesity, compare visceral obesity and Body Mass Index (BMI) association with Blood Pressure BP.

Study design: Research was a cross-sectional survey study design.

Place and Duration of Study: The study took place in Osun State College of Education, Ilesa, Osun State, Nigeria, between November 2015 and April 2016.

Methodology: We included 231 members of staff. A semi-structured instrument (questionnaire) was used for data collection. Waist circumference was measured with a measuring tape. BMI was measured with a weighing scale and stadiometer. BP levels were determined with electronic sphygmomanometer. Data collected were analyzed using Statistical Package for Social Science (SPSS) version 21.

Results: There were 117 (50.7%) male and 99 (42.9%) female participants between the ages of 26 and 66 years. The results revealed that the prevalence of visceral obesity was high (71.4%) among the respondents while prevalence of 88.9% was recorded among the female participants. Dietary pattern of respondents was poor (35.16%), it was significantly associated with the development of visceral obesity [$R^2 = 0.190$; $F = 7.47$; $P < .001$]. Social pressure from immediate family, was detected to be significantly associated with respondents' intention towards reducing visceral obesity [$R^2 = 0.166$; $F = 45.704$; $P < .01$]. Findings also revealed that visceral obesity was significantly associated with BP [$R^2 = 0.155$; $F = 20.989$; $P < .01$] and also significantly associated with BMI [$R^2 = 0.044$; $F = 5.235$; $P = 0.006$].

Conclusion: Visceral obesity was more significantly associated with high BP than BMI; hence visceral obesity was a more accurate predisposing risk factor than BMI, in predicting susceptibility to hypertension. This study will improve health promotion, while creating awareness about visceral obesity.

Keywords: Abdominal obesity, hypertension, cardiovascular diseases, visceral obesity and BMI

1. INTRODUCTION

Obesity affected up to 33% of the industrialized world population [1]. It was stated that chronic diseases accounted for 24% of all deaths in Nigeria with cardiovascular disease accounting for 7% of this [2]. As at 2005, only 29% of men were overweight and 39% of women were overweight. The prevalence of overweight was predicted to increase in both men and women, to 39% and 49% respectively by 2015 [3]. Researches have revealed different risk factors of cardiovascular disease (CVD), among which was obesity. Abdominal obesity was the type of obesity revealed to be an important factor to diabetes and other cardiovascular diseases as they are unique pathogenic fat depot [4-5]. The risk of CVD has been revealed to be strongly associated with Visceral Adipose Tissue, measurable using waist circumference [6]. Visceral fat distribution has been associated with higher prevalence or incidence of CVD influencing the need for waist circumference measurement to estimate the full impact of the worldwide obesity epidemic [7-8].

A study [9] revealed that there were two different depots of abdominal fat: the intra-abdominal (visceral) obesity (excess fat in the abdominal cavity) and abdominal subcutaneous fat (the fat located just under the skin). It is believed that these fat serves as energy sink, where excess energy is stored. The inability of the subcutaneous adipose tissue to store the excess energy due to elevated visceral fat, may cause the fat to move and accumulate at undesired locations like the heart, liver and muscles (ectopic fat deposition) thereby increasing the risk of CVD [10].

The combination of cardio-metabolic risk factors are referred to as 'Metabolic syndrome (MetS); and in 2005, the American Heart Association proposed that any 3 of the following 5 criteria constitute diagnosis of MetS: elevated BP, elevated waist circumference, elevated triglycerides, reduced High Density Lipoprotein cholesterol and elevated fasting glucose [11], this was further reviewed by the International

Diabetes Federation (IDF), a new definition of MetS was established concluding that for an individual to be defined as having the MetS, they must have visceral obesity plus any two of the remaining four risk factors [12]. Elevated waist circumference (visceral obesity) has been proven to be an independent risk factor for some chronic diseases: type 2 diabetes mellitus, CVD (hypertension, coronary artery disease, and stroke), kidney cancer, non-alcoholic fatty liver disease (NAFLD) [13-15]. Visceral fat however is predisposed by inappropriate dietary habit and is greater in men than in women [16]. In determining waist circumference for visceral obesity, there is variability according to ethnicity and nationality; the International Diabetes Federation established a standard waist circumference according to ethnicity and waist circumference of less than 94cm and less than 80cm was recommended for male and female of sub-Saharan African region respectively [17]. (Table 1)

It is possible that some obese patients, with lower-than-expected visceral fat may not have clinical signs of CVD, whereas patients who may be moderately overweight with higher-than-expected visceral fat have metabolic profile that may predispose them to CVD as disease risk are more closely related with visceral fat rather than the total body fat volume [10]. BMI has been used in evaluating body weight and likelihood of developing CVD, it has been indicated that waist-hip circumference is a better measurement of body weight in association with CVD than BMI [18]. A further study [19] revealed that Waist Circumference is a superior anthropometric measurement in diagnosing MetS to both Waist Hip Ratio and BMI using both NCEP and WHO criteria.

There has been a reported rise in prevalence of obesity in sub-Saharan African countries especially in sub-urban populations [20]. The rise has been revealed to be attributed to physical inactivity, unhealthy weight gain behaviors, increased sedentary nature, intake of high caloric fast food and sugar sweetened beverages, relative household wealth and decreased mental health [21-24].

Urbanization is a critical factor that has influenced the traditional ideal body image among Africans, who have always been inclined towards a larger, fuller body shape as obesity was also associated with dignity, health, wealth and respect in view of the “big is beautiful” mindset [25-26]. Obesity has been revealed to be particularly common among urban women [27].

Researches have been carried out in Nigeria to evaluate the prevalence of obesity, with results indicating 10.5% prevalence of obesity in Remo local government [28], another research [23] revealed 62.2% prevalence of obesity among nurses in Akwa Ibom state. Also a 49.34% prevalence of obesity was recorded among the Kalabaris in the Niger Delta region of South-South Nigeria [29].

Table 1: Values of waist circumference according to ethnicity [17]

ETHNICITY	WAIST CIRCUMFERENCE VALUE	
European	Male	>94 cm
	Female	>80 cm
South Asians	Male	>90 cm
	Female	>80 cm
Chinese	Male	>90 cm
	Female	>80 cm
Japanese	Male	>85 cm
	Female	>90 cm
South and Central Americans	Male	>90 cm
	Female	>80 cm
Sub Saharan Africans	Male	>94 cm
	Female	>80 cm
Eastern Mediterranean and Middle East	Male	>94 cm
	Female	>80 cm

Obesity is one of the strongest predictor of hypertension [30]. An increase in rate of obesity subsequently leads to high prevalence of cardiovascular diseases such as high BP, diabetes and stroke. Some studies have revealed that some ethnic groups in Africa historically preferred overweight women and embraced cultural practices that encouraged female obesity and attribute it to health and beauty [31], our society nowadays encourages overweight and obesity as it is seen as a sign of wealth [25-26]. This study aims at increasing awareness on the association of visceral obesity with BP and also to advocate for the use of waist circumference measurement in evaluating obesity.

This research will determine prevalence of visceral obesity, compare visceral obesity and BMI's association with BP in the study area. Results of this study will enhance early diagnosis of hypertension and its control.

2. MATERIAL AND METHODS

Participants were drawn from members of staff of the Osun State College of Education (OSSCOED) Ilesha, all participants reside in Ilesha community and work at the institution. They were within an age range of between 20 and 70 years. This choice of respondents were made because of the increasing prevalence of high BP among members of staff of the institution.

Random sampling method was used in selecting the participants for this study. Sample proportion was drawn using Cochran's formula [32]:

$$n_0 = \frac{Z^2 pq}{e^2}$$

The sample size was approximated to 250 participants. This was done to take care of attrition, in order to account for loss or missing questionnaires (instrument).

2.1 Instruments for data collection

The instrument for data collection in this study was a self-developed questionnaire, measuring tape, weighing scale, stadiometer and sphygmomanometer. The measuring tape was used in measuring the waist circumference of individuals in centimeter (cm) with cutoff points of 94cm for male and 80cm for female; the weighing scale and stadiometer was used in measuring weight and height of respondents which was used in calculating respondents' BMI in unit value of kilogram per meter square (kg/m^2) with cutoff points of above 30kg/m^2 as obese and the sphygmomanometer was used in measuring the BP of respondents with cutoff points at 140 mmHg and 90 mmHg for systolic BP (SBP) and diastolic BP (DBP) [33].

The questionnaire evaluated participants' demographic factors, three multiple answers questions measured respondents' knowledge of visceral obesity, six questions measured attitude on visceral obesity and eight questions measured subjective norms on visceral obesity both measured on a 5-point likert scale of Strongly Agree (SA), Agree (A), Undecided (U), Disagree (D) and Strongly Disagree (DS), six questions measured perceived behavioral control on visceral obesity reduction and seven questions measured behavioral intention and dietary pattern both measured on a 2-point likert scale of 'yes' and 'no'. Seven questions measuring dietary pattern of respondents was also assessed, measured on a 5-point rating scale of Never (N), Not Sure (NS), Small Proportion (SP), Medium Proportion (MP) and Large Proportion (LP).

Reliability of the questionnaire was determined by pre-testing the questionnaire among populace of similar characteristics as the study sample with 30 respondents and the data obtained was analyzed using test statistics of Cronbach's reliability score with a score of 0.665.

2.2 Data Collection

Questionnaires were randomly administered among members of staff of the institution in the five schools which are schools of science, arts, languages, education and vocational and technical studies, members of staff in the registry were also administered the questionnaire for 3 weeks, with no fraction distribution across teaching and nonteaching staff.

The researcher recruited six research assistants, they were students from each of the schools mentioned above, and recruitment was done by the researcher from the 24 students nominated by the deans of each school using the simple random technique as they are more conversant with the members of staff. The six research assistants were trained on measurement and calculations of body mass index by converting measurement of height to meters and also using tape measurement for waist circumference correctly.

Blood pressure levels were taken by the researchers as most members of staff were not willing to be subjected to being measured by their students. The measurement was taken in the morning between 9am and 10am on all working days during the duration of data collection with an average of 15 respondents daily.

The questionnaire were administered to the respondents randomly and collected back upon taking the anthropometric measurements of waist circumference, body mass index and blood pressure levels.

A total number of 250 questionnaires were administered but 231 questionnaires were returned back to the researcher as some of the respondents did not give consent to the measurements and opted out of the study.

2.3 Data Analysis

The data collected were analyzed using Statistical Package for Social Science (SPSS) version 21. Frequency distribution was used in evaluating the demographic characteristics of respondents. Analysis of Variance (ANOVA), bivariate correlation, Regression, Independent Sample T-test and relevant descriptive statistics were used in analyzing the variables.

3. RESULTS AND DISCUSSION

The respondents were predominantly of Yoruba ethnicity (94.4%) with 117 (50.6%) males and 99 (42.9%) females. A high proportion of the respondents 86 (43.0%) were between the ages of 41 and 50 years and 200 (86.6%) respondents were married. Data collected revealed 148 (64.1%) academic members of staff and 68 (29.4%) non-academic members of staff. The study had 53 (45.3%) male respondents who were between the age of 41 and 50 years and 63 (63.6%) female respondents between the ages of 51 to 55 years. There were 18 (7.8%) unmarried respondents, who were of ages between 26 and 35 years with 9 (50%) between 26 and 30 years and 9 (50%) between 31 and 35 years; of the 200 (86.6%) married respondents, 86 (43%) were of ages 41 to 50 years. It was shown that 60 (40.5%) respondents who were academic members of staff between the ages of 41 to 50 years and 32 (47.1%) non-academic members of staff respondents within 51 and 55 years of age. Result indicated 86 (37.2%) respondents between the ages of 41 and 50 years and 81 (35.1%) respondents between the ages of 51 and 55 years. (Table 2) Visceral obesity was prevalent among respondents with 165 (71.4%) respondents with waist circumference higher than expected measurement (Fig. 1). An average percentage (53.0%) of male respondents with visceral obesity (WC > 94cm) when compared to the high percentage of 88.9% (WC > 80cm) in female respondents (Table 3).

Table 2: Socio-demographic characteristics of respondents

	GENDER		MARITAL STATUS		STAFF LEVEL		ETHNIC	TOTAL
	Male	Female	Single	Married	Academic	Non Academic	Yoruba	
	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)
AGE								
No respond	13 (11.1)	0 (0.0)	0 (0.0)	13 (6.5)	13 (8.8)	0 (0.0)	13 (0.0)	26 (11.3)
26 – 30	9 (7.7)	0 (0.0)	9 (50.0)	0 (0.0)	9 (6.1)	0 (0.0)	9 (4.1)	9 (3.9)
31 – 35	9 (7.7)	5 (5.1)	9 (50.0)	5 (2.5)	9 (6.1)	5 (7.4)	14 (6.4)	14 (6.1)
36 – 40	8 (6.8)	0 (0.0)	0 (0.0)	8 (4.0)	1 (0.7)	7 (10.3)	8 (3.7)	8 (3.5)
41 – 50	53 (45.3)	31 (31.3)	0 (0.0)	86 (43.0)	60 (40.5)	24 (35.3)	86 (39.4)	86 (37.2)
51 – 55	18 (15.4)	63 (63.6)	0 (0.0)	81 (40.5)	49 (33.1)	32 (47.1)	81 (37.2)	81 (35.1)
56 – 60	6 (5.1)	0 (0.0)	0 (0.0)	6 (3.0)	6 (4.1)	0 (0.0)	6 (2.8)	6 (2.6)
60 and above	1 (0.9)	0 (0.0)	0 (0.0)	1 (0.5)	1 (0.7)	0 (0.0)	1 (0.5)	1 (0.4)
TOTAL	117 (50.6)	99 (42.9)	18 (7.8)	200 (86.6)	148 (64.1)	68 (29.4)	218 (94.4)	231 (100)




Fig. 1: Visceral obesity among respondents

Table 3: Prevalence of visceral obesity among respondents

	Gender			Total
	No Response	Male	Female	
Non Obese	0	55	11	66
	0.0%	83.3%	16.7%	100.0%
Obese	15	62	88	165
	9.1%	37.6%	53.3%	100.0%
Total	15	117	99	231
	6.5%	50.6%	42.9%	100.0%

The level of knowledge of respondents on visceral obesity was below fifty percent at a score of 4.29 from a possible score of 9 (47.67%) with a low level of knowledge of 1.27 score from a possible 4 (31.75%) on the causes of visceral obesity. Among the respondents, 71% of the respondents had below fifty percent of knowledge regarding visceral obesity followed by 17.7% who had average knowledge and only 11.3% had good knowledge about visceral obesity. (Table 4)

Table 4: The percentage distribution of knowledge of respondents

LEVEL OF KNOWLEDGE	KNOWLEDGE SCORE	RANGE OF SCORE	F (%)
Good	71 – 90%	7 – 8	11.30%
Average	50 – 70%	5 – 6	17.70%
Below Average	Below 50%	0 – 4	71.00%
Minimum score = 0		Maximum score = 9	

There was an evaluated high level of intake of salt, pastries (meat pie, doughnuts, burger, cake etc.) and animal protein (red meat, fried chicken and turkey), and also a low intake of fruits and vegetable in daily meals. (Table 5)

Table 5: Dietary intake of respondents

Level of consumption	
Consume Salt	84.00%
Consume Pastries	81.25%
Consume Animal Protein	82.00%
Consume Soft Drinks	57.25%
Consume Alcohol	37.75%
Consume Fruits	9.75%
Consume Vegetable	17.00%

3.1 Discussion

Since abdominal obesity was significantly linked with cardiovascular diseases like type2 diabetes and hypertension [19, 34], it was necessary to compare relationship of waist circumference to BP and also relationship of BMI (BMI) to BP. This was important to further establish the importance of visceral obesity in diagnosing cardiovascular diseases [8].

The results obtained from this study, which was similar to the findings of other studies [35-39] showed a more significant association of Waist Circumference with BP compared to association of BMI with BP. This indicated that there was a higher chance of developing hypertension with elevated waist circumference, than with high BMI.

The results of this study indicated that visceral obesity was prevalent among the community studied. Marital strain has been shown to have deleterious effects on cardiovascular functions and significantly associated with development of visceral obesity [24, 40] this was reflected in the findings of the study as unmarried respondents were without visceral obesity compared to a high percentage of married respondents being viscerally obese, this might be as a result of change in diet as prepared by significant other.

Also, several studies have established that prevalence of visceral obesity among the female gender [28, 34, 41] this was reflected in the findings of the present study, with a higher proportion of the female respondents having visceral obesity as compared to the proportion of male respondents, this however

was contrary to the findings of [42] who found visceral obesity to be more prevalent among male respondents.

The present study indicated that there was a poor dietary pattern among respondents, with excessive consumption of salt, pastries, animal protein and a low intake of fruits and vegetables and significant relationship with development of visceral obesity was also established, this was similar to other studies [42-43] who established a relationship between dietary pattern and development of visceral obesity.

This study had two major implications to health promotion; Health promotion intervention which will be directed towards reduction of the prevalence of visceral obesity should not only be directed towards the obese individuals but towards their significant order (wives, husbands, children and other immediate family members). This can be done in organizing a family-theme health promotion program where it will have in attendance the members of family of the staff including spouses and children, and the risk associated to development of visceral obesity will be given and awareness will be created not only to the obese individual but the immediate family giving them increased level of knowledge. This will help especially in dietary pattern control and pressure to increase likelihood of the individuals to reduce their visceral obesity. On the community level, awareness can be created via the media houses like radio, television and print media. And the adoption of Waist Circumference as a preferred anthropometric measurement in diagnosis of obesity over Body Mass Index, since it has been established to be more accurate in measuring obesity and susceptibility to cardiovascular diseases.

4. CONCLUSION

This study has highlighted visceral obesity (using waist circumference) as an accurate anthropometric measurement for obesity, over BMI (BMI). This study has shown a more significant association of visceral obesity with high BP (hypertension) than BMI. This finding indicated that there was an increased likelihood of developing hypertension when an individual has elevated waist circumference (visceral obesity) compared to increased body mass index. This finding is remarkable because it will improve health promotion, while creating awareness about visceral obesity and its association with high BP.

CONSENT

Consent of respondents was sought before administering the questionnaire (instrument).

ETHICAL APPROVAL

Ethical clearance was obtained from Babcock University Health Research Ethics Committee (BUHREC). Participants were informed of the purpose of the study and they were assured that confidentiality would be maintained. Their consent was sought for and granted, also participation was voluntary. They were informed of and their right to refuse to answer any of the questions or withdraw from the study at any point in this study. However, there was no reward or compensation for participating.

REFERENCES

1. Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999–2004. JAMA. 2006;295:1549–55.
2. World Health Organization. Noncommunicable Diseases (NCD) Country Profiles, 2014.
3. World Health Organization. The Impact of Chronic Disease In Nigeria; Facing The Facts. 2005. Accessed 23 December 2015. Available: http://www.who.int/chp/chronic_disease_report/en/
4. Mahabadi AA, Massaro JM, Rosito GA, Levy D, Murabito JM, Wolf PA, *et al.*. Association of pericardial fat, intrathoracic fat, and visceral abdominal fat with cardiovascular disease burden: The Framingham Heart Study. European Heart Journal Advance Access. 2009. doi:10.1093/eurheartj/ehn573
5. Sandeep S, Gokulakrishnan K, Velmurugan K, Deepa M, Mohan V. Visceral & Subcutaneous Abdominal Fat In Relation To Insulin Resistance & Metabolic Syndrome In Non-Diabetic South Indians. Indian J Med Res. 2010.131:629-635.
6. Fox CS, Massaro JM, Hoffman U, Pou KM, Maurovich-Horvat P, Liu Chun-Yu, *et al.*, Abdominal Visceral and Subcutaneous Adipose Tissue Compartments Association With Metabolic Risk Factors in the Framingham Heart Study. Semmelweis Univ Medicine. 2013. Available: <http://circ.ahajournals.org>
7. James PT, Leach R, Kalamara E, Shayeghi M. The Worldwide Obesity Epidemic. Obesity Research. 2001.9:

8. Scaglione R, Chiara TD, Argano C, Corrao S, Licata G. Hypoadiponectinemia: A link between visceral obesity and metabolic syndrome. *Journal of Diabetes and Endocrinology*. 2010.1(3):27-35
Available: <http://www.academicjournals.org/JDE/>
9. Fang L, Kojima K, Zhou L, Crossman DK, Mobley JA, Grama J. Analysis of the Human Proteome in Subcutaneous and Visceral Fat Depots in Diabetic and Non-diabetic Patients with Morbid Obesity. *J Proteomics Bioinform*. 2015.8:133-141. doi:10.4172/jpb.1000361
10. Despres JP, Lemieux I. Abdominal obesity and metabolic syndrome: *Nature*. 2006.444:881-887
11. Grundy SM. Metabolic syndrome scientific statement by the American Heart Association and the National Heart, Lung, and Blood Institute. *Arterioscl. Thromb. Vasc. Biol*. 2005.25: 2243-2244
12. International Diabetes Federation. The IDF consensus worldwide definition of the metabolic syndrome. Memo. Avenue Emil De Mot, Brussels, Belgium.
13. Haslam DW, James WPT. Obesity. *The Lancet*. 2005.366(9492):1197-1209.
14. Guh DP, Zhang W, Bansback N, Amarsi Z, Birmingham CL, Anis AH. The incidence of co-morbidities related to obesity and overweight: A systematic review and meta-analysis. *BMC Public Health*. 2009. 9(1):88.
15. Leitzmann MF, Moore SC, Koster A, Harris TB, Park Y, Hollenback A *et al*. Waist Circumference as Compared with Body-Mass Index in Predicting Mortality from Specific Causes. *PLoS ONE*. 2011.
16. Adefule AK, Adu TO, Ogundahunsi OA, Caxton-martins EA, Otulana OJ & Adesanya OA. Prevalence of Overweight and Central Obesity Among South Western Nigerian Young Adults. *Nova Journal of Medical and Biological Sciences*. 2014.2(3):1-10.
17. Shrivastava S, Shrivastava P, Ramasamy J. Assessment of knowledge about obesity among students in a medical college in Kancheepuram district, Tamil Nadu. *Prog Health Sci*. 2013.3(1):54-60
18. Nkeh-Chungag BN, Mxhosa TH & Mgoduka PN. Association of waist and hip circumferences with the presence of hypertension and pre-hypertension in young South African adults. *African Health Sciences*. 2015.15(3): 908-916.
19. Ghazali SM, Sanusi RA. Waist circumference, waist to hip ratio, and BMI in the diagnosis of metabolic syndrome in Nigerian subjects. *Nigerian Journal of Physiological science*. 2010.25:187-195
20. Bakari AG, Onyemelukwe GC, Sani BG, Aliyu IS, Hassan SS, Aliyu TM. Obesity, Overweight and Underweight in Suburban Northern Nigeria. *International Journal of Diabetes and Metabolism*. 2007.15:68-69
21. Duda RB, Dark R, Seffah J, Adanu RMK, Anarfi JK, Hill AG. Prevalence of Obesity in Women of Accra, Ghana. *African Journal of Health Sciences*. 2007.14:154-159.
22. Kivimaki M, Batty D, Singh-Manoux A. Association between common mental disorder and obesity over the adult life course. *Br J Psychiatry*. 2009.195(2):149-155.
23. Ogunjimi LO, Ikorok MM, Yusuf, Olayinka O. Prevalence of obesity among Nigeria nurses: The Akwa Ibom State experience. *International NGO Journal*. 2010.5:45-49
Available: <http://www.academicjournals.org/ingoj/>
24. Benkeser RM, Biritwum R, Hill AG. Prevalence Of Overweight And Obesity And Perception Of Healthy And Desirable Body Size In Urban, Ghanaian Women. *Ghana Medical Journal*. 2012.46(2):66-75
25. Szabo CP, Grange DL. Eating disorders and the politics of identity. *Eating disorders and cultures in transition*. New York: Brunner Routledge. 2001: 24-39.
26. Puoane T, Fourie JM, Shapiro M. 'Big is beautiful': An exploration with urban black community health workers in a South African township. *South African Journal of Clinical Nutrition*. 2005.18(1):6-14.
27. Katchunga P, Jean-René M, Buyamba-Kayamba, Masumbuko E, Lemogoun D, Kashongwe Z *et al*. Hypertension in the adult Congolese population of Southern Kivu: Results of the Vitaraa Study. *Presse Med*. 2011.40(6):315-323.
28. Raimi TH, Odusan O, Fasanmade O. High prevalence of central obesity in rural south-western Nigeria: need for targeted prevention. *Journal of Diabetes and Endocrinology*. 2015.6:12-18
29. Adienbo OM, Hart VO, Oyeyemi WA. High Prevalence of Obesity among Indigenous Residents of a Nigerian Ethnic Group: The Kalabaris in the Niger Delta Region of South-South Nigeria. *Greener Journal of Medical Sciences*. 2012.2:152-156.
30. Kaufman JS, Asuzu MC, Mufunda J, Forrester T, Wilks R, Luke A *et al*. Relationship between BP and BMI in Lean Populations. *Hypertension*. 1997.30:1511-1516
31. Prinsloo EAM, Joubert G, Mohale M, Nyindi N, Matu N, Ntechane L. *et al*. The prevalence and

perception of obesity and its association with the lifestyle of women at the Mangaung University
Community Partnership Project healthcare Centre, Bloemfontein. *S Afr Fam Pract.* 2011.53(4): 366-372

32. Cochran WG. Sampling Techniques. 1963. 2nd Ed., New York: John Wiley and Sons, Inc.

33. ESH/ESC Guidelines on the Management of Arterial Hypertension. *European Heart J* 2013.34:2159-219 doi:10.1093/eurheartj/eh151

34. Freedman DS, Rimm AA. The relation of Body Fat Distribution, as assessed by six Girth measurement, to Diabetes Mellitus in women. *American Journal of Public Health.* 1989.79(6):715-720.

35. Abolfotouh MA, Daffallah AA, Khan MY, Khattab MS, Abdulmoneim I. Central obesity in elderly individuals in South-western Saudi Arabia: prevalence and associated morbidity. *La Revue de santé de la Mediterranee orientale.* 2001.7:716-724

36. Martins IS, Marinho SP. The potential of central obesity anthropometric indicators as diagnostic tools. *Rev saudePublica.* 2003.37(6)

37. Pimenta AM, Kac G, Gazzinelli A, Correa-Oliveira R, Velasquez-Melendez G. Association Between Central Obesity, Triglycerides and Hypertension in Rural area in Brazil. *Arq Bras Cardiol.* 2008.90(6):386-392

38. Giday A, Wolde M, Yihdego D. Hypertension, obesity and central obesity in diabetics and non diabetics in Souther Ethiopia. *Ethiop J. Health Dev.* 2010.24 (2):145-147

39. Okafor CI, Fasanmade O, Ofoegbu E, Ohwovorile AE. Comparison of the performance of two measures of central adiposity among apparently healthy Nigerians using the receiver operating characteristics analysis. *Indian Journal of Endocrinology and Metabolism.* 2011.15(4):320-326.

40. Robles TF, Kiecolt-Glaser JK. The physiology of marriage: pathways to health. *PhysiolBehav.* 2003.79(3):409-416.

41. Mogre V, Aleyira S, Nyaba R. Factors associated with central overweight and obesity in students attending the University for Development Studies in Tamale, Ghana: a cross-sectional study. *South African Journal of Clinical Nutrition.* 2014.27(2):69-74

42. Adefule AK, Adu TO, Ogundahunsi OA, Caxton-martins EA, Otulana OJ, Adesanya OA. Prevalence of Overweight and Central Obesity Among South Western Nigerian Young Adults. *Nova Journal of Medical and Biological Sciences.* 2014.2(3):1-10

43. Moreira PL, Corrente JE, Boas PJFV, Ferreira LA. Dietary patterns are associated with general and central obesity in elderly living in Brazilian City. *Rev Assoc Med Bras.* 2014.60(5):457-464

INFORMED CONSENT FORM

RELATIONSHIP BETWEEN VISCERAL OBESITY, BMI AND BP IN ILESA COMMUNITY, OSUN STATE, NIGERIA.

My name is Posi Aduroja, an MPh student in the Department of Public Health, Babcock University, Ilishan-Remo, Ogun state. The title of my research work as shown above is self-sponsored. The purpose of this study is to determine the prevalence of visceral obesity among members of staff of the Osun state college of education, Ilesa, Osun state. It is also suggest substantial recommendation of reducing visceral obesity, in order to prevent prevalence of cardiovascular diseases. The research work which will remain in the domain of academic environment will serve the interest of students and the entire academic community and may be published in an academic journal.

The outcome of this research will help health officers, parents and lecturers in living a healthy live and implementing program that will encourage healthy society free of cardiovascular diseases.

The survey method will be used for data collection and analysis. Should you decide not to continue with the research, for any reason, be assured that you will not be penalized in any way. You are free to withdraw at any stage of the research if you so wish, as there is no conflict of interest whatsoever.

You are however, assured that your response will be treated with utmost confidence and will be used only for academic purposes.

Please bear with me as there will be no reward or compensation for your participation in this research work. If you agree with these terms, please write your name and sign this consent form on the space provided below. Thank you.

Name, Signature & Date

Witness Signature

SECTION A: SOCIO-DEMOGRAPHIC CHARACTERISTICS

INSTRUCTION: Please kindly tick [✓] appropriately and supply adequate responses to the questions provided.

1. Age group: 20-25 [] 26-30 [] 31-35 [] 36-40 [] 41-50 [] 51-55 [] 56-60 []
60 and above []
2. Gender: Male [] Female []
3. Marital status: Single [] Married [] Divorced/Separated []
4. Level of staff: Academic Staff [] Non-Academic Staff []
5. Ethnic: Yoruba [] Igbo [] Hausa [] Others _____

SECTION B: ANTHROPOMETRIC MEASUREMENTS (for official use only)

6. Waist Circumference: _____ cm
7. BP: _____/_____ mmHg
8. Weight: _____ kg Height: _____ cm BMI: _____

SECTION C: KNOWLEDGE OF VISCERAL OBESITY

INSTRUCTION: Please kindly tick [✓] appropriately and supply adequate responses to the questions provided.

9. Visceral obesity is the accumulation of fat in the abdominal areas of the body.

True [] False []

10. The following are the causes of visceral obesity

- | | |
|--|-----|
| Excessive soft drink and alcohol consumption | [] |
| Mental disorder | [] |
| Insufficient sleep / Stress | [] |
| Lack of physical activities | [] |

11. Visceral obesity causes the following

- | | |
|------------------------|-----|
| High BP (hypertension) | [] |
| Typhoid | [] |
| Diabetes | [] |
| Glaucoma | [] |

SECTION D: ATTITUDE ON VISCERAL OBESITY**INSTRUCTION:** Please kindly tick [✓] appropriately and supply adequate responses to the questions provided.

SA = STRONGLY AGREE A = AGREE NS = NOTSURE D = DISAGREE SD = STRONGLY DISAGREE

S/N	STATEMENT FOR CONSIDERATION	SA	A	NS	D	SD
12	Big tummy helps you to gain respect					
13	Big tummy gives you self-confidence in social gathering					
14	Big tummy is a sign of healthy living					
15	Big tummy increases individual's chances of cardiovascular disease					
16	Reducing big tummy takes longer time than gaining					
17	Exercising and eating healthy allows you reduce big tummy					

SECTION E: SUBJECTIVE NORM**INSTRUCTION:** Please kindly tick [✓] appropriately and supply adequate responses to the questions provided.

SA = STRONGLY AGREE A = AGREE U – UNDECIDED D = DISAGREE SD = STRONGLY DISAGREE

S/N	STATEMENT FOR CONSIDERATION	SA	A	U	D	SD
18	My friends think I should reduce my big tummy					
19	I don't think it's necessary to reduce my big tummy					
20	My family members think I should engage in regular exercises					
21	Regular exercises is a waste of time					
22	TV programs suggest I should engage in healthy eating					
23	I don't believe in eating healthy					
24	My spouse think I should lose weight					
25	I don't think I need to lose weight					

SECTION F: PERCEIVED BEHAVIORAL CONTROL**INSTRUCTION:** Please kindly tick [✓] appropriately and supply adequate responses to the questions provided.

S/N	STATEMENT FOR CONSIDERATION	YES	NO
26	I'm willing to start eating healthy but I don't know which food is healthy		
27	Do you have enough discipline to maintain eating healthy?		
28	I want to exercise regularly but I don't have enough time		
29	If I know the benefits, I will be motivated to reduce my big tummy		
30	Unavailability of gym in my neighborhood doesn't allow for regular exercises		
31	I know big tummy is bad for my health but there is nothing I can do about it		

SECTION G: BEHAVIORAL INTENTION**INSTRUCTION:** Please kindly tick [✓] appropriately and supply adequate responses to the questions provided.

S/N	STATEMENT FOR CONSIDERATION	YES	NO
32	Will you engage in healthy eating subsequently?		
33	Will you engage in regular exercise subsequently?		
34	Will you stop consuming excessive animal protein?		
35	Will you start taking more vegetables and fruits?		
36	Will you stop taking excessive soft drinks and alcohol?		
37	Are you willing to reduce your intake of salt?		
38	Are you willing to reduce your consumption of pastries?		

399 **SECTION H: DIETARY PATTERN**
 400 **INSTRUCTION:** Please kindly tick [✓] appropriately and supply adequate responses to the questions provided.
 401 **N=NEVER NS=NOT SURE SP=SMALL PROPORTION MP=MEDIUM PROPORTION LP=LARGE PROPORTION**

S/N	STATEMENT FOR CONSIDERATION	N	NS	SP	MP	LP
39	I use salt in cooking my food					
40	I consume pastries like meat pie, doughnuts, burger, cake etc.					
41	I consume animal protein like red meat, fried chicken and turkey.					
42	I take soft drinks daily					
43	I take alcohol daily					
44	I include fruits in my daily diet					
45	I include vegetables in my daily diet					

402
403