

**Original Research Article**

**NUTRITIONAL QUALITY OF WEANING FOODS  
FORMULATED FROM MAIZE GRUEL 'OGI' AND  
CRAYFISH USING COMBINED TRADITIONAL  
PROCESSING TECHNOLOGY.**

**ABSTRACT**

**Aims:** To investigate the nutritional quality of weaning foods produced from maize gruel 'ogi' and crayfish using combined traditional processing techniques (germination, fermentation and toasting).

**Study design:** Randomized block design

**Place and Duration of Study:** Department of Food Science and Technology, Federal University of Technology, Akure, Ondo State, Nigeria, between January 2013 and November 2014.

**Methodology:** Maize grains were germinated at room temperature for three days after which they were fermented for 24 h. The maize grains were milled into slurry and divided into two portions. The first portion was oven-dried at 50°C and milled into flour while the second portion was toasted at 80°C and milled into flour. The two flours were separately mixed with crayfish powder to obtain oven-dried crayfish-ogi blend and toasted crayfish-ogi blend. The microbiological quality of the blends was determined. The nutritional qualities of the crayfish enriched ogi blends were assessed biologically using animal feeding experiment to determine the growth rate, feed intake, protein quality parameters and haematological properties. A commercial weaning food (cerelac) and traditional weaning food, ordinary ogi (maize gruel), were used as control diets

**Results:** The total mesophilic bacteria count of the ogi blends ranged from  $1.2$  to  $2.5 \times 10^3$  cfu/g. Mold ( $1.0 \times 10^3$  cfu/g) were found in both oven-dried and toasted crayfish enriched ogi blends. Yeasts were found only in oven-dried enriched ogi blend ( $1.0 \times 10^3$  cfu/g). Coliform, *Staphylococcus* and *Salmonella* were not detected in all the formulated diets. The growth rate of animals fed with crayfish enriched-ogi blends were lower than those fed with the cerelac, but higher than those fed with ordinary ogi. The protein efficiency ratio of animals fed with crayfish enriched ogi blends was similar ( $p=0.05$ ) to those fed with cerelac diet. The net protein ratio, true digestibility, biological value and net protein utilization of animals fed with crayfish enriched ogi diet were significantly lower ( $p<0.05$ ) than those fed with cerelac diet. The weight of the heart, liver, spleen and kidney of animals fed with crayfish-enriched ogi blends were significantly higher ( $p=0.05$ ) than those fed with ordinary ogi but similar to the rats fed with casein and cerelac diets. The haematological variables of animals fed with crayfish enriched ogi diets, commercial weaning food (cerelac) and casein diet were not significantly ( $p>0.05$ ) influenced by the dietary treatment.

**Conclusion:** Crayfish enriched ogi has potential as a functional weaning food with adaptable production technology (toasting) especially among rural dwellers.

**Keywords:** Quality evaluation, crayfish, enrichment, Ogi, germination, fermentation, toasting

## 14 1. INTRODUCTION

15 When breast milk is no longer enough to meet the nutritional needs of the infant at  
16 the age of four or six months and above, complementary foods (i.e., traditional or  
17 commercial weaning foods) should be added to the diet of the child. Several commercial  
18 weaning foods are marketed in Nigeria, but they are too expensive for people of low socio-  
19 economic status, especially those in the rural areas [1]. The most popular traditional weaning  
20 food in Nigeria which is fermented maize gruel known as 'ogi' has been implicated in the  
21 etiology of protein – energy malnutrition (PEM) in children during weaning period. This may  
22 be due to the low nutritive value characterized by low protein, low energy density and high  
23 bulk [2]. There is therefore a need to develop weaning foods with adequate protein that will  
24 promote growth in children from cheap raw material using processing methods that are  
25 adaptable to village level or at home.

26 Food processing techniques such as roasting, germination, milling, cooking, drying,  
27 fermentation and extrusion have the potential to enhance the nutrient bioavailability, nutrient  
28 density, food safety, storage stability, palatability, and convenience of supplementary foods  
29 suitable for infant mixtures [3]. Germination, fermentation and toasting of cereals are  
30 affordable and widely practiced processing techniques in Africa [4]. Fermentation enhances  
31 the nutrient of foods through biosynthesis and bioavailability of vitamins, essential amino  
32 acids, reduction of antinutrients improving the protein quality and fibre digestibility [5, 6].  
33 Germination unlocks many nutrients which are in bound forms in the food, thereby  
34 increasing nutrient bio-availability, energy density and acceptability of the food [7, 8].  
35 Toasting reduces anti-nutrients, improves the taste and nutrient quality of the food product  
36 and lowers the moisture content of such food product thereby increasing its shelf life [9]. An  
37 integrated approach that combines a variety of the traditional food processing techniques in  
38 the preparation of weaning food, including the addition of small amount of animal-source  
39 foods has been reported to be the best strategy to improve the nutrient content and  
40 bioavailability of micro- nutrients in plant-based diets in resource-poor settings [10]. The  
41 combination of two or more food processing techniques is more effective in removing  
42 antinutritional factors in cereal, thereby producing high nutrient dense weaning food [11]

43 Crayfish which is classified as an animal polypeptide is a freshwater crustacean resembling  
44 small lobster and it is commonly found in Nigerian coastal water. Crayfish is relatively cheap,  
45 affordable and readily available throughout the year. A review of nutritional value of crayfish  
46 showed that it is a good source of protein (36 -45%) with a superior biological value, true  
47 digestibility, net protein utilization, high content of essential amino acid, and protein  
48 efficiency is favourable compared to casein [12,13]. It is very low in carbohydrate but rich in  
49 vitamin D, A and mineral elements such as calcium, potassium, copper, zinc and iodine, [14,  
50 15].

51 In the effort to curb problem of protein-energy malnutrition (PEM) among the infants in  
52 Nigeria, a number of weaning foods have been formulated from locally available food  
53 materials [2,16,17,18]. Most of these formulated complementary foods are still not  
54 accessible to many nursing mothers, as a result of the high cost of food materials and  
55 production processes [19]. The present study is therefore aimed at producing weaning foods  
56 from ogi flour and crayfish flour mixes using a combination of traditional processing  
57 techniques and evaluates the microbiological and nutritional quality of the formulated diets.

## 2. MATERIALS AND METHODS

### 2.1 Materials

White maize (*Zea mays*), white crayfish (*Euastacus spp*) and commercial weaning food (cerelac) were purchased from Oba market in Akure Ondo state Nigeria.

### 2.2 Preparation of crayfish enriched ogi blends

The maize grains were soaked overnight after which they were germinated for 3 days. The germinated grains were dried at 60 °C for 14 hours and the radicles were removed. The germinated grains were steeped in water for 24 hours for fermentation to take place. The germinated - fermented grains were wet-milled, sieved and the slurry obtained was allowed to settle after which it was dewatered using muslin cloth. Ogi cake obtained was pulverized, sieved and divided into two portions. The first portion was oven dried at 50 °C for 24 hours while the second portion was toasted at 70 °C to 80°C using open cast iron. The crayfish were cleaned and milled into flour. The two ogi flours were separately mixed with crayfish powder in ratio of 80:15 respectively to obtain oven-dried crayfish-ogi blend and toasted crayfish-ogi blend. The choice of these mixing ratios was based on the target protein which is 18%. This mixing ratio was determined by using Quarto pro 8 software programme.

### 2.3 Microbiological analysis

The formulated weaning diets were examined microbiologically using the procedure of Olutiola *et al.*, [20] after serial dilution. The total microbial load was determined using nutrient agar in a plate count while molds and yeasts were examined using potato dextro agar. *Staphylococcus aureus*, coliform and *Salmonella* were determined using manitol salt agar, macconkey agar and deoxycholate citrate agar respectively.

### 2.4 Experimental Diets

The experimental diets which consist of formulated diets (crayfish enriched-ogi diets), commercial weaning food (cerelac) and casein were prepared at 10% protein level (iso-nitrogenous diets). A Basal diet (ordinary ogi) was also prepared. Composition of experimental diets is shown in Table 1. Diet 1 is the basal diet (ordinary ogi), diet 2 is the control (casein diet) while diets 3, 4 and 5 are cerelac, oven dried crayfish enriched ogi and toasted crayfish enriched ogi respectively.

Table 1: Composition of experimental diet (g/100g)

Ingredients	N-free diet	Casein diet	Cerelac diet	Oven-dried enriched ogi diet	Toasted enriched ogi diet
Ordinary ogi	71.80	60.30	5.63	16.24	16.24
Casein	-	11.50	-	-	-
Cerelac	-	-	66.67	-	-
Oven-dried enriched ogi	-	-	-	55.56	-

Toasted enriched ogi	-	-	-	-	55.56
Glucose	5.00	5.00	5.00	5.00	5.00
Sucrose	10.00	10.00	10.00	10.00	10.00
Non-nutritive cellulose	5.00	5.00	5.00	5.00	5.00
Vegetable oil	5.00	5.00	5.00	5.00	5.00
Mineral mixture	2.00	2.00	2.00	2.00	2.00
Vitamin mixture	1.00	1.00	1.00	1.00	1.00
NaCl	0.2	0.2	0.2	0.2	0.2
Total	100.00	100.00	100.00	100.00	100.00

90

## 91 2.5 Animal Experiment

92 In this study, thirty weanling albino rats of the Wistar strain weighing between 30 – 65g at the  
93 beginning of experiment were obtained from the Department of Biochemistry, University of  
94 Ilorin, Kwara State, Nigeria. The rats were weighed and divided into five groups. They were  
95 randomly distributed in metabolic cages and fed on normal (pellet) diets for a period of 7  
96 days for proper acclimatisation to the environment before commencement of the  
97 experiments. After the acclimatisation period, the animals were then re-weighed and  
98 grouped into five groups of six rats each per group such that the differences in their mean  
99 weights were  $\pm 2g$ . Two groups of animals were administered with the formulated diets  
100 (oven-dried crayfish enriched ogi and toasted crayfish enriched ogi). The remaining three  
101 groups of animals were administered with cerelac (a commercial weaning food), ordinary ogi  
102 and casein. Food and water were provided *ad libitum* to the rats for 28days. During this  
103 period dietary intake per day and weight of the animals were recorded. Five days before the  
104 end of feeding experiment, the faeces and urine were collected separately from each rat and  
105 pooled together at the end of the experiment. Pooled samples of faeces were dried in an  
106 oven at 80°C for 12 hours, cooled and weighed. A few drops of dilute sulphuric acid ( $H_2SO_4$ )  
107 were added to the urine, which was kept under frozen conditions. Nitrogen in the urine and  
108 faeces was determined by micro-Kjeldahl method [21]. The biological value (BV), true  
109 digestibility (TD), net protein utilization (NPU), protein efficiency ratio (PER), feed efficiency  
110 ratio (FER) and net protein ratio (NPR) were calculated.

## 111 2.6 Haematological Evaluations

112 At the end of the experiment, all the rats were starved for 3 hours and weighed after which  
113 each rat was anaesthetised and sacrificed. Blood samples from each rat were collected into  
114 sample bottles containing a few milligram of EDTA prior to haematological analysis. The  
115 packed cell volume (PCV) was estimated by spinning about 75 $\mu$ l of each blood sample in  
116 heparinised capillary tubes in a haematocrit microcentrifuge for 5 minutes, and the total red  
117 blood cell (RBC) and white blood cell (WBC) counts were determined. The haemoglobin  
118 concentration (Hb) was estimated using the cyano-methaemoglobin concentration method,  
119 while the lymphocyte, neutrophil, monocyte, basophil and eosinophil were determined [22,

120 23]. The heart, lungs, spleen, kidneys and liver were removed, blotted free of blood and  
121 weighed [22]. The values were subsequently expressed in g/kg of body weight

## 122 2.7 Statistical analysis

123 Data were collected as means of three separate determinations and subjected to one-way  
124 analysis of variance using Statistical Package for Social Statistics (SPSS 15.0). The  
125 significant differences ( $p \leq 0.05$ ) between the mean values were determined using the  
126 Duncan's Multiple Range Test.

## 127 3. RESULTS AND DISCUSSION

### 128 3.1 Microbial analysis

129 The result of coliform, *Staphylococcus*, *Salmonella*, mould, yeast and total viable count of  
130 the formulated diets are shown in Table 2. Coliform, *Staphylococcus* and *Salmonella* spp  
131 were absent in the formulated diets. This shows that the food will be fit for human  
132 consumption. The total viable count in all the formulated diets are below the maximum level  
133 of  $1.0 \times 10^5$  recommend by PAG [24]. However, all the formulated weaning diets would  
134 require cooking before feeding to children during which most of these microorganisms would  
135 be destroyed. The reduction in the total viable count of toasted enriched ogi diets may be  
136 due to toasting which was done at high temperature. (70-80°C) and might have destroyed all  
137 the pathogenic microorganisms.

138

139

140 Table 2: Microbiological quality of enriched ogi (cfu/g)

Micro-organism	counts	
	Ovendried	Toasted
Coliform	0	0
<i>Staphylococcus</i>	0	0
<i>Salmonella</i>	0	0
Molds	$1 \times 10^3$	$1 \times 10^3$
Yeast	$1 \times 10^3$	0

$1.2 \times 10^3$

161 D = Days of feeding


162 Fig. 1: Growth rate of rats fed with formulated weaning diets, cerelac and casein


163 Table 3: Nutrient utilization of rats fed with enriched ogi, casein and cerelac

Parameters	Casein	Cerelac	Oven-dried enriched ogi	Toasted enriched ogi
Food intake (g)	164.6 <sup>a</sup>	182.2 <sup>a</sup>	148.5 <sup>b</sup>	149.6 <sup>b</sup>
Protein intake (g)	17.8 <sup>a</sup>	18.5 <sup>a</sup>	14.5 <sup>b</sup>	15.0 <sup>b</sup>
Nitrogen consumed	0.61	0.62	0.47	0.56
Feecal nitrogen	0.15	0.20	0.20	0.23
Urinary nitrogen	0.09	0.13	0.14	0.17
Nitrogen retained	0.37	0.28	0.13	0.16


164 Values followed by different superscript on the same row are significantly different (p=0.5)

165 The results of feed efficiency ratio (FER), protein efficiency ratio (PER) and net protein ratio (NPR) of the  
 166 experimental animals are shown in Figure 2. The feed efficiency ratio (FER) of different diets varied from  
 167 0.22g (toasted enriched ogi diet) to 0.26g (casein diet). The FER of rats fed with casein and cerelac were  
 168 not significantly different (P=0.05) but were higher than those of rats fed with toasted crayfish enriched ogi  
 169 and oven-dried crayfish enriched ogi diets. The corrected PERs of the different diets varied from 2.3 (toasted  
 170 crayfish enriched ogi) to 2.5 (casein diet). The corrected PERs of the formulated diets were similar to those  
 171 of cerelac and casein diets. The PAG (Protein Advisory Group) and U.S. Department of Agriculture  
 172 guidelines recommend a PER of not less than 2.1 and preferably greater than 2.3 for weaning food and  
 173 corn-based blends [18,26,27]. The net protein ratio (NPR) of the diets ranged from 2.6 (toasted enriched ogi)  
 174 to 3.25 (casein diets). NPR is a more accurate measure of protein quality than PER as it allows for the  
 175 evaluation of maintenance requirement and results are independent of feed intake. The NPR of the  
 176 formulated diets was lower than those of casein and cerelac diets. Similar report was obtained by Fashakin,  
 177 [28].





Casein Cerelac  
oven-dried toasted



Casein Cerelac  
oven-dried toasted

179



180  
181

182 Fig. 2: (a) Feed Efficiency Ratio (FER), (b) Protein Efficiency Ratio (PER) and (c) Net Protein Ratio (NPR) of  
183 rats fed on different weaning diets and casein

184 The results of true digestibility (TD), biological value (BV), net protein utilization (NPU) and protein retention  
185 efficiency (PRE) are illustrated in (Fig 3). TD, BV, NPU and PRE ranged from 76.0 to 89.8%, 66.5 to 87.6%,  
186 50.5 to 77.4% and 42 to 51 respectively. The TD, BV and NPU of casein and cerelac were higher ( $p \leq 0.05$ )  
187 than those of oven-dried and toasted crayfish enriched ogi. The results obtained in the present study are  
188 similar to those observed by Obizoba [29], who reported BV values of 67.6 to 75.9% and NPU values of 51.8  
189 to 62.3% for the weaning food prepared from malted corn plus crayfish. The lower values of TD, BV and



190 NPU in the toasted diet may be due to roasting as it affects the availability of some amino acids. Similar  
191 report was obtained by Dahiya and Kapoor, [9] who showed that PER, TD, BV and NPU decreased on  
192 roasting. The effect of roasting on availability of amino acid can be minimized by roasting at a reduced  
193 temperature. Since protein retention efficiency (PRE) was obtained by multiplying NPR by 16, the trend of  
194 the result obtained for PRE is similar to that of NPR (Fig. 2.).







Fig. 3: (a) The True Digestibility (TD), (b) Biological Value (BV), (c) Net protein Utilization (NPU) and (d) Protein Retention Efficiency of rats fed with different weaning diets and casein

### 3.3 Organ weights and haematological parameters of animals fed with cerelac, casein, formulated diets and ordinary ogi

The weight of some vital organs of animals fed with cerelac, casein, formulated diets and ordinary ogi are shown in Table 4. The heart weight, the liver weight, the spleen weight and kidney weight ranged from 0.20 to 0.37g, 2.19 to 4.4g, 0.13 to 0.42 g and 0.35 to 0.78g respectively. The weights of the heart, the kidney, the spleen and liver of animals fed with oven dried and toasted crayfish enriched ogi compared favorably with those of standard diet (Casein and Cerelac). This indicates that the formulated diets may not result in abnormal development of the vital organs.

209 The results of haematological parameters of animals fed with cerelac, casein, formulated diets and ordinary  
 210 ogi are shown in Table 5. The blood indices varied: packed cell volume (PCV) 30.25 to 33.0 %, haemoglobin  
 211 concentration (Hb) 10.03 to 11.28%, red blood cell (RBC) 64.45 to 73.08 x10<sup>5</sup>, white blood cell (WBC) 52.0  
 212 to 101.5 x 10<sup>5</sup>, erythrocyte sedimentation rate 1.23 to 1.65, lymphocyte 50.18 to 53.15%, basophil 2.0 to  
 213 11.25%, neutrophil 30.5 to 37.35% and monocytes 7.75 to 10.0%. PCV measures the ratio of the volume  
 214 occupied by red blood cell to the volume of whole blood cell. It is a convenient and rapid measure of the  
 215 degree of anaemia [18]. Low PCV, Hb and serum protein have been associated with protein deficiency [25].  
 216 The PCV, Hb and RBC of rats fed with basal diet were lower than those fed with casein, cerelac and  
 217 formulated diets. Similar results were reported by Osundahunsi and Aworh [30]. The values obtained for  
 218 PCV, RBC, WBC and Hb of the rats fed with formulated diets were similar to those fed with casein and  
 219 cerelac. The results show the adequacy of the formulated diets in blood formation. This suggests that the  
 220 feeding of formulated diets will support haematopoietic activities of the body.

221 **Table 4: Organ weights (g) of rats fed with crayfish enriched-ogi, casein, cerelac and ordinary 'ogi'**

Dietary group	Heart (g)	liver (g)	Spleen (g)	Kidney (g)
<b>Cerelac</b>	0.28 <sup>b</sup>	4.02 <sup>ab</sup>	0.31 <sup>ab</sup>	0.72 <sup>ab</sup>
<b>Casein</b>	0.37 <sup>a</sup>	4.49 <sup>a</sup>	0.42 <sup>a</sup>	0.78 <sup>a</sup>
<b>Oven-dried</b>	0.29 <sup>b</sup>	3.79 <sup>b</sup>	0.29 <sup>b</sup>	0.55 <sup>ab</sup>
<b>Enriched 'ogi'</b>				
<b>Toasted</b>	0.31 <sup>ab</sup>	3.80 <sup>b</sup>	0.29 <sup>b</sup>	0.53 <sup>ab</sup>
<b>Enriched 'ogi'</b>				
<b>Ordinary'Ogi'</b>	0.20 <sup>c</sup>	2.19 <sup>c</sup>	0.13 <sup>c</sup>	0.35 <sup>b</sup>
<b>diet</b>				

222

223 Values with different superscript on the same column are significantly different (p=0.5)

224

225 **Table 5: Haematological parameters of rats fed with formulated diets, casein and cerelac**

Parameters	Toasted	Oven-dried	Casein	Cerelac	Ogi
	enriched ogi	enriched ogi			
<b>Packed cell volume</b>	30.50	30.75	32.00	33.00	30.25
<b>(%)</b>					

<b>Haemaglobin</b>	10.38	10.25	10.73	11.28	10.03
<b>(g/100ml)</b>					
<b>Red blood cell (x 10<sup>5</sup>)</b>	67.55	67.30	70.98	73.08	64.45
<b>White blood cell</b>	76.63 <sup>b</sup>	52.01 <sup>c</sup>	66.51 <sup>bc</sup>	68.01 <sup>bc</sup>	101.5 <sup>a</sup>
<b>(x10<sup>2</sup>)</b>					
<b>Erythrocyte</b>	1.23 <sup>b</sup>	1.35 <sup>ab</sup>	1.38 <sup>ab</sup>	1.35 <sup>ab</sup>	1.65 <sup>a</sup>
<b>sedimentation rate</b>					
<b>Lymphocytes (%)</b>	53.15	51.00	50.18	51.00	54.25
<b>Monocytes (%)</b>	9.00	8.00	7.75	9.00	10.00
<b>Eosinophil (%)</b>	2.50	2.75	2.00	2.00	3.50
<b>Basophil (%)</b>	1.25 <sup>ab</sup>	1.75 <sup>a</sup>	1.25 <sup>ab</sup>	1.50 <sup>a</sup>	2.01 <sup>a</sup>
<b>Neutrophil (%)</b>	36.75 <sup>a</sup>	30.51 <sup>b</sup>	37.25 <sup>a</sup>	36.01 <sup>a</sup>	31.25 <sup>b</sup>

226

## 227 Conclusion

228 The study showed that the formulated diets promote growth better than ordinary ogi. The haematological  
 229 indices and organ weight measurement of the rats fed the formulated diets were better than that of ordinary  
 230 ogi and compared favourably with that of rats fed with standard casein and **cerelac**. The study indicated that  
 231 oven dried enriched ogi and toasted enriched ogi may support growth in children than ordinary ogi which is  
 232 currently in use as traditional weaning foods in Nigeria. The implications of these findings are far reaching  
 233 since all the components used in the formulation are obtained from local market and toasting is a processing  
 234 method that can easily be practiced at home. Adoption of toasted enriched ogi may make the product a  
 235 potentially more functional and more accessible weaning food.

236

237 **COMPETING INTERESTS**

238 We declared no competing interests exist

239

240 **ETHICAL APPROVAL**

241 This study was approved by the ethical review committee of the Federal University of  
242 Technology, Akure, Ondo State, Nigeria

243

244 **REFERENCES**

- 245 1. Fetuga GO (2001) Nutritional Evaluation of maize-soyabens based weaning food  
246 fortified with some local raw food materials. Proceedings of 15th Annual NIFST  
247 conference. November, 5-9 . (Lagos).
- 248 2. Fashakin J B, Ogunsola F (1982) The utilization of local foods in the formulation of  
249 weaning foods. Journal of Tropical Pediatrics 28:93-96.
- 250 3. Mila'n -Carrillo J, Valde'z-Alarco'n C, Gutie'rrez-Dorado R, Ca'rdenas-Valenzuela  
251 OG, Mora-Escobedo R, Garzo'nN-Tiznado JA, & Reyes-Moreno C (2007) Nutritional  
252 properties of quality protein Maize and chickpea extruded based weaning food. Plant  
253 foods for Human Nutrition. 62:31-37
- 254 4. Ongol MP Niyonzima E, Gisanura I, Vasanthakaalam H (2013) Effect of  
255 germination and fermentation on nutrients in maize flour. Pak J Food Sci 23:183-  
256 188
- 257 5. Gabriel RAO, Akharaiyi FC (2007) Effect of spontaneous fermentation on the  
258 chemical composition of thermally treated jack beans (*Canavalia ensiformis* L.). Int J  
259 Biol Chem 1:91-97.
- 260 6. Ijarotimi OS (2012) Influence of germination and fermentation on chemical  
261 composition, protein quality and physical properties of wheat flour (*Triticum*  
262 *aestivum*). Journal of Cereals and Oil seeds 3:35-47.
- 263 7. Sangronis E, Machado CJ (2007) Influence of germination on nutritional  
264 quality of *Phaseous vulgaris* and *Cajanus cajan*. J Sci Tech 40:116-120.
- 265 8. Obizoba IC, Egbuna HI (1992) Effect of germination and fermentation on the  
266 nutritional quality of bambara groundnut (*Vaandzenia subterranea* L. Thauars) and  
267 its product (milk). Plant Foods for Human Nutrition 42: 13-23.
- 268 9. Dahiya S, Kapoor AC (1993). Nutritional evaluation of home processed weaning  
269 foods based on low cost locally available foods. Journal of Food Chemistry 48: 179-  
270 182
- 271 10. Hotz C, Gibson RS (2007) Traditional food-processing and preparation practices to  
272 enhance the bioavailability of micronutrients in plant-based diets. J. Nutr. 137:  
273 1097–1100.
- 274 11. Hassan EE, Babiker AH, Tinay EI (2007) Content of antinutritional factors and HCl-  
275 extractability of minerals from white bean (*Phaseolus vulgaris*) cultivars: Influence of  
276 soaking and/or cooking. Food Chem100: 362-368
- 277 12. FAO/WHO/UNU (1985, 1998, 2002) Preparation and use of food-based dietary  
278 guidelines. Report of a Joint.FAO/WHO Consultation. WHO Technical Report series  
279 880. Geneva.
- 280 13. Ibironke SI, Fashakin JB, Badmus OA (2012) "Nutritional evaluation of  
281 complementary food developed from plant and animal protein sources." Nutrition &  
282 Food Science 42: 111-120.
- 283 14. Zaglol NF, Eltadawy F (2009) Study on chemical quality and nutrition value of fresh  
284 water cray fish (*Procambarus clarkii*). Journal of the Arabian Aquaculture Society 4:  
285 1-18.

- 286 15. Ibironke SI, Fashakin J B, IGE MM (2014) Nutritional quality of animal polypeptide  
287 (Crayfish) formulated into complementary foods. American Journal of Food and  
288 Nutrition 2: 39-42.
- 289 16. Ikujenlola VA & Fashakin JB (2005). The physicochemical properties of a  
290 complementary diet prepared from vegetable proteins. Journal of Food Agriculture  
291 and Environment 3:23-26.
- 292 17. Ijarotimi OS (2006). Evaluation of nutritional quality and sensory attributes of home-  
293 made processed complementary diet from locally available food materials (*Sorghum*  
294 *bicolor* and *Sphenostylis stenocarpa*). Journal of Food Technology 4:334-338.
- 295 18. Abiose SH, Ikujenlola AV, Abioderin FI (2015). Nutritional Quality Assessment of  
296 Complementary Foods Produced from Fermented and Malted Quality Protein Maize  
297 Fortified with Soybean Flour. Pol. J. Food Nutr. Sci. 65 (1): 49–56.
- 298 19. Ijarotimi OS (2008). Protein and haematological evaluations of infant formulated  
299 from cooking banana fruits (*Musa* spp, ABB genome) and fermented bambara  
300 groundnut (*Vigna subterranean* L. Verdc) seeds. Nutrition Research and Practice.  
301 2(3):165-170
- 302 20. Olutiola PO, Famurewa O, Sontag HG(1991) An Introduction to General  
303 Microbiology. Heidelberger Veringsinstall and Druekeriegmbll Heidelberg  
304 Germanybll Heidelberg Germany.
- 305 21. AOAC (1995). Association of official analytical Chemists Official Methods of Analysis  
306 16th ed. Washington D.C.
- 307 22. Agbede JO, Aletor VA (2003) Comparative evaluation of weaning foods from  
308 *Glyricidia* and *Leucaena* leaf protein concentrates and some commercial brands in  
309 Nigeria. Journal of the Science of Food and Agriculture 84: 21- 30.
- 310 23. Lamb G.N (1981) Mammal of Veterinary Laboratory Technique. CIBA-GEIGY.  
311 Kenya Pp96-97.
- 312 24. PAG (1995) PAG Ad Hoc working group meeting on clinical evaluation and  
313 acceptable nucleic acid-levels of SOP for human composition. Geneva
- 314 25. Ibironke SI (2014) Formulation of Infant Weaning Foods from Vegetable Proteins  
315 and Cereal. American Journal of Food Technology 9: 104-110.
- 316 26. PAG, Protein Advisory Group of the United Nations. Guideline no. 8: Protein rich  
317 mixtures for use as weaning foods. New York: Food and Agriculture Organization of  
318 the United Nations/World Health Organization/United Nations Children's Funds,  
319 1971, pp. 1–7.
- 320 27. Annan NT, Plahar WA (195). Development and quality evaluation of a soy fortified  
321 Ghanaian weaning food. Food Nutr. Bull. 16(3):263–267.
- 322 28. Fashakin JB (2006). Bioassay and Physico-chemical properties of complementary  
323 food prepared from cowpea (*Vigna unguiculata*) and melon seeds as man source of  
324 proteins Proceeding of Annual Conf. School of Agric and Agric Technology. Federal  
325 University of Technology Akure, Nigeria, 24th May 2006.
- 326 29. Obizoba IC (1988) Nutritive value of malted, dry or wet milled sorghum and corn.  
327 cereal Chem 65: 447-449.
- 328 30. Osundahunsi OF, Aworh OC (2003) Nutritional evaluation with emphasis on protein  
329 quality of maize – based complementary food enriched with soybean and cowpea  
330 tempe. International Journal of Food Science and Technology 38: 809 – 813.