Review Article

Phosphate Fertilizer Sources and Use in Africa: Challenges and Opportunities

4

1

5 ABSTRACT

Most African soils are low in phosphorus (P) and farmers are increasingly using more P fertilizers to improve crop production. Phosphate rocks (PR) deposits which are the major source of P are unfortunately finite. This is likely to cause world food crisis, especially in Africa due to its high human population growth rate. Conflicting information on the quantity of existing PR reserves has made it difficult to accurately predict how low they would last so as to plan for its judicious use. World PR deposits are currently estimated at 290 billion tonnes and potentially 490 billion tonnes. These reserves at the current production rate of 160 - 170 million tonnes per year are likely to be depleted between years 2311 and 2411 and Africa will be most affected due to high human population grow rate. With the eminent threat of PR deposits depletion, African therefore needs to adopt farming practices that will reduce the use of P fertilizers without negatively affecting its crop productivity. Practices such as soil erosion control, use of P efficient crop germplasms, P solubilizing organisms and organic materials are perceived to reduce soil P loss and increase its use efficiency by plants.

6

7 Key words: Soil fertility, phosphorus, phosphate rocks, depletion, reserves,

8 1. INTRODUCTION

9 1.1 Importance of Phosphorus in Plant Nutrition

10 Phosphorus (P) is an essential plant nutrient and as a result crop response to P fertilizer applications 11 are widespread. Total P in plant tissue ranges from about 0.1 to 1% and it plays both metabolic and 12 structural roles in plants [1]. Metabolic roles include: photosynthesis, synthesis and breakdown of carbohydrates and energy transfer processes within the plant. The energy obtained during 13 14 photosynthesis and carbohydrate metabolism is stored in energy rich phosphates compounds namely; 15 adenosine diphosphate (ADP) and adenosine triphosphate (ATP). Phosphorus is a structural component of nucleic acids, coenzymes, nucleotides, phosphoproteins, phospholipids and sugar 16 17 phosphates [2]. Large amounts of P are deposited in reproductive cells; therefore, it is essential for 18 seed and fruit formation, faster grain maturity, quality and strong cereal straws. Phosphorus is also 19 important for good root development and growth [2, 3]. Common P deficiency symptoms include 20 purple or bronze colouration appearing on lower leaf tips, progressing along the leaf margins until the

21 entire leaf is discoloured. Since P is mobile within the plant, its deficiency symptoms are first

- 22 expressed on lower leaves [3]. Phosphorus deficiency in soils therefore interferes with photosynthesis,
- 23 protein synthesis, respiration and biomass production in plants.
- Crop responses to application of various P sources in Africa soils are enormous. Inorganic P fertilizers
 have increased soil P levels and crop productivity in many African countries [8].

26 1.2 The Process of Soil Phosphorus Depletion in Africa

27 The major contributing factors to soil fertility depletion in Africa are breakdown in traditional 28 practices and low priority given to the rural areas. Increasing pressure on land due to high human 29 population has led to breakdown in traditional farming systems whereby fallowing, cereal-legume 30 intercropping, mixed crop-livestock farming and opening of new lands maintained soil fertility [4]. 31 Limited attention is given by African governments to rural areas where farming is carried out. As a 32 result most smallholder farmers, who produce about 90% of its food, lack the credit to purchase 33 fertilizers to replenish soil fertility. In 30 years (i.e. from the year 1967 – 1997), about 75 kg P/ha was 34 lost from about 200 million cultivated land in 37 African countries [5]. The continent is now loosing 35 0.5 million tons of P every year from its cultivated lands which is much higher than its annual 36 consumption of 0.26 million tons P [6].

Nearly three-quarters of farmlands in Africa are nutrient depleted, lowering crop yield to one-quarter of the global average [7]. At the same time, more nutrients continue to be removed each year than are added in the form of fertilizer, crop residues and manure. Nutrient balance studies in the 1990s suggested average annual P depletion of 2.5 kg P/ha [5]. Intensively cultivated highlands in East Africa loose an estimated 5 kg P/ha year, whereas croplands in the Sahel loose 2 kg P/ha [8]. Therefore, most African soils have low levels of soil available P to support high crop production required for its already high and increasing human population.

44 1.3 Sources of Phosphorus for Soil Fertility Replenishment

Phosphorus is the second most limiting nutrient to crop production after nitrogen (N) in many tropical soils including Africa [9]. While soil N can be replenished through biological nitrogen fixation from atmospheric sources, P sources are not renewable through such biological means [5, 10]. Therefore, soil P replenishment is mainly through inorganic fertilizer sources from phosphate rocks (PR) with minor sources from manures, guano and human excreta [11]. The main source of P fertilizer is finite and this poses a great danger to world food production, especially in African due to its high human population growth rate compared to other parts of the world.

52 1.4 Phosphorus Fertilizer Use in Africa Compared to Other Parts of the World

53 Farmers in Africa are becoming aware of the importance of using fertilizer to increase crop 54 production [12]. As a result the demand for fertilizers such as those containing phosphorus is on the 55 increase. Between the years 1950 and 2000, global use of fertilizers that contain P, N and K increased 56 by 600% [13]. The increase is linked to soil fertility depletion. Average annual fertilizer use in Africa 57 is only about 17 kg per ha, compared, for example to 85 kg/ha in North America, 96 kg/ha in Latin 58 America and 196 kg/ha in Asia. Even this low rate of consumption is restricted to just a few African 59 countries. Sub-Saharan Africa, excluding South Africa, uses about 5 kg per ha per year of fertilizer, of 60 which less than 30 per cent is phosphorus [14]. With this background, it is apparent that on average 61 Africa uses about 5.1 kg P fertilizer per ha/year. The continent looses between 2 and 5 kg P/ha 62 through crop harvest from its farmlands [8]. On African's high potential areas with acid soils crops 63 recover only 10 - 25 % of the applied P fertilizers [15, 16]. Crops therefore recovers only about 0.5 - 10064 1.3 kg P per ha of the applied P fertilizers. As a result, the rates of P fertilizer applied by African 65 farmers are insufficient to balance off the amounts taken up by crops as well as those fixed in acid 66 soils. A combination of high cost and low accessibility to credit prevents many African farmers from 67 acquiring fertilizers. Poor transport, low trade volumes and lack of local production or distribution 68 capacity resulting in farm-gate fertilizer prices two to six times higher than the world average. 69 Nevertheless, fertilizer is needed to achieve adequate and sustainable crop yields. The Africa 70 Fertilizer Summit [17] concluded that a lasting solution requires policies to sustain robust distribution 71 networks; including adequate credit sources, retail outlets and transportation, as well as the transfer of 72 technology and knowledge for efficient fertilizer use.

73

74 2. WORLD PHOSPHATE RESERVES

75 It is not easy to ascertain the quantity of world phosphate reserves. Knowledge of phosphate rock 76 deposits is evolving, along with technology and the economics of production [18]. Compared to fossil 77 fuel, most deposits of PR are found in very few countries. Most reserves are found in Morocco, the 78 USA and China (Table 1). The reserves are estimated to be about 16 billion tons [19]. These reports 79 also suggest that estimates are not comprehensive, as they do not include deposits in all countries. A 80 recent report from the International Fertilizer Development Center (IFDC) on reserves and resources 81 provisionally revised the estimate of phosphate rock reserves from the United States Geological 82 Survey (USGS) estimate of around 16 to 60 billion tonnes [19], which is somehow consistent with the 83 most recent USGS report [20].

The IFDC report estimates world's phosphate reserves to be approximately 290 billion tonnes and potentially as much as 490 billion tonnes [19]. It seems the world phosphate reserves are underestimated given the fact that they are continually being revised upwards as more reserves are discovered. At the same time the deposits with small quantities in Africa such as Minjingu in 88 Tanzania, Busumbu and Sukulu in Uganda among others are not listed [21]. Since not all the 89 reserves are quantified, it's likely that the deposits are still under-estimated even with these recent 90 upward predictions. Therefore, there is need to accurately estimate the quantity of all the deposits for 91 proper planning for use of this vital resource.



92 Table 1. World phosphate reserves

94

95 3. LIFESPAN OF WORLD PHOSPHATE ROCK RESERVES

96 There is conflicting information on how long PR reserves would last. However, how long they will 97 last is dependent on their quantity, quality and rate of mining. Knowledge of phosphate rock deposits 98 is evolving, along with technology and the economics of production [18]. It is predicted that peak 99 phosphorus production will occur between the years 2030 and 2040 [11]. The estimate was based on 100 USGS data for global phosphate reserves [20]. It is suggested that world PR reserves depletion 101 would occur in 50-100 years [22] meaning that it will be between the years 2058 and 2108. 102 However, recent upward estimates of the extent of the PR reserves to about 60 - 160 billion metric 103 tonnes have pushed upward when the reserves would be depleted [19, 20]. A report by IFDC 104 indicates that there are sufficient PR reserves to produce P fertilizers for the next 300-400 years at 105 current production rates of 160 to 170 million tonnes per year. This prediction therefore indicates 106 that the PR deposits would be depleted between the years 2311 and 2411. To cater for the increasing 107 demand for P fertilizer due to population increase, the rate of PR mining is likely to increase making 108 the reserves to be depleted sooner than later. This is likely to pose food crisis in Africa given its high 109 population growth rate compared to other parts of the world. This date of depletion is with an 110 assumption that the rate of mining and consumption will remain the same, which is unlikely 111 scenario. Small PR reserves are not included and more discoveries are expected to be reported in 112 future, therefore it is likely that the depletion would take longer than expected. A number of deposits 113 particularly in Africa are not economically viable since they have very low P content [21] and 114 therefore not worth exploiting.

4. OPPORTUNITIES FOR AFRICA TO IMPROVE PHOSPHATE FERTILIZER USE EFFICINCY

118 Rock phosphate sources are non-renewable, hence the need for Africa to adopt best practices to help 119 prolong the lifespan of existing PR deposits. Morocco has the highest world PR reserve, however, it 120 has export restrictions to preserve its deposits [19]. A part from Morocco and Egypt, the rest of 121 African countries import phosphate fertilizers. This therefore calls for African as a continent to put in 122 place practices that will minimize P fertilizer losses as well as improving it use efficiency by plants. 123 Such practices include soil erosion control, use of P efficient crop germplasms, P solubilizing 124 organisms and organic materials among others as discussed below. The practices are likely to reduce 125 the amount of P fertilizer required by the continent.

126 4.1 Soil Erosion Control

127 Most plant nutrients are found in the topsoil and therefore removal of topsoil through erosion reduces 128 soil fertility. Protecting the topsoil from soil erosion therefore minimizes nutrient losses. Run-off and 129 erosion combined are responsible for 48 and 40 per cent of phosphorus losses in intensively cultivated 130 highland areas and in parts of the Sahel, respectively [8]. Africa looses about 0.47 tons per haper year 131 of its topsoil [23]. Soil erosion accounts for about 75-90% soil P losses in Africa estimated at 1.0 kg P 132 loss ha per year [8, 25]. A number of soil erosion control techniques exist. Ploughing across rather 133 down the slope and planting of hedgerows on steep lands greatly reduce soil erosion. Soil vegetation 134 cover is one of the best ways to control soil and nutrient losses due to erosion. African farmers need to 135 use mulches, cover crops and fertility enhancing systems on low-fertility soil to minimize soil erosion 136 losses [26].

Given the extent of soil fertility losses through erosion, there is need for African countries to put in place measures to curb these losses. Farmer education on soil erosion control and creation of awareness on the available control measures are important. African countries also need to formulate and put in place policies on soil erosion control measures. These will help maintain high soil fertility levels through minimization of nutrient losses such as P.

142

143 **4.2 Use P Efficient Crop Germplasms**

Most African soils are inherently low in soil available P. This is exacerbated by the fact that a vast majority of this P is not readily available to plants. Traditional systems of farming thus, unknowingly, relied on growing crop species with low P requirements [27]. Large proportion of P in African soils is unavailable for plants uptake due to its fixation particularly in high to medium agricultural areas with

115

acid soils [5]. In modern agriculture, continuous use of P fertilizers over many years has increased the
total P levels in the soils but the available P remains low [28]. The unavailable P can represent a
reserve which can be exploited by crops that are well adapted to extraction of P from less available
soil fractions [29].

152 Many trees, shrubs and important crop species grown in Africa have the ability to exude organic acids 153 from their roots or have mycorrhizal associations that help dissolve inorganic P that are not otherwise 154 available to plants [5]. Other P acquisition strategies that are used by adapted species include 155 excretion of phosphatases to release the organically bound P and provision of extra carbon as a 156 booster of microorganisms, which in turn, also produce organic acids as well as phosphatase [30]. 157 There is, therefore, a campaign in some quarters to tailor plants to fit the soil through genetic 158 improvement in the belief that it is more economical than changing the soil. There may be reasonably good prospects for improving the efficiency of P use by plants by selecting appropriate genotypes 159 160 with characteristics for root hair length, organic acids production in the rhizosphere, and mycorrhizal 161 associations for soils with low P [31]. It has been reported that some of the genotypes express a 162 protein kinase gene called phosphorous starvation tolerance gene (Pstol1), which enables acquisition 163 of P and other nutrients [32] even in P deficient soils.

164

165 To deal with low soil available P related problems, plant breeding programs have developed 166 germplasms tolerant to low soil P [33, 34]. Studies in Africa have reported maize and sorghum 167 germplasms that are P use efficient [35-37]. These elite materials provide a good foundation for 168 breeding for P use efficiency. Currently, there are no commercial maize, sorghum or other crop 169 varieties available to farmers that are adapted to low P soils [35]. There is need therefore, to develop 170 crop varieties for Africa that are P use efficient to enhanced crop productivity on low P soils. Use of 171 P efficient crop germplasms will make both the native and the applied P fertilizer normally fixed on 172 medium to high potential areas with acid soils available for plants uptake [3, 5]. It is important to note 173 this management option is not sustainable without application of P fertilizers, because the removal of 174 P in the harvested produce will eventually lead to a decline in total soil P levels.

175

176 4.3 Use of Phosphorus Solubilizing Organisms

Almost 75–90% of added phosphatic fertilizer is precipitated by metal cation complexes present in soils [38] and has led to accumulation of P in forms that are not available for plants uptake. Further, it has been suggested that the accumulated phosphates in agricultural soils are sufficient to sustain high crop yields worldwide for about 100 years [39]. Thus, the dependence of fertilizer production on a fossil energy source and the prospects of the diminishing availability of costly input of fertilizer production in years to come have obviously brought the subject of mineral phosphate solubilization to the forefront [40]. P-solubilizing activities in agricultural soils are considered to be environmentally 184 friendly alternative to further applications of mineral based P fertilizers [41]. Under diverse soil and 185 agro-climatic conditions, the organisms with phosphate-solubilizing abilities have proved to be an 186 economically sound alternative to the more expensive superphosphates and possess a greater 187 agronomic utility [40]. Use of phosphate solubilizing bacteria as inoculants increases soil P 188 availability, P uptake by plants that reduces external P-fertilizer application rates, reduces 189 environmental pollution and promotes sustainable agriculture. The introduction of mycorrhizae into 190 soils has also been suggested for improving the availability of soil P, but initial enthusiasm for these 191 has waned [5]. Mycorrhizae are important for many plant species when grown in P-deficient soils, but 192 much less effective where soil P status is adequate. Enhancing the availability of soil fixed P through 193 use of P solubilizing organisms is one way farmers in Africa can reduce the use of fertilizers, thus 194 prolonging the lifespan existing PR reserves.

195

196 4.4 Use of Organic Materials

197 With increasing costs of the inorganic fertilizers and the finite nature of PR, it is imperative to explore 198 alternative phosphate sources. Before the advent of inorganic P fertilizers, crop production relied on 199 native soil P and the addition of locally available organic matter, mainly animal manures [11]. The 200 unavailable P could be made available to other subsequent crops after decomposition of the residues 201 of P use efficient plants [42]. Organic P sources vary widely in terms of P concentration, chemical 202 form and state (solid, liquid or sludge). They include animal manures, composts, crop residues, green 203 manures and human excreta. In most cases however, their P content is often too low to meet the crop 204 nutrient demands [43]. Organic materials (OMs) can improve plant P use efficiency of both the 205 native soil P and applied P fertilizers; therefore reducing on the need for fertilizer P inputs [44]. Use 206 of OMs for soil fertility management in Africa face challenges such as inadequate amounts available 207 to farmers and their low nutrient levels. Their low nutrient contents requires application of large 208 volumes which increases the labour costs that cannot be offset by the crop yields obtained [44, 46]. 209 Therefore, OMs suitable for use as P sources should have a high P content and low cost of production 210 to make them economically viable to farmers [47]. A part from using from high quality OMs, the 211 quality of organic materials can be enhanced through pit storage and manure storage under shade [48]

A lot work in Africa on use of OMs has not focused on use of human wastes. Human beings produce large amounts of excreta (faeces and urine), that can provide adequate amount of organic materials for soil fertility management. Human urine has been reported to contain P, N and potassium (K) in the correct ratios, necessary for plant nutrition [49]. Studies in African countries such as Zimbabwe have revealed that nutrients content in one person's urine are adequate to produce 50-100% of the food requirement for another person [49]. Challenges on human excreta handling has been resolved by development of guidelines that minimize health risks [50]. Many Africans consider food produced from human excreta unfit for human consumption. As a result there is need to create awareness among Africans that food produced using human excreta is safe if the guidelines are properly followed.

222 **5.** CONCLUSION

223 Most African soils have low soil available P and crop responses to its applications are widespread. 224 Use of fertilizers such as P in African as a continent to feed in its human population is on the increase. 225 Future use of P fertilizer faces a major change since its major source, the PR deposits are finite. This 226 is likely to cause food crisis in Africa due to its high population growth rate. It has not been easy to 227 accurately quantify PR reserves, thus making it difficult to predict how long they will last so as to 228 plan for its judicious use. However, with estimates of 290 billion tonnes and a potential of 490 billion 229 tonnes reserves at the current production rate of 160 - 170 million tonnes per year they are likely to be 230 depleted between years 2311 and 2411. With eminent PR reserves depletion, Africa needs to adopt 231 farming practices that will minimize P losses as well as improving its availability to plants to help 232 reduce use of inorganic P fertilizers. Soil erosion control is one of such measures that should be 233 adopted to minimize top soil P losses. Large amounts of soil P in Africa soils particularly in acid soils 234 are fixed. Use of P use efficient crop germplasms and P solubilizing organisms are some of the 235 practices that can help African farmers exploit fixed soil P thereby reducing external P fertilizer use. 236 Unfortunately there are no improved P use efficient crop germplasms in Africa, therefore the need to 237 develop such germplasms. Another practice that can help reduce use of inorganic P fertilizers is 238 application of organic materials since it increases soil P and improves its availability in soils. Use of 239 organic materials in Africa faces challenges such as low quantities available to farmers and low 240 nutrient contents. Human urine as organic source should be exploited since it has right nutrient 241 contents necessary for healthy plant growth.

242

243 **REFERENCES**

- 1. Tisdale SL, Nelson WL, Beaton JD. Soil fertility and fertilizers. 5th edition. New York:
 Macmillan;1990.
- Sanginga N, Woomer PL., editors. Integrated Soil fertility management in Africa:
 principles, practices and development processes. Tropical Soil Biology and Fertility
 Institute of the International Centre for Tropical Agriculture, Nairobi, Kenya; 2006.
- 3. Havlin JL, Tisdale SL, Beaton JD, Nelson WL. Soil fertility and fertilizers. Upper Saddle
 River (NJ); 2005.

8

- 4. Sanders JH, Shapiro JI, Ramaswamy S. The economics of agricultural technology in semiarid sub-Saharan Africa. John Hopkins University Press Baltimore MD; 1996.
- 5. Sanchez PA, Shepherd KD, Soule MJ, Place FM, Buresh PJ, Izac A-MN, et al. Soil
- 254 fertility replenishment in Africa: An investment in natural resource capital. In: Buresh RJ,
- et al. editors. Replenishing soil fertility in Africa. Soil Science Society of America, SSSA
- 256 Madison Wisconsin, USA: Special Publication No.51; 1997.
- 6. FAO. FAO fertilizer year book 1994. Vol. 4. Food and Agricultural Organization of the
 United Nations; 1995.
- 7. Henao J, Baanante C. Agricultural production and soil nutrient mining in Africa:
 Implications for resource conservation and policy development. International Fertilizer
 Development Center (IFDC), Muscle Shoals, Alabama, USA; 2006.
- BH. 262 8. Smaling, EMA, Nandwa SM, Janssen Soil fertility in Africa 263 is at stake. In: Buresh RJ, et al. editors. Replenishing soil fertility in Africa. Soil Science Society of America, SSSA Madison Wisconsin, USA: Special Publication No.51; 1997. 264
- 9. Bekunda MA, Bationo A and Sali H. Soil fertility management in Africa. A review of
 selected research trials. In: Buresh RJ, et al. editors. Replenishing soil fertility in Africa.
 Soil Science Society of America, SSSA Madison Wisconsin, USA: Special Publication
 No.51; 1997.
- 10. Okalebo JR, Othieno CO, Woomer PL, Karanja NK, Sesmoka JRM, Bekunda MA, et al.
 Available technologies to replenish soil fertility in East Africa. Nutrient Cycling in
 Agroecosystems. 2006; (76):153-170.
- 272 11. Cordell D, Drangert J.-O and White S. The Story of Phosphorus: Global food security and
 273 food for thought. Global Environmental Change. 2009; (19): 292-305
- 12. Okalebo JR, Simpson JR, Okwach EG, Probert ME, McCrown LR. Conservation of soil
 fertility under intensive maize cropping in semi-arid eastern Kenya. Proceedings of the
- Third African Crop Science Conference, 1 October 1997, Pretoria, South Africa; 1997.
- 13. International Fertilizer Industry Association. Production and International Trade Statistics.
 International Fertilizer Industry Association, Paris; 2006
- 14. International Fertilizer Industry Association. Annual Phosphate Rock Statistics. International
 Fertilizer Industry Association, Paris; 2009
- 281 15. Bhal GS, Singh NT. Phosphorus diffusion in soils in relation to some edaphic factors and
- its influence on P uptake by maize and wheat. Journal Agricultural Science Cambridge.
- 283 1986; (107): 335–341.

- 16. Kisinyo PO. Constraints of soil acidity and nutrient depletion on maize (*Zea mays L.*)
 production in Kenya. Ph. D Thesis, Moi University, Kenya; 2011
- 17. Africa Fertilizer Summit. Africa Fertilizer Summit Proceedings. International Fertilizer
 Development Center (IFDC), Muscle Shoals, Alabama, USA; 2006
- 18. International Fertilizer Development Center (IFDC)/United Nations Industrial Development
 Organization (UNIDO), Fertilizer Manual. Prepared by the International Fertilizer
 Development Center (IFDC) and the United Nations Industrial Development
 Organization (UNIDO). Kluwer Academic Publishers, Dordrecht, the Netherlands; 1998.
- 292 19. van Kauwenbergh S. World Phosphate Rock Reserves and Resources, IFDC
 293 Technical Bulletin 75. International Fertilizer Development Center (IFDC), Muscle
 294 Shoals, Alabama, USA; 2010
- 20. Jasinski SM. Phosphate Rock. In: Mineral Commodity Summaries 2011. United States
 Geological Survey. United States Government Printing Office, Washington, D.C; 2011.
- 297 21. Jama B, van Straaten. Potential of East African phosphate rock deposits in integrated
 298 nutrient management strategies. Annals of Brazilian Academy of Sciences. 2006; (78):
 299 781–790.
- 22. Vaccari DA. Phosphorus Famine: The Threat to Our Food Supply. Scientific
 American, June 1; 2009; 54-59.
- 23. Cordell D. The story of phosphorus: missing global governance of a critical resource.
 Paper prepared for SENSE Earth Systems Governance, Amsterdam, 24-31 August, 2008.
 2008a.
- 24. El-Swaify SA, Dangler EW, Armstrong CL. Soil erosion by water in the tropics.
 Research Extension Series 024, College of Tropical Agriculture and Human Resources,
 University of Hawaii; 1982.
- 308 25. Sharpley A, Rekolainen S. phosphorus in agriculture and its environmental implications.
 309 In: Tunney H, et al. editors. Phosphorus loss from soil to water. CAB International,
 310 Wallingford, UK; 1997.
- 26. Stocking M. Rates of erosion and sediment yield in the African environment. In: Walling
 DE et al. editors. Challenges in African Hydrology and Water Resources (Proceedings of
 the Harare Workshop Symposium in July 1984). International Association of
 Hydrological Sciences (IAHS) Publication No. 14; 1984
- 27. Mokwunye AU, Chien SH, Rhodes E. Phosphorus reactions in tropical African soils. In:
- Mokwunye AU, Vex PLG, editors. Managing phosphorus and nitrogen fertilizers in sub-
- 317 Saharan Africa. Martinus Nijhof Publishers. Dordretch, The Netherlands; 1986.

- 28. Li M, Hou YL, Zhu B. Phosphorus sorption-desorption by purple soils of China in
 relation to soil properties. Australian Journal of Soil Research. 2007; (45): 182-189.
- 29. Buresh RJ, Smithson PC, Hellums DT. Building soil phosphorus capital in Africa. In:
- Buresh RJ, et al. editors. Replenishing soil fertility in Africa. Soil Science Society of
 America, SSSA Special Publication No.51. Madison Wisconsin, USA; 1997.
- 30. Droulin M, Merckx R. The role of citric acid as a phosphorus mobilization mechanism in
 highly phosphorus fixing soils. Gayana Botany. 2003; (60): 55-62.
- 31. FAO. Efficiency of soil and fertilizer phosphorus use: Reconciling changing concepts of
 soil phosphorus behaviour with agronomic information. Fertilizer and Plant Nutrition
 Bulletin no. 18. Rome, Italy; 2006
- 328 32. Gamuyao R, Chin JH, Pariasca-Tanaka J, Pesaresi P, Catausan S, Dalid C, et al. The
 protein kinase PSTOL1 from traditional rice confers tolerance of phosphorous deficiency.
 Nature. 2012; (488): 535-338.
- 33. Parentoni SA, Alves VMC, Gamma EEG, Colelho AM, Guimariaes CT, Guimariaes,
 PEO, et al. Breeding maize for Al tolerance and P us se efficiency and acid soil
 adaptation for the cerrado areas of Basil: the Embrapa's experience, In: Alves et al.
 editors. 3rd International Symposium on Phosphorus Dynamics in Soil-Plant continuum:
 Integrating marginal lands into productive agricultural Systems by means of improving
 soil and fertilizer phosphorus efficiency. 14 19th May, 2006. Uberlandia, Minas Gerais,
 Brazil; 2006.
- 338 34. Donswell CR, Paliwal RL, Cantrell RP. Maize in Third World. West View Press. Inc.
 339 1996
- 340 35. Ouma E, Ligeyo D, Matonyei T, Were, Agalo J, Too E, et al. Development of maize
 single cross hybrids for tolerance to low phosphorus. African Journal of Plant Science.
 2012 (14): 394-402.
- 36. Ligeyo DO. Genetic analysis of maize (*Zea mays* L.) tolerance to aluminium toxicity and
 low phosphorus stress and development of synthetics for use in acid soils of western
 Kenya. Ph.D. Thesis. Moi University, Eldoret, Kenya; 2007.
- 37. Too EJ. Physiological and molecular characterization of resistance to Aluminium stress in
 selected grain sorghum. PhD Thesis Moi University, Kenya; 2011
- 348 38. Stevenson FJ. Cycles of soil carbon, nitrogen, phosphorus, sulphur micronutrients, Wiley,
 349 New York; 1986

11

- 350 39. Goldstein AH, Rogers RD, Mead G. Mining by microbe. Biotechnology. 1993; (11):
 1250–1254.
- 40. Khan MS, Zaidi A, Wani PA. Role of phosphate-solubilizing microorganisms in
 sustainable agriculture A review. Agronomy for Sustainable Development, Springer
 Verlag. 2007; (27): 29-43.
- 41. Sharma, BS, Sayyed RZ, Trivedi MH, Gobi TA. Phosphate solubilizing microbes:
 sustainable approach for managing phosphorus deficiency in agricultural soils. Springer
 Plus 2013; (2): 587
- 42. Baligar VC, Fageria NK. Nutrient use efficiency in acid soils: Nutrient management and
 plant use efficiency. In: Moniz AC et al. editors. Plant soil interactions at low pH.
 Brazilian Soil Science Society Vicosa. 1997: 75-95.
- 43. Palm CA. Contribution of agroforestry trees to nutrient requirements of intercropped
 plants. Agroforestry Systems 1995; (30): 105-124.
- 44. Opala PA, Okalebo JR, Othieno CO, Kisinyo P. Effect of organic and inorganic
 phosphorus sources on maize yields in acid soils in western Kenya. Nutrient Cycling in
 Agroecosystems. 2010; (86):317–329.
- 45. Jama B, Swinkels RA, Buresh RJ. Agronomic and economic evaluation of organic and
 inorganic sources of phosphorus in western Kenya. Agronomy Journal. 1997; (89): 597604.
- 46. Opala PA, Jama BJ, Othieno CO, Okalebo JR . Effect of phosphate fertilizer application
 methods and nitrogen on maize yields in western Kenya. An agronomic and economic
 evaluation. Experimental. Agriculture. 2007; (43): 477-487.
- 47. Nziguheba G, Merckx R, Palm CA, Mutuo P. Combining *Tithonia diversifolia* and
 fertilizers for maize production in phosphorus deficient soil in Kenya. Agroforestry
 Systems. 2002; (55): 165-174.
- 48. Murwira HK, Nzuma JK, Manure management options for increasing crop production in
 communal areas of Zimbabwe. In: Delving JK, editors. Eco-agriculture initiatives in
 Eastern and Southern Africa. Weaver press, Avondale, Harare, Zimbabwe; 1999.
- 49. Cordell D. The story of phosphorus: 8 reasons why we need to rethink the management ofphosphorus resources in the global food system. Sustainable Phosphorus Futures website.
- 380 2008b. Accessed 27th January, 2016. <u>http://phosphorusfutures.net/why-phosphorus</u>.
- 50. EcoSanRes. Guidelines on the use of urine and faeces in crop production. EcoSanRes
 Factsheet 6. EcoSanRes Programme and the Stockholm Environment Institute. 2008.
 Accessed 22nd January, 2016. <u>http://www.ecosanres.org/pdf_files/ESR-factsheet-06.pdf</u>.