

Effect of cow dung slurry and Termite mount as seed treatment on germination and seedling characteristics of Red sanders (*Pterocarpus santalinus* L.f.)

Abstract

Aims:- *Pterocarpus santalinus* L.f. is a highly valued timber species, because of its “heavy, dark claret-red heartwood,” especially that possessing a ‘wavy’ grain. Propagated through mainly seeds, it faces a problem in seed germination because of hard seed coat, so limits poor seed germination and seedling production. current study was carried out to find out best germination for enhancement treatment.

Study design: —The data were then analyzed by the ‘F’ test for significance at 0.05 levels by using statistical software agrees with completely randomized block design.

Place and duration of study: - Forest college and research institute Mettupalayam, Tamil Nadu agricultural university, one year study.

Methodology: —Mature pods collected, were subjected to 4 treatments in 4 replications and the experiment was conducted in completely randomized block design.

The results: —The results showed that Cow dung slurry 24h and 72h resulted more synchronized germination of 51 percent, followed by Cow dung slurry 48h (44 %) as against 33% in control.

Conclusion: —Among all the treatment Cow dung slurry 24h resulted more synchronized germination of 51 percent.

Keywords: —Cow dung slurry, Termite mounts, Speed of germination, microbe.

1. Introduction

Red sanders distribution is largely confined to the southern portion of the Eastern Ghats in Andhra Pradesh, India (Shilpa *et al.*, 2012). The reddish and fragrant heartwood has range of medicinal, pharmaceutical industrial and timber value and thus economically placed in the same range as tusk and amber. The natural habitats of red sanders in India (the major supplier) are extensively exploited to the point of near extinction thus placing it in the red list of endangered species under IUCN guidelines. The species is propagated through seeds (Dayanand and Lohidas, 1988), seed propagation encountered with number of problems owing to low fruit set, hard pod and seed coat, dormancy of the seed; extended germination period up to 90 days; low poor germination of 20% and conversion of 34% restricted the area expansion (Gopinatha *et al.*, 2004).

Failure in seed propagation may adversely affected the important regeneration mechanism through quality seed, leaving only the coppicing mode for the survival of the species. Seed possessed with dormancy up to six months to one year, type of dormancy has not yet been elucidated (Rao and Raju, 2002). Presence of dormancy cause prolonged germination. The growth of the seedling also not to the expected speed and vigour, due to number of reasons, resulting in poor crop establishment after transplanting (Kalimuthu and Lakshmanan, 1995).

Comment [BB1]: Please, look at Authors: <http://www.sciencedomain.org/page/guideline-for-authors>

Comment [BB2]: ABSTRACT BOLD, 11 FONT, LEFT ALIGNED
Formatted: Line spacing: single

Comment [BB3]: 1. INTRODUCTION (ARIAL, BOLD, 11 FONT, LEFT CAPS)

(Arial, normal, 10 font, justified) instruction about this section is given. After reading these instructions, copy this paragraph and begin typing. If you are using copy-paste option, use 'match destination formatting' in OR use 'paste special' option and 'unformatted Unicode text' option.

— Conventional vegetative propagation techniques such as grafting and air-layering have limitations in large-scale multiplication of this species and rooting of cutting was also found to be poor (Kesava Reddy *et al.*, 1990). Tissue culture has proved to be a promising technique for conservation and large scale multiplication of several woody species. However the members of *Fabaceae* have been difficult to culture *in vitro* owing to their recalcitrant nature, roots were robust and vigorous in air layers compared to stem cuttings, but the rate of manipulation is comparatively low and not enough to transplant in the nursery and main field (Rao and Raju, 2002). Based on the above reasons, the multiplication of the species largely depends on seed (Dayanand and Lohidas, 1988).

Formatted: Font: Italic

2. Materials and method:

Comment [BB4]: 2. MATERIALS AND METHODS / EXPERIMENTAL DESIGN / MATERIALS AND METHODS (ARIAL, BOLD, LEFT ALIGNED, CAPS)

The study was carried out during 2015-16 at Forest College and Research Institute, Mettupalayan Tamil Nadu, India.

Comment [BB5]: Please follow the guidelines of the reviewers (Rev_AIR_34246_lbr.doc; Rev_AIR_34246_Moh.doc). It is too much subheadings and subheadings is not necessary.

2.1. Seed source

Seeds of *Pterocarpus santalinus* were collected during June, 2015 from the Chittoor, Andhra Pradesh sources.

2.2. Treatment details

2.2.1. Cow dung slurry

Four hundred pods from source were separately mixed with cow dung slurry and kept (1:2 ratio of water and cow dung) for different duration *viz.*, 24, 48, and 72 h.

Preparation of cow dung slurry

About 500 g of cow dung slurry was mixed with 500 ml of water and mixed well to make slurry.

2.2.2. Effect of Termite digestion on seed germination and seedling germination

About 400 pods were exposed to live termite mound for ten days; after ten days pods were collected.

2.3. The observations made are described as follows.

2.3.1. Days to initial germination

The nursery bed was observed daily, for seedling emergence. The day on which the first seedling emerged was expressed as days to initial germination.

2.3.2. Days to final germination (Mauromicale and Cavallaro, 1995)

The number of days on which the last seedling emerged was recorded and expressed as days to final germination.

2.3.3. Speed of germination

Speed of germination was calculated by the following formula,

$$\text{Speed of germination} = n_1/d_1 + n_2/d_2 + n_3/d_3 + \dots$$

Where, n = number of germinated seeds; d = number of days

2.3.4. Germination per-cent

The number of normal seedlings produced in each replication was counted, and average was expressed in per cent.

$$\text{Germination percentage} = \frac{\text{Number of normal seedlings}}{\text{Total number of seed sown}} \times 100$$

2.3.5. Seedling length

All normal seedlings of each treatment were measured for length from root tip to shoot tip and the average was expressed in cm.

2.3.6. Dry weight

All normal seedlings were dried under shade for 24 h and then dried in hot air oven maintained at $85 \pm 1^\circ\text{C}$ for 48 h. It was cooled in a desiccator for 30 minutes and weighed. The values were expressed as 'g seedlings⁻¹'.

2.3.7. Vigour Index (Abdul-Baki and Anderson, 1973)

Vigour index (VI) was computed using the following formula and expressed as whole number.

$$\text{VI} = \text{Germination percentage} \times \text{dry weight (g/seedling)}$$

2.3.8. Survival percentage

One month old seedlings were transplanted to polythene bags (size 23 cm x 15 cm) containing nursery mixture; after one month of transplanting number of survived seedlings were counted and expressed in percentage.

2.4. Statistical analysis

Result data (in per-cent) were transformed to arcsine values before statistical analysis in order to unify the variance of the data (Ansari *et al.*, 2012). The data were then analyzed by the 'F' test for significance at 0.05 level by using statistical software AGRESS.

3. Result and Discussion

—All the observed parameters were significant. Exposure of the pod to termite mound from 1 to 10 days did not show any remarkable increase either for germination percentage and speed of germination (Table 1).

Comment [BB6]: statistically?

-Among the observed parameters for the influence of cow dung slurry on seed germination and seedling characters, days to initial germination, final germination, speed of germination, germination percentage, seedling length, vigour index and survival percentage showed significant difference for treatmental effect (Table 2).

-The pods exposed to termite mound did not have any influence for all the recorded parameters and evidenced through statistical analysis. Even exposure of pod for a duration of 10 days did not have any positive effect and this might be due to hard veins on the surface of pod, shiny shell or due to presence of high quantity of phenols which might have prevented the termite activity. Such a non productive effort due to termite was reported by Sivaprakash (2003) in *Terminalia chebula* and *T. bellerica*,—but in many cases termites have an influence of increasing the germination through weakening of the coat and make tiny holes which facilitated the entry of water. Absence of such positive mechanism in *P. santalinus* is yet to be studied.

The use of bio-regulators in enhancing seed germination and seedling vigour is well known (Pampanna and Sulikeri, 2001). Presence of biologically active substances, microbes, weak acids of some bioregulators like cow dung resulted in enhanced germination in *Calophyllum inophyllum* (Rajesh *et al.*, 2011); Khirmi (Shinde and Malshe, 2015) and *Melia azadirachta* (Sujatha and Manjappa, 2015)

4. Conclusion

In the present study, the pods mixed with cow dung slurry for 24 h, resulted in 17 per-cent increased germination than the control, apart from germination enhancement more uniform germination with higher seedling vigour and survival percentage was observed. This might be due to the corrosion of pod coat by the weak acids, digestion of thin and strong veins by the microbes present in cow dung, both together might have resulted in the opening of pores; entry of

growth stimulants of cow dung and adequate water through the opened pores might have resulted in positive performance.

Table 1. Effect of treatment on seed germination characteristics

| Treatment | Days to initiate germination | Days to final germination | Speed of germination | Germination % |
|----------------|------------------------------|---------------------------|----------------------|---------------|
| T ₀ | 14.00 | 53.00 | 00.22 | 34(35.66) |
| T ₁ | 12.00 | 32.75 | 00.87 | 51(45.57) |
| T ₂ | 10.00 | 28.75 | 00.78 | 50(45.00) |
| T ₃ | 10.25 | 31.75 | 00.86 | 51(45.57) |

Comment [BB7]: Please follow reviewer: Rev_AIR_34246_Moh.doc

| | | | | |
|----------------------|-------------|-------------|-------------|-------------|
| T₄ | 13.00 | 50.75 | 00.21 | 36(36.86) |
| Mean | 11.85 | 39.40 | 00.58 | 44(41.73) |
| SE.D | 0.23 | 1.67 | 0.01 | 1.59 |
| CD (P ≤0.05) | 0.46 | 3.40 | 0.01 | 3.23 |

T₀- Control, T₁ - Cow dung slurry 24h, T₂ - Cow dung slurry 48h, T₃- Cow dung slurry 72h,
T₄ - Effect of Termite digestion

Table 2. Effect of treatment on seedling characteristics

Comment [BB8]: Please follow reviewer: Rev_AIR_34246_Moh.doc

| Treatment | Seedling length (cm) | Dry weight (g) | Vigour Index | Survival (%) |
|----------------------|-----------------------------|-----------------------|---------------------|---------------------|
| T₀ | 11.22 | 0.10 | 07.41 | 91.00 |
| T₁ | 17.57 | 0.16 | 08.36 | 92.50 |
| T₂ | 17.27 | 0.17 | 08.40 | 89.25 |
| T₃ | 17.55 | 0.15 | 08.33 | 89.50 |
| T₄ | 10.85 | 0.21 | 06.99 | 89.25 |
| Mean | 14.89 | 0.16 | 7.90 | 90.30 |
| SE.D | 0.61 | 0.01 | 1.58 | 0.4 |
| CD (P ≤0.05) | 1.25 | 0.02 | 2.60 | 0.8 |

References:-

Comment [BB9]: Please, look at Authors: <http://www.sciencedomain.org/page/guideline-for-authors>

Abdul Baki, A.A. and J.D. Anderson. 1973. Vigor determination in soybean seed by multiple criteria. **Crop Sci.**, **13**: 630-633.

Ansari, O., H.R. Choghazardi, Z.F. Sharif and H. Nazarli. 2012. Seed reserve utilization and seedling growth of treated seeds of mountain ray (*Secale montanum*) as affected by drought stress. **Cercetări Agronomice în Moldova**, **2(150)**: 43-48.

- Dayanand, T. Lohidas, T. 1998. Effect of different treatments on pod germination of red sanders. (*Pterocarpus santalinus*) L. **Indian J. Forester** 11: 87-88.
- Gopinatha, K., B.G. Reddy, P.K. Jha and R.U. Kashappa. 2004. Working plan for the management of forests of Kadapa division. Andhra Pradesh Forest Department, Government of Andhra Pradesh, India.
- Kalimuthu K, Lakshmanan KK. 1995. Effect of different treatments on pod germination of *Pterocarpus* species. **Indian Journal of Forestry** 18,104
- Kesava Reddy, K. and K.P. Srivasuki. 1990. Vegetative propagation of red sanders (*Pterocarpus santalinus* Linn.). **Indian Forester**, **116**: 536-540.
- Mauromicale, G. and V. Cavallaro. 1995. Effects of seed osmo priming on germination of tomato at different water potential. **Seed Sci. Technol.**, **23**: 393-403.
- Pampanna, Y. and G.S. Sulkeri. 2001. Effect of growth regulators on seed germination and seedling growth of Sapota. **Karnataka Journal of Agricultural Science**, **14**: 1030- 1036
- Rajesh, P., D. Gunaga and R. Vasudeva. 2011 Influence of seed size on germination and seedling growth in *Mammea suriga*. **Karnataka J. Agric. Sci.**, **24(3)**: 415-416
- Rajesh, P., D. Gunaga and R. Vasudeva. 2011 Influence of seed size on germination and seedling growth in *Mammea suriga*. **Karnataka J. Agric. Sci.**, **24(3)**: 415-416.
- Rao S. Purnachandra and A. J. Solomon Raju 2002. Pollination ecology of the Red Sanders *Pterocarpus santalinus* (Fabaceae), an endemic and endangered tree species. **Current Science**, VOL. 83, NO. 9, 10
- Shilpa, B., C.S. Reddy, K. Sudha, S. Sudhakar and V.S. Raju. 2008. Vegetation cover mapping and landscape level disturbance gradient analysis in Warangal district, Andhra Pradesh: India using satellite remote sensing and GIS. **Space Res. J.**, **1**: 29–38.
- Shinde, V.V. and K.V. Malshe. 2015. Effect of cattle urine and cowdung slurry as seed treatment on germination and growth of Khirni (*Manilkara hexandra*) **Journal of Eco-friendly Agriculture**, **10(2)**: 128-130.
- Sivapraksh, M. and P.R. Renganayaki. 2003. Germination improvement in *Terminalia bellerica* and *Terminalia chebula*. **M.Sc. Thesis**, Tamil Nadu Agricultural University, Coimbatore.

Sujatha, V.N. and K. Manjappa. 2015.Effect of pre-sowing treatments on seed germination of *Melia azedarach* L.**Karnataka J. Agric. Sci.**, **28(2)**: 269-270.