

## **Original Research Article**

# **AGRO-POTENTIALITY OF TREATED PAPERBOARD MILL EFFLUENT ALONG WITH ORGANIC AMENDMENTS ON GROWTH AND YIELD CHARACTERISTICS OF OKRA**

## **ABSTRACT**

Paper and pulp industry is categorized under 17 most polluting industries due to discharge of large volumes of black liquor with high nutrients. It ranks third in the world in terms of utilization of fresh water for processing. The paper mills generating appreciable quantities of solid wastes and effluent every day. The average quantity of water consumed for each tonne of paper produced is about 300 m<sup>3</sup> and this significant amount recur as effluent causing wide spread of environmental pollution. Results show that irrigation with treated paperboard mill effluent increased the plant growth characters as well as the fruit quality parameters. Compared to well water, reuse of treated paperboard mill effluent increased the plant vegetative growth vigorously. This is because of the nutrients present in the treated paperboard mill effluent. The scientific ways and means of recycling this wastewater in an integrated, eco friendly manner is the main objective of this study.

**Key words:** Treated Paperboard Mill Effluent, Okra, Growth characteristics, Organic Amendments

## **1. INTRODUCTION**

Paper and pulp industry is categorized under 17 most polluting industries due to discharge of large volumes of black coloured liquor with high nutrients. It ranks third in the world in terms of utilization of fresh water for processing. The paper mills generating appreciable quantities of solid wastes and effluent every day. The average quantity of water consumed for each tonne of paper produced is about 300 m<sup>3</sup> and this significant amount recur as effluent causing wide spread of environmental pollution [4]. Utilization of this wastewater for irrigation purpose makes the soil healthier due to the presence of plant nutrients [15]. Some researchers had reported that the nutrient status of soil like N, P and K has been increased with proper application of paper mill wastewater [3]. The crop yield and growth also showed positive correlation with the increasing dose of irrigating paper mill waste water and application of sludge compost up to certain limits [10]. Keeping the aforementioned points in mind, the study has been designed to assess the influence of paper board mill effluent irrigation on growth and yield of okra.

## **2. MATERIAL AND METHODS**

### **2.1. Experimental Design**

Field experiment was conducted in ITC (PSPD) model farm, Thekkampatti, Mettupalayam, Coimbatore District, Tamil Nadu to assess the impact of ITC treated effluent and solid wastes on crop growth, yield and quality of bhendi and soil. The experiment was conducted under factorial randomized designed and replicated by three times. The number of treatments was used for the cultivation of okra is seven with two factors viz., well water and effluent irrigation. The seven treatments are T<sub>1</sub>-Control (100 % NPK), T<sub>2</sub>-FYM 25 t ha<sup>-1</sup>+NPK, T<sub>3</sub>-ETP Sludge 5 t ha<sup>-1</sup>+NPK, T<sub>4</sub>-Biochar 2.5 t ha<sup>-1</sup>+NPK, T<sub>5</sub>-Vermicompost 3.5 t ha<sup>-1</sup>+NPK, T<sub>6</sub>-Pressmud 6 t ha<sup>-1</sup>+NPK and T<sub>7</sub>-Fly ash 5 t ha<sup>-1</sup>+NPK. The spacing between the crops is 45 x 30 cm and irrigated at weekly intervals.

### **2.2. Plant Biometric Observation**

Plant height, Number of leaves per plant, Number of branches, Internodal length, Leaf Area Index was measured at different stages of plant. The height of the plant from the ground level to the tip of the main stem was measured at 30, 60 DAS and at harvest stage and expressed in cm. Total number of fully opened leaves in a plant was recorded at 30, 60 DAS and at harvest stage and expressed as number of leaves per plant. The total number of main branches whose origin is in the leaf axils and the main stem was recorded on 10 randomly selected plants at 30, 60 DAS and at harvest stage.

### 2.3. Yield and Quality Parameters

Individual fruit weight, Fruit length, Fruit girth, Fruit yield and Dry matter production (DMP) is recorded at matured stage of the plant. Fruit yield was calculated by weighing of all the fruits in the plants from each plot and expressed in  $\text{t ha}^{-1}$ . Five plants at random were cut close to the ground level for the estimation of DMP. Samples were sun dried for three days followed by oven drying at  $70^{\circ}\text{C}$  till constant weight obtained. The dry weight of the plant samples were recorded and expressed in  $\text{kg ha}^{-1}$ . The leaf sample of 500 mg was taken and total phenol content in the leaves was estimated using Folin-Ciocalteu reagent and expressed as  $\text{mg } 100 \text{ mg}^{-1}$  of the material [7]. Crude fibre content was estimated by the method [2] and expressed in percentage. The protein content of fruits was estimated by the method suggested by Lowry *et al.* (1951) [6] and expressed in per cent.

### 2.4. Statistical Analysis


The data generated during this investigation for various characters were statistically analysed by the method given by Gomez and Gomez (1984). Results are presented and discussed at five per cent probability level uniformly. Treatment differences that are not significant were noted as non significant (NS).

## 3. RESULTS AND DISCUSSION

### 3.1. Plant Biometric Observation

Effluent irrigation produced taller and stronger plant than the well water irrigation. Among the treatments, VC + NPK had a favorable influence on plant growth attributes in bhendi at all the stages of growth period with effluent irrigation compared to well water irrigation (Fig. 1). This is because the treated effluent supplies appreciable amounts of plant nutrients than well water. Similarly, Udayasoorian *et al.* (2003) [14] reported that effluent irrigated bhendi recorded higher plant height and yield than the well water irrigated bhendi. In the present study, the vermicompost might have contributed all essential plant nutrients which helped to maintain the soil moisture ultimately resulted in the promotion of the plant height of bhendi. Najar and Khan (2013) [8] reported that the application of vermicompost increased the number of clusters per plant, fruits per cluster, number of fruits per plant, mean fruit weight and yield in tomato. Roseta and Innocent (2012) [9] found that agro industrial effluents and agricultural wastes like poultry manure increased the growth parameters like plant height, number of leaves per plant, stem girth and dry matter of the bhendi in different combinations.


**Fig. 1. Effect of treated paperboard mill effluent irrigation and solid waste on plant height and number of branches of bhendi**



### 3.2. Yield Parameters

In the present study, application of vermicompost along with NPK had favorable influence on yield attributing characters of bhendi viz., fruit weight, fruit length and fruit girth with effluent irrigation. The treatment receiving vermicompost along with NPK along with effluent irrigation registered the highest fruit weight and fruit yield compared to other treatments (Fig. 2). This could be due to increased level of major nutrients in the effluent which might have contributed to higher fruit yield of bhendi. Similar observations were recorded in cowpea [1], chillies and brinjal [14].

**Fig. 2. Effect of treated paperboard mill effluent irrigation and solid waste on fruit weight and yield of bhendi**




### 3.3. Quality Parameters

The quality parameters of bhendi viz., total protein, total phenol and crude fibre content were higher in paperboard mill effluent irrigation when compared to well water irrigation. In bhendi, the quality parameters were better in treatment receiving VC + NPK (Fig. 3). This might be due to combined use of treated effluent along with amendments, which might have provided enough nutrients with better physical and microbial environment and thus improving the soil fertility and ultimately resulted in improved quality parameters.

Similar results were also reported in radish and onion [12], chillies and brinjal (Udayasoorian and Ponmani, 2009) when these crops are grown in organic amended soil along with paperboard mill effluent irrigation. In *Allium cepa*, application of vermicompost combined with mineral fertilizers increased the bulb qualities like bulb size, total number of bulbs and fresh weight of bulbs [11]. Najar *et al.* (2013) reported similar findings that the application of vermicompost increased the number of marketable fruits and in the same way decreased the non marketable fruits which are infested. Biochemical components of bhendi like crude proteins, crude fibre and crude carbohydrates were found maximum with 25 per cent paperboard mill effluent fertigation [5].

**Fig. 3. Effect of treated paperboard mill effluent irrigation and solid waste on total phenol, total protein and crude fibre of bhendi**



### 4. CONCLUSION

Application of vermicompost along with NPK under treated paperboard mill effluent irrigation had a favorable effect on the growth of bhendi. The yield attributes viz., fruit length, fruit girth and individual fruit weight were significantly increased in VC + NPK under treated paperboard mill effluent irrigation. The yield increase was 37.8 per cent in bhendi under VC + NPK than that of control (100 %

NPK). The quality parameters of bhendi viz., total phenol, crude fibre and total protein were not affected due to effluent irrigation along with solid waste application. But a slight increase was noticed in quality parameters. Based on the results, it could be concluded that the treated paperboard mill effluent can be used as an effective irrigation source with addition of vermicompost along with NPK. Treated effluent irrigation and vermicompost had provided necessary plant nutrients to bhendi, resulting in higher yield and without causing any adverse effect on soil health and crop quality parameters. These findings conclude that the future perspective of treated effluent in agriculture is favorable due to its effect on increased crop yield and growth, but there is also a possible accumulation of various nutrients and heavy metals in soil and in the ground water that may cause potential problems after long term reclaimed wastewater irrigation.

## References

1. Chopra, A. K., S. Srivastava and V. Kumar. Comparative study on agro-potentiality of paper mill effluent and synthetic nutrient (DAP) on *Vigna unguiculata* L. (Walp) Cowpea. *J. Chem. Pharm. Res.*, 2011; 3(5):151-165.
2. Chopra, S.L and J.S. Kanwar. *Analytical Agricultural Chemistry*. Kalyani Publishers, New Delhi, p.162, 1982.
3. Dhevagi, P., G. Rajannan and G.Oblisami. Effect of paper mill effluent on soil microflora of maize. *J. Indus. Pollu. Contr.*, 2000; 16: 95-105.
4. Hazarika S., Talukadar N.C., Borah K., Barman N., Medhi B.K., Thakuria D. and Barooah A.K. Long term effect of pulp and paper mill effluent on chemical and biological properties of a heavy textured acidic soil in Assam. *J. Indian Soc. Soil Sci.*, 2007; 55, 45-51.
5. Kumar, V and A.K. Chopra. Ferti-irrigational effect of paper mill effluent on agronomical characteristics of *Abelmoschus esculentus* L. (Okra). *Pakistan J. Bio. Sci.*, 2013; 16(22):1426-1437.
6. Lowry, O.H., N.J. Rosebrough, A.L. Farr and R.J. Randall. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.*, 1951; 193(1): 265–275.
7. Malick, C.P and M.B. Singh. *Plant enzymology and Histoenzymology*. Kalyani publishers, New Delhi, p.286. 1980.
8. Najar, I.A and A.B. Khan. Effect of vermicompost on growth and productivity of Tomato (*Lycopersicon esculentum*) under field conditions. *Acta Biologica Malaysiana*, 2013; 2(1): 12-21.
9. Roseta E.C and I.C. Innocent. Agro industrial effluents and agricultural wastes effects on soil chemical properties and yield of okro (*Abelmoschus esculentus* L. Moench). *J. Environ. Earth Sci.*, 2012; 2(5): 85-89.
10. Saha, N., A.C. Das and D. Mukherjee. Effect of decomposition of organic matter on the activities of microorganisms and availability of nitrogen, phosphorus and sulphur in soil. *J. Indian Soc. Soil Sci.*, 1995; 43: 210-215.
11. Srivastava, P., M. Gupta, R. Upadhyay, S. Sharma, S. Shikha, N. Singh, S. Tewari and B. Singh. Effects of combined application of vermicompost and mineral fertilizer on the growth of *Allium cepa* L. and soil fertility. *Z. Pflanzenernähr. Bodenk.*, 175: 2012; 101-107.
12. Srivastava. R.K. Effect of Paper Mill Effluent on Seed Germination and Early Growth Performance of Radish and Onions. *J. Ecotoxicol. Environ. Monitoring*, 1991; 1(1): 13-18.
13. Udayasoorian, C and P. C. Prabu. Impact of treated paper mill effluent irrigation on quality of soil and ground water. *Madras Agric. J.*, 2003; 90(7-9): 540-541.
14. Udayasoorian, C and S. Ponmani. Effect of treated paperboard mill effluent irrigation on soil health and yield of vegetable crops. *J. Environ. Res. Develop.*, 2009; 3(3):879-889.

- 170 15. Velusamy, K, C. Udayasoorian, and Praghadeesh Manivannan. "Influence of Treated Paperboard Mill  
171 Effluent Irrigation on Soil Micro Flora and Enzyme Activities." *Advances in Life Sciences* 5.6 (2016):  
172 2103-2109.