

Original Research Article**Evaluation of Radiation Health Risk Due to Gamma Exposure From River Water Around Oil Bunking Centre In Rivers State, Nigeria****ABSTRACT:**

Aim: The aim of this study is to assess radiation health risk due to gamma exposure from river water around oil bunkering centers in Rivers state, Nigeria. **Study design:** This study was purely an experimental work. **Place and Duration of Study:** Sampling started from the meeting point of Otamiri tributary and Imo River at the Abia/Rivers boundary to over seven kilometer along the Imo River; between July and January, 2017. **Methodology:** 20 samples of river water were collected along coastal shore of Imo River with pre-washed 1.5 ml Polypropylene bottles. The bottles were rinsed with the water before collection and acidified immediately after collection with few drops of nitric acid. The bottles were sealed tightly with vinyl tapes and kept in the laboratory for 4 weeks for secular equilibrium of the radionuclides. The activity concentration of the radionuclides were measured using well calibrated Sodium Iodide detector. **Results:** The mean activity concentrations of ^{238}U , ^{232}Th and ^{40}K were 2.02 ± 0.02 , 3.59 ± 0.21 and $10.43 \pm 1.13 \text{ BqL}^{-1}$. The mean annual effective dose estimated for infants, children and adult citizen that ingest river water sampled were 58.64, 0.19 and 0.24 mSv respectively. The values of annual effective dose for infant and adults exceeded the reference level of 0.26, 0.2 and 0.10 mSv^{-1} respectively while that for children are within the safe reference level. The estimated total cancer risk to adults citizen and the lifetime hereditary effects show that 53 out of 10,000 citizens may suffer some form of cancer fatality and 596 out of 1000,000 citizens may suffer some form of hereditary effect since the values exceeded the USEPA recommended range. **Conclusion:** The result of this study show that the river water under study have been radiologically impacted by oil bunking activities and may cause significant health risk. Hence few recommendations were made in this work which will help to reduce radiation exposure and possible health impact.

Keywords: Radionuclide, lifetime cancer risk, committed dose, hereditary effect, Imo river..

1. Introduction

River water does not exist in a pure form for any appreciable length of time in nature. Even while waterfalls as rain, water picks up small amount of contaminants from the atmosphere and move as it filters through the ground^[1]. Those contaminants may be natural or anthropogenic including biological, chemical, physical and radiological impurities such as industrial and commercial solvents, heavy metals, acid salts, and radioactive materials. The major sources of natural radionuclides in water results from weathering and recycling of terrestrial minerals and rocks that give rise to ^{40}K , ^{232}Th , ^{235}U and ^{238}U . The later three decay naturally to produce other important radioactive isotopes of elements including radium, radon, polonium and lead^[2, 3].

Water can pick up radioactive materials as it flows through rocks, soils or cracked cement surrounding of water source thereby contaminating the water sources.

In a closed system the progeny of thorium and uranium are present in concentrations determined by the concentration of parent uranium and thorium isotopes and the time since the system became closed to nuclide migration. In nature closed systems rarely exist and predictions regarding nuclide concentrations in water bodies invariably include large uncertainties. These nuclides and their decay products are found in ground and spring waters in element specific concentrations dependent on complex hydrogeologic processes and conditions (dissolution, transport and ion-exchange processes as well as redox potentials and pH-conditions of the aqueous system). These hydrogeological processes result in non-equilibrium conditions between parent nuclides and their progeny. However, characteristic behaviour in the natural environment can provide a basis for assumptions regarding probable behaviour of nuclides used in the radioactivity screening assessment ^[4].

In the oxidised zone of the earth's near-surface environment thorium and uranium may both be mobilised, but in different ways. Thorium has an extremely low solubility in natural waters. There is a close correlation of thorium concentration and detrital content of water. Thorium is almost entirely transported in particulate matter and is bound in insoluble resistant minerals or is adsorbed on the surface of clay minerals. Even when thorium is generated in solution by radioactive decay of U-234 it rapidly hydrolyses and adsorbs on to the nearest solid surface. Products of radioactive decay in the U and Th series include radon (Rn) gas of which three isotopes exist. Rn-222 is the longest-lived and most abundant. Loss of radon will cause disequilibrium between members of a decay chain. Rn-222 has an appreciable solubility in water and is often found in concentrations far in excess of the parent nuclide radium, Ra-226. An Rn-222/Ra-226 activity ratio of 450 has been observed in ground waters from central England^[5]. Aeration of water and short half-lives make the contribution of radon negligible in ingestion dose calculations.

Illegal bunkering activities and crude method of refining crude oil along Imo river course has introduced a lot of hazardous waste into the water bodies of the area. Recently the entire Rivers state is experiencing massive air pollution (black soot) covering everywhere within Port Harcourt metropolis. Some speculations are pointing towards the illegal refining of crude oil in all those oil bunkering centres which produces some kind of explosions in the process. The inadvertent discharges of petroleum hydrocarbons or petroleum derived wastes streams from oil and gas productions activities are toxic to the coastal waters, soils and sediment near the discharge point^[6]. For human race, water is essential to life as air to breath. Thus, the importance of investigating the levels of radionuclide element in river water is very important ^[1] as river water serves as a major source of drinking water to many communities along Imo River. Estimation of radiation dose distribution is vital in assessing the health risk to a population and serves as a reference for documenting changes in environmental radioactivity due to anthropogenic activities^[7]. Hence, the aim of this work is to determine the radiological health risk of the populace from the activity concentration of ²³⁸U, ²³²Th and ⁴⁰K in river water collected from Imo

River near the bunkering sites in Rivers state. The result will help in assessment of the health impact of oil bunkering activities in Rivers state.

2. Materials and Methods

2.1 Study Area

The Imo River is located in the northern part of Rivers State in South eastern Nigeria. The study area is the boundary between Abia State and River State in the Niger Delta region. It lies between longitude $007^{\circ} 08' 11.9''$ and $007^{\circ} 11' 35.5''$ East and latitudes $04^{\circ} 54' 11.9''$ and $04^{\circ} 51' 37.8''$ North of equator (Figure 1). It is in southeastern Nigeria and flows 240 km into the Atlantic Ocean. Its estuary is around 40 km wide and the river has an annual discharge of 4 km^3 with 26,000 hectares of wetlands. The Imo's tributaries are the Otamiri and Oramirukwa[8]. The Imo River serves as drinking water sources for the surrounding communities.

Two geologic formations are covered in the study area, namely: Imo shale and Ameki formations

respectively. Imo shale consists of a thick sequence of blue and dark grey shales with occasional bands of clay-ironstones and subordinate sandstones [9]. It dips at angles 17° to 25° to the south-west and South [10]. It includes three constituent sandstones: the Igbabu, Ebenebe and Umuna Sandstones with the last two outcropping in the Imo River Basin. The Umuna sandstone is composed of thick sandstone units and minor shales and is generally less than 70m thick. The Ebenebe Sandstone occurs as a lens in the northwestern extremity of the Imo River Basin. It is similar in lithology to the Umuna sandstone but is relatively thicker with a maximum thickness of 130m [10]. Ameki Formation (Eocene) consists of sand and sandstones. The lithologic units of the Ameki Formation fall into two general groups [11, 12, 13]; an upper grey-green sandstones and sandy clay and a lower unit with fine to coarse sandstones, and intercalations of calcareous shales and thin shelly limestone.




Fig. 1: Map showing sampling points and industrial study areas

2.2 Sample Collection and Preparation

Sampling started from the meeting point of Otamiri tributary and Imo River at the Abia/Rivers boundary to over seven kilometer along the Imo River. In order to measure the specific activity concentration of natural radionuclides in river water, 20 samples of the water were collected. All the water samples were collected with 1.5litres linear polypropylene bottles which has been carefully washed before sampling. Containers for the samples were washed with a solution of detergent and then rinsed with freshly distilled Hydrochloric acid (HCl) to remove an inorganic material that might have stuck to the walls of the container before the samples were collected. About 1.5 L of each water sample were taken and 20 mL of 1 M HNO_3 added immediately^[14]. This is necessary to fix the radioactive elements in the samples. The samples were taken to the National Institute of Radiation Protection and Research (NIRP & R) University of Ibadan for preparation and analysis. The containers were sealed tightly and wrapped with thick vinyl tapes around their screw necks. Some 250 mL of each water samples in tightly covered cylindrical containers were stored for 4 weeks to reach secular equilibrium between ^{238}U and ^{232}Th and their respective progeny.

2.3 Gamma Spectroscopy

Activity measurements of radionuclides in river water collected were performed at the National Institute of Radiation Protection and Research Centre (NIRP & R), university of Ibadan with a gamma-ray spectrometry system with a thallium activated $3'' \times 3''$ sodium iodide on a NaI(Tl)

detector connected to ORTEC 456 amplifier^[15,16]. The detector in a 100 mm thick lead shield, was connected to a computer program called SAMPO 90 window that matched gamma energies to a library of possible isotopes. Since the accuracy of the quantitative measurements is dependent on the calibration of the spectrometry system and adequate energy, background measurement and efficient calibration of the system was made using Cs-137 and Co-60 standard sources from IAEA, Vienna. The analysis was performed using a Canberra S 100 computer analyzer. Standard of natural origin were prepared in the same manner as the samples, these standards are uranyl nitrate ($\text{UO}_2(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$) 502.18 mol/g, potassium chloride (KCl) 0.55 mol/g and thorium nitrate ($\text{Th}(\text{NO}_3)_4 \cdot 5\text{H}_2\text{O}$) 570.13 mol/g. One gram of each of the standard was taken and dissolved into a 200 mL distilled water to form a standard solution. It is subtle that 1 g of uranyl nitrate contains 0.474 g of uranium which has activity of 0.0294 Bq/L, so 1 g of potassium chloride contains 0.534 g of potassium which has activity of 0.706 Bq/L and 1g of thorium nitrate contains 0.859 g of thorium with activity of 0.0175 Bq/L^[17].

Spectrum were accumulated for background for a period of 29000 s at 900 volts to produce strong peaks at gamma emitting energies of 1460 Kev for ^{40}K ; 63.0 Kev of ^{214}Bi and 92.5 Kev of ^{228}Ac , which were used to estimate the concentration of ^{238}U and ^{232}Th respectively^[16]. The detector was calibrated with cesium-137 and cobalt-60 sources and the energy resolution is 39.5 and 22.2%, respectively^[18]. The configuration and geometry was maintained throughout the analysis

3. Radiological Risk Estimation

The annual effective dose from ingestion of radionuclide in water samples was estimated on the basis of the mean activity concentration of the radionuclides. This was done for different age categories. Assumptions on the rate of ingestion of water were made. In this work, the rate of water intake rates based on UNSCEAR^[19] recommendation of 0.5, 1.0 and 2.0 L/d infants, children and adults (≥ 17 years) respectively, were used for calculations. The conversion factors for ^{238}U , ^{232}Th and ^{40}K as reported by ICRP^[20] and presented in Table 3 were used for all the age groups.

The total annual effective dose due to ingestion of water was computed using the following formula^[21, 22](ICRP, 1996, Ndontchueng et al., 2013).

$$H_{\text{ing}} (\text{mSv} \cdot \text{y}^{-1}) = \sum_{i=1}^{i=3} DCF_{\text{ing}} (i) \times A_i \times I_i \quad (2)$$

Where $DCF_{\text{ing}} (i)$ is the dose coefficient of a particular radionuclide in Sv/Bq for a particular age categories. A_i is the specific activity concentration of radionuclide in the water sample measured in Bq/L and I_i is the radionuclide intake in liters per year for each age categories.

In addition to the estimated annual effective dose, the cancer and hereditary risk due to low dose without any threshold doses known as stochastic effect were estimated using the ICRP cancer risk methodology^[23]. Radiation risks to members of the public results from exposure to low dose radiation are normally known as chronic risk of somatic or hereditary damage of human tissues, thus much emphasis is always placed on the reduction of these radiological

risks to natural radiation. The nominal lifetime risk coefficient of fatal cancer recommended in the 2007 recommendations of the members of the public is $5.5 \times 10^{-2} \text{ Sv}^{-1}$. For hereditary effects, the detriment-adjusted nominal risk coefficient for the whole population as stated in^[23] for stochastic effects after exposure to low dose rates was estimated at $0.2 \times 10^{-2} \text{ Sv}^{-1}$.

The risk to population was then estimated using the 2007 recommended risk coefficient of ICRP report and assumed 70 years lifetime of continuous exposure of the population to low level radiation. According to ICRP methodology;

$$\text{Cancer Risk} = \text{Total annual Effective Dose (Sv)} \times \text{Cancer risk factor (Sv}^{-1}) \quad (3)$$

$$\text{Hereditary Effects} = \text{Total annual Effective Dose (Sv)} \times \text{Hereditary effect factor (Sv}^{-1}) \quad (4)$$

The recommended reference levels of the effective dose for infants, children and adults corresponding to one year consumption of drinking water are 0.26, 0.20 and 0.1 mSv^{-1} respectively.

Table 1: Effective Dose Coefficients (Sv/Bq) for ingestion of Radionuclides for members of the public to 70 years of age (ICRP, ^[20]; Publication 119)

S/N	Radioisotopes	Infant $\leq 1 \text{ year}$	Children 10 years	Adult > 17 years
1	^{238}U	1.4 E-07	6.8 E-08	4.5 E-08
2	^{232}Th	1.6 E-06	2.9 E-07	2.3 E-07
3	^{40}K	5.2 E-05	1.3E-08	6.2 E-09
Water intake		0.5 L/day	1.0 L/day	2.0 L/day

4.0 Results and Discussion

The activity concentration of ^{238}U , ^{232}Th and ^{40}K determined in river water from Imo River and the associated annual effective dose to infant, children and adult population of the communities are presented in Table 2 while the estimated cancer risks and hereditary effects of adult member of the public are shown in Table 3.

179 **Table 2: Activity Concentration of ^{238}U , ^{232}Th and ^{40}K in Water Samples and Total Annual**
 180 **Effective Dose for Different Age Categories**

S/N	Sample ID	Location	Activity Concentration (Bq/l)			Total Annual Effective Dose (mS)		
			^{238}U	^{232}Th	^{40}K	Infant	Children	Adult
1	SW1	Otamiri-Imo River	BDL	BDL	BDL	0.00	0.00	0.00
2	SW 2	NNPC-Alscon	BDL	BDL	3.50±0.27	33.0	0.0167	0.0158
3	SW 3	Obigbo Bridge	BDL	0.85±0.089	20.33±1.50	193.0	0.186	0.235
4	SW 4	Mama Town	BDL	BDL	14.36±1.08	137.0	0.068	0.0650
5	SW 5	Old Imo River	BDL	3.77±0.37	22.11±1.71	212.0	0.504	0.0417
6	SW 6	Imo River Village	1.93±0.50	7.89±0.76	BDL	0.169	0.883	1.388
7	SW 7	Back of Kom-Kom	BDL	BDL	BDL	0.00	0.00	0.00
8	SW 8	Imo River Railway	3.08±0.84	BDL	BDL	0.079	0.077	0.0604
9	SW 9	NNPC Pipeline	4.36±1.07	BDL	3.84±0.30	13.63	0.034	0.0426
10	SW 10	Imo River	BDL	BDL	13.34±0.95	9.018	0.064	0.061
11	SW 11	Imo River Division 1	BDL	4.34±0.43	BDL	1.271	0.459	0.728
12	SW 12	Imo River Division 3	1.10±0.30	BDL	1.43±0.10	13.63	0.034	0.0426
13	SW 13	Imo River Division 5	BDL	4.12±0.41	BDL	1.206	0.436	0.692
14	SW 14	Imo River Division 7	BDL	BDL	7.99±0.60	76.0	0.038	0.0362
15	SW 15	Imo River Banks 1	1.27±0.34	BDL	2.17±0.17	20.60	0.042	0.0515
16	SW 16	Imo River Banks 2	BDL	4.02±0.40	BDL	1.177	0.425	0.675
17	SW 17	Imo River Banks 3	BDL	BDL	14.98±1.11	142.55	0.071	0.068
18	SW 18	Imo River Banks 4	0.39±0.09	BDL	9.81±0.73	93.40	0.056	0.057
19	SW 19	Imo River Banks 5	BDL	BDL	11.32±0.82	107.72	0.054	0.0512
20	SW 20	Mmiri-Nwayi Division 14)	BDL	0.50±0.05	BDL	0.146	0.053	0.084
		Mean	2.02	3.59	10.43	58.64	0.19	0.24
		WHO, 2008 Standard	10.0	1.0	10.0	0.26	0.20	0.10

182 **Table 3: Estimated Cancer Risks and Hereditary Effects of Adult Member of the Public**

183

S/N	Sample ID	Total Annual Effective Dose (mSv)			Fatality cancer risk to Adult per year	Lifetime fatality cancer risk	Severe hereditary Effects in Adult per/y	Estimated lifetime hereditary Effects
		Infant	Children	Adult	$\times 10^{-6}$	$\times 10^{-4}$	$\times 10^{-7}$	$\times 10^{-6}$
1	SW1	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2	SW 2	33.0	0.017	0.016	0.87	0.61	0.32	2.22
3	SW 3	193.0	0.186	0.235	12.93	9.05	4.70	32.90
4	SW 4	137.0	0.068	0.065	3.57	2.50	1.30	9.09
5	SW 5	212.0	0.504	0.042	2.29	1.60	0.83	5.83
6	SW 6	0.169	0.883	1.388	76.35	53.44	27.76	194.30
7	SW 7	0.00	0.00	0.000	0.00	0.00	0.00	0.00
8	SW 8	0.079	0.077	0.060	3.32	2.32	1.21	8.45
9	SW 9	13.63	0.034	0.043	2.34	1.64	0.85	5.96
10	SW 10	9.018	0.064	0.061	3.35	2.34	1.22	8.522
11	SW 11	1.271	0.459	0.728	40.06	28.04	1.46	101.98
12	SW 12	13.63	0.034	0.043	2.34	1.64	0.85	596.4
13	SW 13	1.206	0.436	0.692	38.03	26.60	13.83	96.80
14	SW 14	76.0	0.038	0.036	1.99	1.39	0.72	5.06
15	SW 15	20.60	0.042	0.052	2.84	1.98	1.03	7.22
16	SW 16	1.177	0.425	0.675	37.12	25.98	1.35	9.45
17	SW 17	142.55	0.071	0.068	3.73	2.61	1.35	9.48
18	SW 18	93.40	0.056	0.057	3.15	2.20	1.14	8.01
19	SW 19	107.72	0.054	0.051	2.82	1.97	1.03	7.17
20	SW 20	0.146	0.053	0.084	4.62	3.23	1.68	1.18
	Mean	58.64	0.19	0.24	13.43	9.40	3.39	61.67

4.1 Specific Activity Concentration of ^{238}U , ^{232}Th and ^{40}K in River waters

The specific activity concentration of ^{238}U , ^{232}Th and ^{40}K in the river watersamples shown in Table 2 and ranges from BDL to $4.36 \pm 1.07 \text{ BqL}^{-1}$ with an average value of $2.02 \pm 0.02 \text{ BqL}^{-1}$, BDL to $7.89 \pm 0.76 \text{ BqL}^{-1}$ with an average value of 3.59 BqL^{-1} and BDL to $22.11 \pm 1.71 \text{ BqL}^{-1}$ with an average value of 10.43 BqL^{-1} . The result clearly show that ^{238}U are sparsely distributed along the coastal shore. This could be due to high mobility of uranium-238 in river water. Uranium - 238 were below detectable limit in most of the locations along the shore. This is in agreement with the fact that uranium in natural environment are variable in uranium content, depending mainly on factors such as contact time with uranium bearing rocks, uranium content of the contact rock, amounts of evaporation and availability of complexing ions. The ability of uranium to undergo inorganic dissociation and re-precipitation is probably the most important process in the natural environment to cause disequilibrium between the nuclides in the decay chains. The large variation of uranium observed in this work could be due to PH values which cause precipitation of uranium from the solution along the flow direction^[4].

The activity concentration of ^{40}K is highest at the old Imo River basin due to illegal oil and gas bunkering activities that releases its wastes into the river. The activity concentration of ^{232}Th in river water was relatively higher than that of ^{238}U because thorium is very insoluble^[24]. The activity concentration of ^{232}Th and ^{40}K are slightly higher than the reference levels of 1.0 and 10.0 BqL^{-1} while that for ^{238}U is within the reference levels. The results obtained in this work was compared with other works done in a similar environment within this country and other countries of the world as presented in Table 4. Figures 2 and 3 shows the comparison of the activity concentration of ^{238}U , ^{232}Th and ^{40}K in river water with standard value prescribed by ICRP, ^[20]. It shows that ^{238}U activity concentrations are lower than the standard value in all the locations while about six locations, activity concentration of ^{232}Th exceeded the standard value. The ICRP ^[20] and WHO, ^[25] regulations for drinking water quality does not include a listing for ^{40}K but specifies that the maximum allowable concentration limit for beta and photon emitters should correspond to a committed effective dose of 1.0 mSvy^{-1} from annual intake at the rate of two liters' of drinking water per day^[26].




Fig.2: Comparison of activity concentration of ^{232}Th with ICRP, 2012 Standard




Fig.3: Comparison of activity concentration of ^{40}K with ICRP, 2012 Standard




Fig. 4: Variations of total annual Effective dose for different age groups

The annual effective dose due to ingestion of the river water sampled was estimated for three different age groups: Infants, children and adults. The calculated total annual effective dose for different age groups as shown in Table 2 range from 0.00 to 212.0 mSv for infants, 0.00 to 0.883 mSv for children and from 0.00 to 1.388 mSv for adult with average values of 58.64, 0.19 and 0.24 mSv⁻¹ respectively. It can be observed from Figure 5 that the radiation dose received by infants is relatively higher than that received by children and adults. The WHO^[25] and UNSCEAR^[19] reference levels of the effective dose for infants, children and adult due to one year continuous ingestion of various drinking water are 0.26, 0.20 and 0.10 mSv⁻¹ respectively. The effective doses obtained were higher than the reference values for infants, and adults that consume river water but that for children are within the reference level of 0.2 mSv⁻¹. From the radiation protection point of view, life-long ingestion of these sampled river waters may cause significant radiological health problems.

In order to determine the radiation risk due to ingestion of ²³⁸U, ²³²Th and ⁴⁰K in river water sampled, ICRP^[20] methodology was adopted in the study and the results are shown in Table 3. The results of the cancer and non-cancer risk components were evaluated from the estimated total annual effective dose of the various age groups. The result of the estimated fatal cancer risk to adult per year in each of the stream water sampled ranged from 0.87×10^{-6} to 76.35×10^{-6} with the associated lifetime fatality cancer risk of 0.61×10^{-4} to 53.44×10^{-4} . The estimated hereditary effect to adult per year varied from 0.32×10^{-7} to 27.76×10^{-7} with its associated lifetime hereditary effect in adult of 1.18×10^{-6} to 596.40×10^{-6} . This means that in terms of the lifetime fatality cancer risk to adult approximately 53 out of 10,000 may suffer some form of cancer fatality and for the lifetime hereditary effect approximately 596 out of 1000,000 may suffer some hereditary effects. The negligible cancer fatality risk value recommended by

USEPA is in the range of 1.0×10^{-6} to 1.0×10^{-4} (ie 1 person out of 1 million to 10,000 persons suffering from some form of cancer fatality is considered trivial).

Comparing the estimated results of the lifetime fatality cancer risk in the present study with the acceptable risk factor, it can be seen that all estimated results of the lifetime fatality risk in adult member of the Nigerian population due to ingestion of radionuclide in the studied stream water are higher than the range of acceptable risk value recommended by USEPA.

Table 4: Comparison of activity concentration of ^{238}U , ^{232}Th and ^{40}K in water samples of Imo River Rivers State Nigeria and other studies in different parts of the world.

Samples	Country	^{238}U (Bq)	^{232}Th (Bq)	^{40}K (Bq)	References
Stream OD W (Nigeria)	Nigeria	0.52	1.8	27	[27]
Stream OW	Nigeria	4.62	4.06	42.57	[27]
Stream water	Nigeria	9.044 ± 3.11	2.28 ± 0.57	10.17 ± 23.47	[1]
Well OD	Nigeria	3.16	2.38	235.64	[27]
Mineral bottled water	Cameron	0.0	0.055	0.107	[22]
Portable water	Nigeria	0.000655	0.0005039	0.4154	[26]
Borehole water	Nigeria	0	0	7.40	[28]
Stream water	Nigeria	2.02	3.59	10.43	This study

4.2 Statistical Analysis

Basic statistics with statistical software package SPSS version 11.0 for windows was used to demonstrate the distribution and behavior of the measured radionuclide in stream waters and presented in Table 5. The statistical parameters determined includes the range (minimum-maximum), arithmetic mean (AM), arithmetic standard deviation (SD), median, mode, skewness, kurtosis and the type of frequency distribution for the three radionuclides for all the water samples.

The frequency distribution curves of ^{238}U , ^{232}Th and ^{40}K are shown in Figure 5. From Table 5, all the radiological parameters have positive skewness which shows that ^{238}U , ^{232}Th and ^{40}K have asymmetric distribution and only ^{40}K has a negative kurtosis indicating relatively flat distribution. Pearson's correlation analysis was also carried out to ascertain if there are mutual

relationship between the pairs of variables by calculating their linear correlation coefficient R^2 . It is important to note that a positive correlation among variables indicates similar source and behavior in the given environment.

Results of the Pearson correlation coefficient among all the three studied radionuclide and the associated radiological parameters are presented in Table 6. From Table 6, it can be observed that positive correlation exists among the three radionuclides and all the radiological parameters except ^{238}U having a negative correlation with $\text{AEDE}_{\text{children}}$ and $\text{AEDE}_{\text{adult}}$ indicating that uranium did not contribute to gamma emission on children and adult. Strong correlation were observed between ^{232}Th and ^{40}K while ^{238}U is weakly correlated with ^{232}Th and ^{40}K .

The strong positive correlation between ^{232}Th and ^{40}K shows that their origin and behavior in the coastal environment are the same while weak positive relationship between ^{238}U and the other two indicates that they may have the same origin but their behavior in the river environment differs. All the three radionuclides have strong positive correlation coefficient with the radiological parameters except for Uranium-238 that showed negative correlation with $\text{AEDE}_{\text{children}}$ and $\text{AEDE}_{\text{adult}}$. This means that two of the radionuclide only contributed significantly to gamma-ray emission at the sampling points.

Table 5: Descriptive statistics of radiological parameters

	U-238	Th-232	K-40	AEDEInfant	AEDEChildren	AEDEAdult
N Valid	18	18	18	18	18	18
missing	0	0	0	0	0	0
Mean	.6739	1.9161	6.9544	58.6442	.1945	.2442
Std. Error of Mean	.29716	.68676	1.774	16.90533	.05726	.08878
Median	.1800 ^a	.3636 ^a	3.6700 ^a	18.2767 ^a	.0660 ^a	.0607 ^a
Mode	.00	.00	.00	13.63	.03	.04
Std. Deviation	1.26076	2.91367	7.52305	71.72324	.24293	.37666
Variance	1.590	8.489	56.596	5144.224	.059	.142
Skewness	2.067	1.431	.746	1.027	1.679	2.087
Std. Error of Skewness	.536	.536	.536	.536	.536	.536
Kurtosis	3.795	1.065	-.741	-.261	2.420	4.132
Std. Error of Kurtosis	1.038	1.038	1.038	1.038	1.038	1.038
Range	4.36	9.00	22.11	211.92	.87	1.37
Minimum	.00	.00	.00	.08	.02	.02
Maximum	4.36	9.00	22.11	212.00	.88	1.39
Sum	12.13	34.49	125.18	1055.60	3.50	4.40

a. Calculated from grouped data.




Fig. 5 : Frequency Distribution of ^{238}U , ^{232}Th and ^{40}K in stream water

Table 6: Pearson Correlations of measured parameters

	²³⁸ U	²³² Th	⁴⁰ K	AEDEinfant	AEDEchild	AEDEAdult
²³⁸ U	1					
²³² Th	-0.12834	1				
⁴⁰ K	-0.36877	-0.18064	1			
AEDEinfant	-0.37325	-0.10098	0.923147	1		
AEDEchildren	-0.03014	0.680037	0.152366	0.0826631	1	
AEDEAdult	0.027668	0.642057	-0.42099	0.3598336	0.89566772	1

Conclusion

The activity concentration of ²³²Th and ⁴⁰K measured in river water collected from the meeting point of Otamiri tributary and Imo River at the Abia/Rivers boundary to over seven kilometer along the Imo River exceeded the reference level of 1.0 and 10.0 Bq l⁻¹ while the activity concentration

of ²³⁸U measured are within the reference level of 10.0 Bq l⁻¹. The meantotal annual effective dose determined for infant, children and adult population that drink river water from the Imo River are 58.64, 0.19 and 0.24 mSv y⁻¹ respectively. AEDE estimated for infant are 94% higher than the reference

Level of 0.26 mSv y⁻¹ and also higher than that for children and adult.

The estimated fatal cancer risk to adult per year and the lifetime hereditary effect shows that 53 out of

10,000 population may suffer some form of cancer fatality and approximately 596 out of 1000,000 might

Suffer some hereditary effects. Statistically all the radionuclide showed positive skewness and kurtosis except ²³⁸U. Pearson correlation of the radionuclides and all the radiological parameters showed positive correlation between ²³²Th and ⁴⁰K which indicate same origin and behavior in the coastal environment. ²³⁸U showed negative correlation with the radiological parameters which shows that uranium -238 did not contribute to gamma emission and probably had a different origin.

The result of this study showed that the activity of oil bunkering along the creeks, river shore has impacted negatively on the river water which in turn might lead to health challenges of the population that depend on such river for their daily living.

References

- [1] Ajayi, J.O, Adedokun, O. and Balogun, B.B. (2012). Levels of Radionuclide Content in stream water of some selected rivers in Ogbomosho land, South West Nigeria. *Journal of Environmental and Earth Sciences* 4(a) 835 – 837.
- [2] Arogunjo, M. A, Farai, I.P. and Furape, I.A (2004). Impact of oil and gas industry to the natural radioactivity distribution in the Delta region of Nigeria. *Nigerian Journal of Physics* 16, p131- 136.
- [3] Avwiri, G.O. and Agbalagba, O.E, (2007). Survey of gross alpha and gross beta radionuclide activity in Okpara creek, Delta state, Nigeria. Asia Network for science Information *Journal of Applied Science* 7(22) p3542 – 3547.
- [4] Kulin, A. V. Wonde S.N. and Mafejane, A. (2002). Radioactivity dose calculation and water quality evaluation guidelines for domestic water use. Department of water affairs and forestry document
- [5] UNSCEAR (2000). United Nations Scientific Committee on the Effects of Atomic Radiation. Sources and Effects of Ionizing Radiation (Report to the General Assembly, United Nations, New York).
- [6] Azpecza, O.S. and Szabo, Zoltan (1988). Radioactivity in ground water a review: U.S. Geological Survey Water Supply Paper 2325 p.50 – 57.
- [7] El-Tahaway M.S, Farouk M.A, Ibraheim M.N and El Mongery S.A.M (1994) Natural and Artificial Radioactivity in the Suez Canal bottom sediment and stream water, *Journal of Environmental Radioactivity*, 47(2) p201- 212.
- [8] Hany El-Gamal and Abdallah I.A. EL- Mageed (2014). Natural Radioactivity in water samples from Assiut city, Egypt. *International journal of pure and Applied Science and Technology*. 22(1):44-52.
- [9] Obed, R.I, Farai I.P. and Jibiri, N.N. (2005). Population Dose Distribution due to Soil Radioactivity Concentration levels in 18 cities across Nigeria. *Journal of Radioactivity Protection*, 2.
- [10] Kalin, M., Wheeler, W.N., Meinrath, G. (2005). The removal of uranium from miningwaste water using algal/microbial biomass, *Journal of Environmental Radioactivity* 78, 151–177.
- [11] West Larry (2006). World Water Day: A billion people worldwide lack safe drinking water. Wikipedia 2006.

- 378 [12] Alfaih. A.A. Osman, IsamSalih, Ibrahim A. Shaddad, Saif El Din , M.B. Siddeeg, Hatem
379 Eltaheb, Hajoldriss, Wadi Hamza, E.H. Yousif (2008). Investigation of natural
380 radioactivity levels in water around Kadugli, Sudan. *Applied Radiation and Isotopes*, 68:
381 1650 – 1653.
- 382 [13] Taskin, H, Karavus, M.A., Topuzoglu, B.A., Hindiroglu, S. and Karahan, G. (2009).
383 Radionuclide Concentrations in soil and life-time cancer risk due to radioactivity in
384 Kirklareli, Turkey. *Journal of Environmental Radioactivity*, 100 p49-53.
385
- 386 [14] Ahmed, N.K. (2004). Natural radioactivity of ground and drinking water in some areas of
387 upper Egypt, Turk. J. Eng. Environ. Sci., 28: 344-354.
- 388 [15] Jibiri, N.N., A.A. Mabawonku and S.J. OriadeUjiagbedion, (1999). Natural radionuclide
389 levels in soil and water around a cement factory in Ewekoro, Ogun State, Nigeria. J.
390 Phy., 5(11): 12-16.
- 391 [16] Guogang J., Giancarlo, T. (2007). Estimation of Radiation Doses to members of the
392 public from intakes of some important naturally occurring radionuclides. J. Environ.
393 Rad., 76:654-72.
- 394 [17] ICRP (1977). Publication 26. Recommendation of the International Commission on
395 radiological protection. Annals of the ICRP 1977.
- 396 [18] Oseji, J.O. (2010). Aquifer systems of Ndokwaland , Delta state, Nigeria. IJRRAS 5(3):
397 344-354.
- 398 [19] UNSCEAR, 2008. Sources and effect of ionizing radiation. Unscear,2008 report to the
399 general Assembly with scientific Annesvol II.
- 400 [20] ICRP publication (2012). Annals of the ICRP. Compendium of dose coefficients based
401 on ICRP publication 119.
- 402 [21] Isinkaye , M.O. and Emelue, H.U. (2015). Natural radioactivity measurements
403 andEvaluation of radiological hazards in sediment of Oguta Lake, South East
404 NigeriaJournal of Radiation Research and Applied Sciences , 8(459- 469)
- 405 [22] Ndontchueng, M.M., Simo, A., Njuelelem, E.J.M., Beyala, J.F., Kryeziu, D. (2013).
406 Preliminary study of natural radioactivity and radiological risk assessment in some
407 mineral bottled water produced in Cameroon. International Journal of Science and
408 Technology, 3(5): 271 – 276.
409
- 410 [23] Jibiri, N.N, Mabawonku A.A. and OriadeUjiaegbedion S.J. (1999). Natural radio nuclide
411 levels in soil and water around a cement factory in Ewekoro, Ogun state, Nigeria, *Journal*
412 *of Physics*, 5(11) 12 – 16.
413

- [24] Senthilkumar B, Dharamani V, Ramkumar S. and Philominathan P. (2010) Measurement of gamma radiation levels in soils samples from Thanjavur using x-ray spectrometry and Estimation of Population exposure. *Journal of Medical Physics*, Jan- Mar, 35 (1) p48-53.
- [25] WHO (2008) Guidelines for drinking water quality. Incorporating first Addendum vol.1. Recommendations 3rd radiological aspect Geneva.
- [26] Onoja, R.A. ,Adeyemo, D.J and Okoh, S. (2014). Determination of Natural Radionuclide concentrations in portable water supply of Northern part of Kaduna state. *Research journal of applied sciences, engineering and Technology* 7(14):2905-2907.
- [27] Jibiri, N.N., Chijioke, M.N., George, O.A. (2010). Radionuclide contents and physicochemical water quality indicators in stream, well and borehole water sources in high radiation area of Abeokuta, Southwestern Nigeria. *Journal of Water resources and Protection*, 2: 291 – 297
- [28] Meltem, D. and Gursel, K. (2010). Natural Radioactivity in various surface waters in Adana, Turkey. *Desalination*, 261:126-130