

Original Research Article

EXPLORATION OF THE ANTIMICROBIAL PROPERTIES OF *FICUS*

EXASPERATA LEAVES FROM AKURE METROPOLIS.

Abstract

Aim: The study was conducted to explore the antibacterial and antifungal properties of the leaf extracts of *Ficus exasperata* (Sand paper tree) *in vitro*.

Study design: Extracts from *Ficus exasperata* leaves were qualitatively screened for the phytochemical constituents, and their *in vitro* antimicrobial potency was evaluated against fourteen (14) fungal and bacterial isolates.

Results: The tested extracts contained tannins, flavonoids, terpenoids, alkaloids and cardiac glycosides whereas, saponin, steroids, phlobatannin and anthraquinone were absent. The acetone extract of the leaf demonstrated better antimicrobial activity against 10 of the test organisms. However, the highest antimicrobial activity (31.27mm) was exhibited by the methanol extract against referenced culture of *Staphylococcus aureus*. In addition, the extracts also displayed better antibacterial than antifungal activity. The minimum inhibitory concentration (MIC) of the extracts ranged between 0.391-1.563mg/mL, with the acetone extract displaying lower MIC values.

Conclusion: The occurrence of the observed phytochemicals in the extracts of *Ficus exasperata* (Sand paper tree) could be involved in the antimicrobial efficacy of the plant. The results from this study thus supports the folkloric use of the plants. Additionally, the plant could also be exploited for the production of drugs especially for staphylococci infections.

Keywords: medicinal plants; antimicrobial; phytochemicals; extracts; *Ficus exasperata*

Introduction

For ages, mankind has faced a constant battle with infectious diseases. This has led to increased morbidity and mortality especially among population from developing countries. Many populations have adopted the traditional healthcare system as a way of preventing and treating diseases of microbial origin [1]. Traditional medicine remains the most sort after, as it is considered safer, affordable, and readily available [1].

30 Due to the upsurge in resistance to conventional drugs by microbial agents, novel
31 antimicrobial agents from different biological sources have been sort after and reported to be
32 effective in combating pathogenic organisms. The use of herbal remedies containing plants or
33 part of plants has in recent years gained ground in developed countries [2].Pharmaceutical
34 companies have thus developed new antimicrobial drugs and also improved on the existing
35 ones through the modification of their structures with a view to increasing their efficacy [3].

36 *Ficus exasperata*otherwise known as the sandpaper tree is native to tropical Africa [4]. The
37 leaves of *F. exasperata*have been employed in folkloric medicine for the treatment of various
38 diseases such as ophthalmic and oral infections, venereal diseases, parasitic infection
39 (cutaneous, subcutaneous), leprosy, and malaria [5,6]. The study therefore investigates the
40 claim of the antimicrobial potential of *F. exasperata*, in a bid to develop novel antimicrobials.

41 **Materials and Methods**

42 **Collection and preparation of extracts from leaves Samples**

43 The leaves of *F. exasperata*were collected from its tree at a building near Life Spring
44 College, Apatapiti layout, Federal University of Technology, Akure, Ondo State
45 (Latitude:7.289N, Latitude:5.150E) Nigeria in the month of April, 2015. Samples of the
46 leaves were taken to the Department of Crop, Soil and Pest, FUTA for authentication.
47 Afterwards the leaves were cleansed with water, shade dried, grinded and stored in airtight
48 container.Thereafter, the powdered leaves (100g) of *F. exasperata*was weighed separately
49 into different plastic containers and 1000mL of 100% acetone and methanol added to the
50 containers for extraction. Aluminium foil was placed on each container before covering. Each
51 solution wasallowed to stand for 3 days with continuous stirring.The extracts were thereafter
52 obtained by filtering the solutions through a funnel fitted with a filter paper. The filtrates
53 were thereafteevaporated to dryness at 50 °C in a rotary evaporator (RE-52A; Union
54 Laboratory, England) with 90 rpm under reduced pressure. The obtained concentrated
55 extracts were stored in dark at 4 °C until further analysis.

56 **Phytochemical screening of leaf extract of *Ficus exasperata***

57 The plant extracts were subjected to qualitative phytochemical screening using standard
58 protocols described by Odebiyi and Sofowora [7], Trease and Evans [8], and Harborne [9].

59

60

61

Measurement of antibacterial and antifungal activities of leaf of *Ficus exasperata*

Varying concentrations of the leaf extracts of *F. exasperata* (3.125-100mg/mL) were prepared by dissolving different amount of the extracts in 5mL of 30% tween 20. For example, concentrations of 100, 50, and 25mg/mL were prepared by dissolving 500, 250 and 125mg of the extracts into 5ml of 30% tween 20 respectively. Afterwards the prepared extracts were sterilized by passing them through a 0.22µm millipore membrane filter. The agar well diffusion method as described by Schinoret *al.* [10] was employed in assessing the antimicrobial activity of *F. exasperata* leaf extracts. A total of 14 clinical and referenced microbial strains were used for the experiment. The test organisms were obtained from the Pathology and Clinical Laboratory (PATHCARE), Lagos State University Teaching Hospital, Lagos State, Nigeria and the Department of Microbiology, FUTA. Active broth cultures of the test organisms were prepared from stock cultures. To 5ml of nutrient broth 0.2 ml of bacterial culture was inoculated and incubated till it reached the turbidity equal to that of the standard 0.5 McFarland solution measured at 600nm which is equivalent to 10^6 – 10^8 CFU/ml. Suspensions of fungal spores were prepared from fresh mature (5days) cultures that grew at $26 \pm 1^\circ\text{C}$ on a Sabouraud dextrose agar. Spores were rinsed with sterile distilled water. The suspensions were then adjusted to 10^6 spores per/ml by microscopic enumeration with a cell counting hematocytometer. An aliquot of 100µL of bacterial and fungal solution was evenly spread on already solidified Mueller Hinton agar plates. Afterwards, wells of 7mm diameter were bored in the solidified Mueller Hinton agar plates using a sterile cork-borer. Thereafter, an aliquot of 100uL of the sterilized extract was added into the bored agar wells. The plates were thereafter incubated at 37°C for 24 hour for bacteria and at $26 \pm 1^\circ\text{C}$ for 48 to 72 hours for fungi. The plates were observed for clear zones of inhibition and the measurements were taken using a ruler calibrated in millimetres. Commercial antifungal drugs (clotrimazole, nystatin and gluseofluvin) and commercial antibacterial drug (ciprofloxacin (10µg), rocephin (25µg)) were used as the positive control, while 30% tween 20 was used as the negative control. To determine the minimum inhibitory concentrations (MIC), the agar diffusion method described above was used to screen the antimicrobial effect of the different concentrations of extracts (0.391-100mg/mL). The MIC value was determined by establishing visible growth of microorganisms. The boundary dilution without any visible growth was defined as the MIC for the tested microorganism at the given concentration.

95 Statistical analysis

96 Experiments were carried out in triplicates were applicable. The results were expressed as
97 mean \pm standard error of three values. Data analysis was carried out using the One Way
98 Analysis of Variance (ANOVA) and treatment means were compared using New Duncan's
99 Multiple Range Test (SPSS version 16). Differences were considered significant at $P < 0.05$.

100 Results

101 Table 1 shows the presence of tannin, flavonoid, terpenoids, alkaloids and cardiac glycosides
102 in *F.exasperata* leaf extracts, and the absence of saponin, steroids, phlobatannin and
103 anthraquinone.

104
105 **Table 1:** Qualitative phytochemical screening of *F.exasperata* leaf extracts

Phytochemical	Extracts	
	FEM	FEA
Saponin	-	-
Tannin	+	+
Flavonoid	+	+
Steroids	-	-
Terpenoids	+	+
Alkaloids	+	+
Phlobatannin	-	-
Anthraquinone	-	-
Cardiac Glycosides		
Legal test	+	+
Keller kiliani	+	+
Salkowski	+	+
Lieberman test	+	+

106

107 **KEYS:** FEA: Acetone leaf extract of *F. exasperata*; FEM: Methanol leaf extract of *F.*
108 *exasperata*.

109 The antimicrobial activity of the leaf extracts of *F. exasperata* showed that the acetone leaf
110 extract exhibited better activity against most of the test organisms used for the study (Table
111 2). However, the highest antimicrobial activity (31.27mm) was exhibited by the methanol
112 extract of *F. exasperata* against referenced *Staphylococcus aureus* and this was found to be
113 slightly higher than observed in the acetone extract (29.40mm) against the same organism. In
114 like manner, the leaves extracts displayed better antibacterial than antifungal activity. The

115 antifungal activity of the acetone extract of the plant was however a better than that of the
116 methanol extract.

117 Upon comparison of the activities of the leaf extracts against organism with Gram reaction
118 positive and Gram negative bacterial isolates, the Gram positive organism were more
119 susceptible than the Gram negative organism in most cases. The extracts antibacterial activity
120 was comparatively better than that of the commercial antibacterial drugs in most of the tested
121 organisms. Reverse was the case for the commercial antifungal drugs as they exhibited better
122 activity than the extracts. The acetone extract of *F. exasperata* was found to exhibit lower
123 minimum inhibitory concentration values than the methanol extracts. The results are
124 displayed in Table 3.

125 **Table 2:** Antimicrobial activity of leaves extracts of *F. exasperata* and commercial drugs

Test Organism	Zone of inhibition (mm)						
	FEA	FEM	CPX	R	CLOT	GRIS	NYST
<i>Salmonella</i> Typhi(ATCC 33489)	15.20 ± 0.12 ^c	12.43 ± 0.18 ^a	12.27± 0.15 ^a	14.30±0.12 ^b	NT	NT	NT
<i>Salmonella</i> Typhi	18.43 ± 0.18 ^c	12.47 ± 0.15 ^a	12.40±0.23 ^a	14.37±0.20 ^b	NT	NT	NT
<i>Staphylococcus aureus</i> (ATCC 43300)	29.40 ± 0.17 ^c	31.27 ± 0.15 ^d	14.40±0.12 ^a	15.50±0.17 ^b	NT	NT	NT
<i>Staphylococcus aureus</i>	27.33 ± 0.24 ^b	28.40 ± 0.12 ^c	15.43±0.20 ^d	12.50±0.12 ^a	NT	NT	NT
<i>Escherichia coli</i> (ATCC 35218)	22.20 ± 0.12 ^b	15.53 ± 0.15 ^a	11.20±0.12 ^b	14.27±0.15 ^c	NT	NT	NT
<i>Escherichia coli</i>	17.50 ± 0.26 ^{ab}	15.60 ± 0.17 ^{ab}	12.50±0.23 ^{ab}	49.97±36.67 ^a	NT	NT	NT
<i>Pseudomonas aeruginosa</i> (ATCC 27853)	12.27 ± 0.15 ^c	15.40 ± 0.12 ^b	15.43±0.15 ^a	16.33±0.18 ^b	NT	NT	NT
<i>Shigelladysenteriae</i>	17.40 ± 0.21 ^a	16.30 ± 0.12 ^b	14.40±0.12 ^b	14.27±0.15 ^c	NT	NT	NT
<i>Bacillus cereus</i>	15.60 ± 0.17 ^c	20.40 ± 0.12 ^d	12.33±0.18 ^a	14.53±0.20 ^b	NT	NT	NT
<i>Bacillus subtilis</i>	21.30 ± 0.12 ^d	12.43 ± 0.15 ^a	14.33±0.15 ^b	15.50±0.12 ^c	NT	NT	NT
<i>Candida albicans</i>	18.37 ± 0.23 ^d	10.27 ± 0.15 ^b	NT	NT	16.65 ± 0.68 ^c	20.50 ± 0.29 ^e	6.40 ± 0.21 ^a
<i>Aspergillus niger</i>	15.60 ± 0.17 ^b	3.47 ± 0.20 ^a	NT	NT	22.33 ± 0.33 ^d	21.67 ± 0.33 ^d	17.47 ± 0.32 ^c
<i>Aspergillus flavus</i>	12.40 ± 0.17 ^c	3.30 ± 0.12 ^a	NT	NT	25.00 ± 0.15 ^e	9.77 ± 0.15 ^b	18.73 ± 0.22 ^d
<i>Aspergillus fumigatus</i>	13.30 ± 0.17 ^c	5.30 ± 0.15 ^a	NT	NT	35.67 ± 0.44 ^e	9.33 ± 0.44 ^b	20.57 ± 0.30 ^d

126

127 Each value is expressed as mean ± standard error (n = 3). Values with different superscript within a row are significantly different at (P=0.05).

128 **Keys:** FEA: Acetone leave extract of *Ficus exasperata*; FEM: Methanol leave extract of *Ficus exasperata*; CPX: Ciprofloxacin (10µg);

129 R:Rocephin (25µg); CLOT: Clotrimazole(1mg/mL);GRIS: Griseofluvin(1mg/mL);NYST: Nystatin(1mg/mL);NT; Not tested.

130 **Table 3:** Minimum inhibitory concentration (mg/ml) of leaf extracts of *Ficus exasperata*

Test organisms	MIC (mg/mL)	
	FEA	FEM
<i>Salmonella typhi</i> (ATC 33489)	0.781	0.781
<i>Salmonella typhi</i>	0.391	1.563
<i>Staphylococcus aureus</i> (ATC 43300)	0.391	0.781
<i>Staphylococcus aureus</i>	0.391	1.563
<i>Escherichia coli</i> (ATC 35218)	0.391	0.781
<i>Escherichia coli</i>	0.391	0.391
<i>Pseudomonas aeruginosa</i> (ATC 27853)	0.781	0.781
<i>Shigelladysenteriae</i>	0.781	1.562
<i>Bacillus cereus</i>	0.391	0.391
<i>Bacillus subtilis</i>	0.391	1.562
<i>Candida albicans</i>	0.391	0.391
<i>Aspergillus niger</i>	0.391	0.391
<i>Aspergillus flavus</i>	0.391	1.563
<i>Aspergillus fumigatus</i>	0.391	1.563

131

132 **Keys:** FEA: Acetone leave extract of *F. exasperata*; FEM: Methanol leave extract of *Ficus*
 133 *exasperata*

134 Discussion

135 Plants remain an inexhaustible source of novel antimicrobials. Africa with its tropical and
 136 subtropical climate is richly blessed with an array of plants that have naturally acquired
 137 secondary metabolites in order to survive the harsh environment [1, 11]. Compounds with
 138 antimicrobial properties that also offer protection against drug resistant microorganisms have
 139 been isolated from medicinal plants [12, 13]. The present study investigated the secondary
 140 metabolite profile and antimicrobial efficacy of leaves of *F. exasperata*.

141 The presence of the observed secondary metabolites in the leaf extracts validates the
 142 medicinal potentials of this plants as these compounds have been reported to play a protective
 143 role against pathogenic organisms [13]. The absence of saponin, steroids, phlobatannin and
 144 anthraquinone in the extracts might be attributed to solubility of the compounds in the
 145 extraction solvent used.

146 The antimicrobial activity of the extracts could be attributed to the observed phytochemicals
 147 in the extracts. In addition, the variation observed in the antimicrobial activity of the extracts
 148 might be linked to differences in the type and amount of phytochemicals present in the

extracts. The structural differences in the cell wall of Gram positive and Gram negative bacteria may account for the higher susceptibility of Gram positive bacteria to the plant extracts. The complexity in the cell wall Gram negative bacteria gives them better buffering capacity thus making their cell wall less impermeable, whereas Gram positive bacteria have only an outer peptidoglycan cell wall which makes them more susceptible [14].

The higher antibacterial activity demonstrated by the extracts than antifungal activity is in consonance with findings of several authors [15, 16] that have reported higher sensitivity of bacteria to antimicrobials. The chitinous cell wall of fungi promotes lesser susceptibility to antimicrobials than bacteria [17]. Antibiotics have been mostly reported to produce better performance against microorganisms than plants as a result their higher purity and smaller molecular sizes which aid their penetration into the cell wall of the organisms [18]. The better activity produced by the extract suggests that they can be explored for potent antimicrobial compounds.

Conclusion

The plant extracts produced an effective performance against the growth of the tested organisms especially *Staphylococcus aureus*. The plant extracts could therefore be exploited for the production of antimicrobial drugs especially for staphylococci infections.

References

1. Mahomoodally MF. Traditional Medicines in Africa: An Appraisal of Ten Potent African Medicinal Plants. *Evidence-Based Complementary and Alternative Medicine*, 2013; Article ID 617459, 14 pages,
2. Chintamunnee V, Mahomoodally MF. Herbal medicine commonly used against infectious diseases in the tropical island of Mauritius. *Journal of Herbal Medicine*, 2012; 2:113–125.
3. Silva NCC, Fernandes JA. Biological properties of medicinal plants: a review of their antimicrobial activity. *The Journal of Venomous Animals and Toxins including Tropical Diseases*, 2010; 16(3): 402-413
4. Arbonnier M. **Trees, shrubs and lianas of West African dry zones**. Margraf Publishers, Paris, France; 2004.
5. Titanji VP, Zofou D, Ngemenya MN. The antimalarial potential of medicinal plants used for the treatment of malaria in Cameroonian folk medicine. *African Journal of Traditional Complementary and Alternative Medicines*. 2008;5:302-321.

6. Joseph B, Raj SJ. Phytopharmacological and phytochemical properties of three *Ficus* species - An overview. *International Journal of Pharmacy and Biological Sciences*, 2010; 1:246-253.
7. Odebiyi OO, Sofowora EA. Phytochemical screening of Nigerian medicinal plants. *Lloyd*. 1978;41(3):234–246.
8. Trease GE, Evans MC. **Pharmacognosy** (14th ed.). New Delhi, India: Elsevier; 2005.
9. Harborne JB. **Phytochemical methods. A guide to modern techniques of plant analysis** (3rd ed.). New Delhi: Springer PvtLtd; 2005.
10. Schinor EC, Salvador MJ, Ito IZ, Dias DA. Evaluation of the antimicrobial activity of crude extracts and isolated constituents from *Chrestascapigera*. *Brazilian Journal of Microbiology*, 2007; **38**: 145-149.
11. Manach C, Scalbert A, Morand C, Rémésy C, Jiménez L. Polyphenols: food sources and bioavailability. *American Journal of Clinical Nutrition*, 2004; 79(5):727–747.
12. Miyasaki Y, Rabenstein JD, Rhea J, Crouch ML, Mocek UM, Kittell PE, et al. Isolation and Characterization of Antimicrobial Compounds in Plant Extracts against Multidrug-Resistant *Acinetobacterbaumannii*. *PLOS ONE*, 2013;8(4): e61594.
13. ManganyiMC, Regnier T, OlivierEI. Antimicrobial activities of selected essential oils against *Fusariumoxysporum* isolates and their biofilms. *South African Journal of Botany*, 2015; 99:115–121.
14. Tadeh H, Mohammed E, Asres K, Gebre-Mariam T. Antimicrobial activities of some selected traditional Ethiopian medicinal plants used in the treatment of skin disorders. *Journal ofEthnopharmacology*, 2005;100:168-175.
15. Modarresi-Chahardehi A, Ibrahim D, Fariza-Sulaiman S, Mousavi L. Screening antimicrobial activity of various extracts of *Urticadioica*. *Revista de Biología Tropical*, 2012; 60(4):1567-1576.
16. Momoh AO, Oladunmoye MK, Adebolu TT. Evaluation of the Antimicrobial and Phytochemical Properties of Oil from Castor Seeds (*Ricinuscommunis* Linn). *Bulletin of Environment, Pharmacology and Life Sciences*, 2012; 1(10):21-27
17. Madigan MT, Martinko JM. **Brock Biology of Microorganisms**. New Jersey, USA, Pearson-Prentice Hall; 2006
18. Raza MW, Shad A, Pedler SJ, Karamat KA. Penetration and activity of antibiotics in brain abscess. *Journal of the College of Physician and Surgeons Pakistan*, 2005; 15(3):165-167.