

Original Research Article

A study on physical properties of okra seed: *ABELMOSCHUS ESCULENTUS* (L.)

Abstract

Physical properties of any seed is vital information for development of its metering mechanism, processing and storage system. An experiment was conducted to evaluate physical properties of okra seed of variety 'Varsha Uphar'. The study revealed that the average value of length, width, thickness and geometrical diameter of okra seed were 5.73 mm, 4.83 mm, 4.49 mm and 4.98 mm, respectively. The bulk density and true density okra seeds were found to be 0.54 g cm⁻³ and 1.07 g cm⁻³, respectively. The average volume of one okra seed was 108 mm³. The average angle of repose of okra seeds was 28.65°, which ranged from 27° to 30°. The study also inferred that the roundness of the seed was 78.2% ranging from 68.4% to 89.5% whereas the sphericity of the seed was 87.9% ranging from 82.4% to 95.8%. The porosity of bulk of okra seeds was 49.1%, which varied between 43.9% and 55.0%. The average weight of 1000 okra seeds varied between 53.2 g and 57 g with an average value of 55.16 g.

Keywords: Okra, seed properties of okra, roundness, sphericity of seeds, density.

Introduction

Okra, commonly known as lady finger, is one of the important vegetable crops in India. In world scenario, it occupies first position in okra production (65.94% of total production) followed by Nigeria [1]. It was sown in 5.04 lacs hectares in India during 2014-15 resulting into a production of 5.7 million metric tonnes [2]. Okra is consumed in form of its immature finger-like green pods. These green pods are the important constituents of diet in developing countries. Its major nutrients are 2.2% protein, 9.7% carbohydrate and 1.0% fibre [3]. It is also rich source of vitamin C (30 mg/100 g), calcium (90 mg/100 g) and iron (1.5 mg/100 g) content [4]. The stems and roots of this vegetable are used to clean cane-juice while making jaggery [5]. Because of its high mucilage content, it is beneficial in curing peptic ulcer, reducing the pains and haemorrhoid effects [6]. In an experiment, its mucilage had been used as alternative to blood plasma [7].

The physical properties of okra is the primary data set required for development of its metering mechanism. This data can also be used for designing of its handling, processing and storage structures. Separation of unwanted material from seeds is easy through oscillating chaffers when size, shape and density of the seed are known [8]. The angle of repose affect the design of mass flow structures. In drying and aeration systems for seeds, bulk density and porosity play a very important role as these properties control the amount of hindrance caused to air flow [9]. This study was aimed to assess physical properties of okra seeds, which includes size, shape, bulk density, and true density, angle of repose, porosity and weight of one thousand seeds.

Materials and Methods

Sample preparation

The okra seed was procured from local market. The variety of the okra seed was *Varsha Uphar*. The sample seeds were cleaned from foreign material, damaged seeds and impurities by manual picking and then passing through a metal screen having square pores of size 10mm. The moisture content of



the seed sample was determined by oven drying method. Three samples of seed taken in aluminum boxes were weighted and placed in hot air oven at 105°C [10]. The moisture content of the seeds was found to be 6.58% on dry basis.

Determination of physical properties

Seed properties essential for development of the metering, processing and storage systems, were identified and determined. The properties identified are discussed below-

Size of okra seed

Seed size is the one of the basic property of seed that is of primary requirement for any seed handling system. The cell size of metering system also depends on it. The seed size is specified by its length, width and thickness. The measurements in these three dimension were taken by using a micrometer having a least count of 0.1 mm. The dimension of 10 randomly selected seeds were measured.

Shape

The shape of seed is important design parameter of flow structures and metering mechanisms. The shape of the seed is expressed by its roundness and sphericity. Roundness of seed, R_p is calculated by formula [11]:

$$R_p = \frac{A_p}{A_c} \times 100$$

where, R_p = Roundness, %

A_p = Projected area, mm²

A_c = Area of the smallest circumscribing circle, mm²

The projected area of the seed was measured using a testing setup which included a thirteen megapixel camera, glass slab, camera stand and graph paper. Experimental setup used for this is depicted in Fig. 1.




Fig. 1: Schematics of experimental setup for finding projection of seed on graph paper

Image of the seed is then processed with the help of computer and projection of seed on graph is created (Fig. 2). Graph paper acts as reference scale in the image taken. The projected area is calculated by graphical method. The area of smallest circumscribing circle is calculated by taking the largest axial dimension of the seed at natural rest position as the diameter of the circle. The procedure

was repeated for 10 seeds selected randomly. The mean was taken as the characteristic value of roundness.




Fig. 2: Projection of okra seed on graph paper

The sphericity of seeds, ϕ is calculated by using the relationship [11]:

$$\phi (\%) = \frac{(\frac{3}{4}) \pi W^2 T}{\pi L^2}$$

where, L = Length of seed, mm

W = Width of seed, mm

T = Thickness of seed, mm

Assuming that volume of solid is equal to the volume of tri-axial ellipsoid with intercepts a, b, c and that the diameter of circumscribed sphere is longest intercept of the ellipsoid,

Length of seed, a = Longest intercept, mm

Width of seed, b = Longest intercept normal to a, mm

Thickness of seed = Longest intercept normal to both a and b, simultaneously

The procedure was repeated for ten seeds selected randomly. The mean was taken as the characteristic value of sphericity.

Bulk density

Bulk density of the seeds is used for seed storage structures designs. Bulk density of seed was measured using an aluminum box having a volume of 120 cm^3 . The box was filled with okra seeds without compaction and then its weight was measured. The bulk density was calculated as follows:

$$\text{Bulk Density } \left(\frac{\text{g}}{\text{cm}^3} \right) = \frac{\text{Weight of seed sample (g)}}{\text{Volume of sample box (cm}^3\text{)}}$$

The procedure was repeated ten times and the average bulk density of the seed was reported.

Volume and true density

The volume of the seed plays an important role in cell design of seed metering mechanisms. The volume and true density of seed are determined by toluene (C_7H_8) displacement method. Toluene was used instead of water because it is absorbed by seeds to a lesser extent. The volume of toluene displaced was found by immersing a weighted quantity of okra seeds in in a graduated glass jar containing known volume of toluene [12]. True density is the ratio of weight of the sample to the volume of the toluene displaced. Observations were taken for ten samples and the mean was calculated separately for volume and true density of seeds.

97 **Angle of repose**

98 The angle of repose is the angle between the base and the slope of cone formed on a free vertical fall
99 of the grain mass to a horizontal plane [13]. The slope of base of the seed flow structures is based on
100 the average angle of repose of seeds to ensure free flow of seed [14]. Three readings were taken with
101 the help of protractor and average was reported.

102 **Porosity**

103 The porosity of seed is calculated using the following expression [13]:

$$\text{Porosity (\%)} = \left(1 - \frac{\text{Bulk density}}{\text{True Density}}\right) \times 100$$

104 Bulk density and true density values obtained from previous experiments were used to calculate the
105 per cent porosity of the seed.

106 **Weight of 1000 seed**

107 The weight of single seed was determined by randomly selecting 1000 seeds of okra and then
108 weighing them one by one. A precision weighing balance having least count of 0.001 g was used.

109 **Results & Discussion**

110 **1. Bulk density**

111 Bulk density of okra seed varied between 0.51 g cm⁻³ and 0.57 g cm⁻³ at seed moisture content of
112 6.58% on dry basis. The average value of bulk density of the seed was 0.54 g cm⁻³. The coefficient of
113 variation of bulk density was 4.55%. Kushwaha et al. [15] reported bulk density of okra seeds as 0.58
114 g cm⁻³ at 11.2% moisture content (d.b.) of okra seed whereas Sahoo and Srivastava [16] reported it as
115 0.59 g cm⁻³ at 8.16% moisture content on dry basis.

116 **2. Volume and true density**

117 The mean volume of okra seeds was 108 mm³ and it varied in range from 99 mm³ to 110 mm³. The
118 mean true density of okra seed was 1.07 g cm⁻³ which ranged between 1 g cm⁻³ and 1.07 g cm⁻³. The
119 coefficient of variation of true density was 5.14%. Kushwaha et al. [15] reported true density of okra
120 as 1.2 g cm⁻³ at 11.2% moisture content (d.b.). At 8.16% moisture content on dry basis, Sahoo and
121 Srivastava [16] reported 1.10 g cm⁻³ as true density of the okra seeds.

122 **3. Angle of repose**

123 The average angle of repose for okra seeds was found to be 28.65° which ranged from 27° to 30° with
124 coefficient of variation as 3.08%. Kushwaha et al. [15] reported angle of repose of okra seed as 28.70°
125 at the 11.2% moisture content (d.b.) whereas Sahoo and Srivastava [16] reported 27.60° as angle of
126 repose of okra seed at 8.16% moisture content (d.b.).

127 **4. Length, width, thickness and geometrical mean diameter**

128 The dimension of okra seed were taken with the help of a micrometer. The average values of length,
129 width and thickness of the seed were found to be 5.73 mm, 4.83 mm and 4.49 mm, respectively (Fig.
130 3). The average geometrical mean diameter of the seed was 4.98 varying between 4.84 to 5.09 with
131 coefficient of variation as 1.55% (Fig. 3). Sahoo and Srivastava [16] reported average geometrical

mean diameter of the seed as 4.98 and the average length, breadth and thickness of the seed as 5.92 mm, 4.71 mm and 4.59 mm, respectively at the 8.16% moisture content (dry basis) of okra seed. Kushwaha et al. [15] reported geometrical mean diameter of okra seed as 4.9 mm.




Fig. 3: Size variation of okra seeds

5. Shape of the okra seed



Shape of seed is represented by its roundness and sphericity. The average value of roundness of the seed was 78.2% ranging from 68.4% to 89.5% with coefficient of variation as 9.03%. The mean sphericity of the seed was recorded as 87.9%, which ranged between 82.4% and 95.8% with coefficient of variation as 5.10%. Sahoo and Srivastava [16] reported the roundness of okra seed as 77.7% and sphericity of okra seed as 74.4% at 8.16 % moisture content (d.b.) whereas Kushwaha et al. [15] reported roundness and sphericity as 79.3% and 86.4%, respectively.




Fig. 4: Shape variation of okra seeds

6. Porosity



The mean value of porosity of okra seed was 49.1% varying between 43.9% and 55.0% with coefficient of variation as 6.79%. Kushwaha et al. [15] found porosity of okra seed as 49.1%. Sahoo and Srivastava [16] reported porosity of okra seed as 46.3%.

7. Weight of 1000 seeds



The average weight of 1000 okra seeds varied from 53.2 g to 57 g with an average value of 55.16 g. The coefficient of variation of weight of 1000 seeds of okra seed was 2.86%. Sahoo and Srivastava [16] reported weight of 1000 okra seeds as 65.78 g.

Conclusion

The study of experiments conducted on okra seeds (*Varsha Uphar*) at 6.58% moisture content on dry basis revealed the following conclusions:

1. The length, width, thickness and geometrical diameter of okra seed were 5.73 mm, 4.83 mm, 4.49 mm and 4.98 mm, respectively.
2. The bulk density and true density of okra seeds was 0.54 g cm^{-3} and 1.07 g cm^{-3} , respectively. The volume of the okra seed was found to be 108 mm^{-3} . The average angle of repose of okra seeds was 28.65° varying between 27° to 30° .
3. The study also concluded that the roundness of the seed was 78.2% ranging from 68.4% to 89.5% whereas the sphericity of the seed was 87.9% ranging from 82.4% to 95.8%.
4. The porosity of okra seed was 49.1% varying between 43.9%-55.0%. The average weight of 1000 seeds varied between 53.2 g and 57 g with average a value of 55.16 g.

References

- [1]. Anonymous. FAOSTAT. Statistics Division, Food and Agriculture Organization of the United Nations, Viale delle Terme di Caracalla, Rome, Italy; 2013.
- [2]. Anonymous. Indian Horticulture Database 2014. National Horticulture Board. Ministry of Agriculture, Government of India; 2015.
- [3]. Saifullah M, Rabbani MG. Evaluation and characterization of okra [*Abelmoschus esculentus* (L.) Moench] genotypes. SAARC J. Agric. 2009; 7: 92-99.
- [4]. Pal BP, Singh HB, Swarup V. Taxonomic relationship and breeding possibilities of species of okra (*Abelmoschus esculentus*). Bt. Gaz.; 1952; 113: 455-464.
- [5]. Chauhan DVS. Vegetable production in India. 3rd Ed., Ram Prasad and Sons, Agra; 1972.
- [6]. Singh N, Singh DK, Pandey P, Panchbhैया A, Rawat M. Correlation and Path Coefficient Studies in Okra [*Abelmoschus esculentus* (L.) Moench]. Int. J. Curr. Microbiol. App. Sci. 2017; 6(7):1096-1101.
- [7]. Siemonsma, JS, Kouame C. Vegetable plant resource of Tropical Africa 2. PROTA Foundation Wageningen. 2004; 21-22.
- [8]. Scherer R, Kutzbach HD. Mechanische Eigenschaften von Koerner-fruechten [Mechanical properties of grains]. Grundlagen der Landtechnik. 1978; 28(1):6-12.
- [9]. Zewdu AD, Solomon WK. Moisture-Dependent Physical Properties of Tef Seed. Biosystems Engineering. 2006; 96(1):57-63.
- [10]. ISI. Indian standard method for analysis of oilgrains. IS: 3579. Indian Standards Institute, New Delhi; 1966.
- [11]. Mohsenin NN. Physical Properties of Plant and Animal Materials (2nd edn.). Gordon and Breach Science Publications, New York; 1986.
- [12]. Tavakkoli H, Rajabipour A, Mohtasebi SS. Moisture dependent some engineering properties of soybean grains. Agricultural Engineering International. 2009; 11: 1110.
- [13]. Chakraverty A. Handbook of Postharvest Technology: Cereals, Fruits, Vegetables, Tea, and Spices. CRC Press, Boca Raton, United States of America; 1981.

- 193 [14]. Jayan PR, Kumar VJF. Planter design in relation to the physical properties of seeds. Journal of
194 Tropical Agriculture. 2004; **42** (1-2):69-71.
- 195 [15]. Kushwaha HL, Srivastava AP, Singh H. A study on physical properties of okra pod and seed.
196 Journal of Agricultural Engineering. 2007; **44**:88-91.
- 197 [16]. Sahoo PK, Srivastva AP. PH-Postharvest Technology: Physical Properties of Okra Seed.
198 Biosystems Engineering. 2002; **83**(4):441-448.