3

4

5

6

## Short Research Article

## The degradation the river Nakanbé banks (Burkina Faso) facing to climate change : a participatory approach for reforestation as a sustainable solution for water resources management

#### 7 8

ABSTRACT

The reforestation of the river banks is a form of resilience of populations facing climate change, soils erosion, silting and siltation of rivers. In its participatory form, the reforestation highlights the primordial role of the riverside populations in the management of water resources. The present study made it possible.

This paper evaluates the soil and vegetation parameters indicating the stabilization of the rivers banks. The methodology is based field description and characterization of vegetation and soils depending the oldness of reforestation.

This study shows that the participatory reforestation approach for the reconstruction of degraded vegetation and soils, reduce silting of the banks and it contribute to fix the banks. However, the participatory approach remains limited by the lack of plants and the accessibility of the plants in the villages for reforestation ; in addition there is a still incomplete mastery of the technique of river banks' reforestation.

### 9

Keywords: banks' degradation; reforestation; Water resources management, participatory approach,
 Nakanbé Basin, Burkina Faso

12

### 13 **1. INTRODUCTION**

14 Since a few decades, precarious climatic conditions have forced farmers to exploit the shallows soils in a context of increasing land pressure, which makes an unsuitable agriculture still largely dominant [1]; [2]. 15 This strong anthropogenic pressure, without respect for the statutory texts, has its corollary, the 16 17 degradation of the rivers banks by (1) the occupation of rivers banks along the watercourse for agricultural and pastoral farming; (2) the anarchic wood harvesting which contributes to the degradation 18 19 of the vegetation cover of the banks; (3) because of the lack of woody plants cover in the banks, water 20 and wind erosion are the fundamental causes of the silting up of water reservoirs and water courses. The 21 silting of water courses fills the water courses and reduces little by little the volumes of the reservoirs and 22 its potential for uses and the survival of ecosystems [3]; (4) pollution by fertilizers (NPK and urea) and

23 pesticides used in agriculture have a negative impact on water guality [4]; [5].

This situation is worrying in some places, particularly at Bam, Dem lakes (Figure 1), where farmers are seeing the progressive reduction of water disponibility and at the same time, the emergence of invasive plants indicative of bad chemical pollution of water [6].

In order to ensure better banks' protection for sustainable management of water resources, projects such
as The Project for Sustainable Management of Forest Resources and the Water Governance
Improvement Project in the Volta Basin ), have included their program of interventions as part of a
concerted and participatory action plan for the protection of shorelines by riverside populations [7]; [8].

31 This study was carried out through a study project entitled "Agro-sylvo-pastoral practices for adaptation to

32 climate change in the Nakanbé basin: effects and impacts on soils and water resources in a context of

33 Integrated Water Resources Management ". The main objective of this study was to evaluate the impact

34 of the participatory approach of reforestation of banks on the stabilization of degraded banks in the

35 context of climate change. In order to achieve our objectives, we hypothesized that the reforestation of 36 the banks allowed to improve the fixation of the banks by the regeneration of the vegetation. 37

#### 38 2. MATERIALS AND METHODS

39 40

### 2.1 Study framework

41 The area of this study is located in the village of Sampéma (Boulgou province) with the following 42 coordinates: 11°17' North and 0°33' West (Figure 1). The climate is of North-Sudanese type one. The 43 average annual rainfall is 900 mm. The winter period is between July and October. This period is followed 44 by a dry season from November to June ([9]).

45

46 The average minimum temperature is 15°C (from december to february) while the average maximum 47 temperature is 34.9°C (april, may). The soils along the banks are hydromorphic soils.

48

49 According to the division of [9], the vegetation and agricultural practice, Sampema belongs to the northern 50 Sudanian phytogeographical zone. The wooded savannah is the bank vegetation species due to the process of degradation along the rivers. Some gallery forests species grow along the main rivers and are 51 composed of Borassus aethiopium, Terminalia sp, Anogeissus leocarpus, Khaya senegalensis, Daniella 52 oliveri, Anogeissus leiocarpus, Diospyros mespiliformis, Pterocarpus erinaceus, Parkia biglobosa, 53 54 Tamarindus indica. The grass carpet is quite large and composed of species such as Andropogon 55 gayanus, Schoenefeldia gracilis and Cymbopogon schoenanthus.

#### 56 57 2.2. Collection of data

#### 58 2.2.1 Placement of plots

59

60 For the placement of the plots, we first delimited the plots according to their age using man hand GPS. 61 Using the "Mapsource" software, we then realized the plots and the sampling plots on the computer. We 62 initially limited blocks of 75mx30m surface along the water course. Three reforested blocks of 6; 4 and 3 63 years old were selected. Within each block, we made three plots along the slope. For each block, the first plot (top of the slope) of 10mx30m is reforested with Azadirachta indica or Senna siamea, the second 64 65 (mean of the slope) of 30mx30m in Mangifera indica and the last (down of slope) of 35mx30m in 66 Sarcocephalus latifolia (Figure 1). 67

### 70 Figure 1: Location of Sampéma site 71

### 72 2.2.2 Soil sampling

In order to analyze the fertility of the reforested banks, four samples were taken using an auger according to the diagonal of each plot and according to the depths 0-10, 10-20, and 20-30cm. From the four samples of the same depth along the diagonal, we mixed to obtain a composite sample, including three composite samples per plot. The samples were dried in the shade during 14 days.

# 782.2.3 Inventory of vegetation79

The woody plants inventory has consisted of identifying the different species and their numbers,
measuring the diameter of the trunk at the base and the total height of the plant.

### 83 2.2.4 Participatory approach

A field survey was possible to know that this protection of the banks by reforestation was possible with a
participatory approach.

### 88 2.3 Data processing

89

The data collected was entered using the Excel spreadsheet. The analysis of the variance was made using the GenStat 4 software. The comparison of the averages was made with the Student Newman Keuls test at the error threshold of 5%.

### 94 3. RESULTS AND DISCUSSION

95

93

## 96 3.1. Wood density of reforested banks

97

98 The analysis of the of the number of woody species by age of reforestation (Figure 2) shows, in high and 99 medium slope, the number of plants shows non-significant growth with the age of reforestation ; at the 100 bottom of the slope, the 4-year-old reforested bank has the most significant tree population (0.075 101 individuals/m<sup>2</sup>).

102



103 104

107

### 105 Figure 2: Ratio of woody species according to the oldness of reforested banks

106 TS = top of slope; MS = mean of slope and DS = down of slope

The analysis of the number of woody species by age of reforestation (Figure 2) show, in high and medium slope, non-significant growth with the age of reforestation ; at the bottom of the slope, the 4-year-old reforested bank have the most significant tree population (0.075 individuals/m<sup>2</sup>).

111

## 112 **3.2.** Richness and specific contribution of the woody species of the reforested river banks

113

114 The analysis of the richness and specific contribution of the woody species of the reforested river banks 115 (Table 1) shows that :

the 3-year-old reforested banks have the most number of species, six (6), with significantly more
 *Sarcocephalus latifolia* (22 plants) followed by *Senna siamea* (12 plants) and Mimosa pigra (10 plants).
 The 4-year-old reforested banks have 5 species with a significant number of *Mimosa pigra* (49 plants)

119 and Sarcocephalus latifolia (29 plants).

the 6-year-old banks have four (4) species with Sarcocephalus latifolia (26 plants), which is significantly
 larger followed by Azadirachta indica and Mimosa pigra (15 plants each). Senna siamea, Ficus
 gnaphalocarpa and Eucalyptus camaldulensis are present only in 3-year old reforested banks while
 Vitellaria paradoxa is only present on the 4-year-old banks.

124 - the analysis of the vegetation distribution shows that *Sarcocephalus latifolia* is the dominant species

125 followed by *Mimosa pigra* and *Azadirachta indica*.

# 126127 Table 1: Frequency of woody species by age of reforested banks

	6 years old bank	4 years old bank	3 years old bank	Probability
Eucalyptus c.	0	0	0. 67	0,079
Ficus g.	0	0	0. 33	0,42
Manguifera I.	9a	6b	0c	<,001
Mimosa pigra	15b	49a	10b	<,001
Sarcocephalus I.	26a	29a	22b	0,007
Senna siamea	0b	0b	12a	<,001
Azadirachta i.	15a	13.67b	1c	<,001
Vitellaria p.	0	0. 33	0	0,422

128

129 Student-Newman-Keuls test; Threshold 5%; values are averages of 3 repetitions;

130 Values followed by the same letter in the same line do not differ significantly at the 5% probability threshold

131 Eucalyptus c.: Eucalyptus camaldulensis; Ficus g .: Ficus gnaphalocarpa; Manguifera I.: Manguifera indica ;

132 Sarcocephalus I. : Sarcocephalus latifolia ; Azadirachta i. : Azadirachta indica; Vitellaria p. : Vitellaria paradoxa.

133

### 134 **3.3. Distribution of wood plants by height and circumference**

The analysis of the distribution of woody plants by height and circumference (Table 2) shows that the differences are significant. The reforested 4-year-old banks show the plants with the greatest circumference and height. The plants of the reforested banks of 6 years old and those of 3 years old do not show a significant difference between them despite the dimensions of the plants higher in the level of the replanted banks of 6 years old.

140

### 141 Table 2 : Height and circumference of woody plants accordind to the age of reforested banks

	6 years old	4 years old	3 years old	Probability
Circumference (cm)	21. 67ab	33. 33a	15. 67b	0, 030
Height (m)	13b	20a	9. 47b	0, 002

142

143 Student-Newman-Keuls test; Threshold 5%; Values are averages of 3 repetitions; values followed by the same letter 144 in the same line do not differ significantly at the 5% probability threshold

145

### 146 3.4. Sieve analysis

147

148 The results of the Sieve analysis are reported on Table 3. We can show a significant difference between 149 fine and coarse elements.

At the bottom of the slope, the silt contents show a significant increase and go from 24.84% (reforested banks of 3 years) to 37.25% (reforested banks of 6 years old). At the top of the slope, the silt content decreases from 43.79% (reforested banks of 3 years old) to 37.90% (reforested banks of 6 years old).
However, at the top of the slope, the difference is not significant between the reforested banks of 6 years

and 4 years old. On mean slopes no significant differences are observed. The bottoms of slope are less silty than high and medium slopes.

The sand contents decrease from the bottom of the slope to the top of the slope, although there are no significant differences, apart from the reforested bottoms of slopes of 4 and 3 years old. At the bottom of the slopes, the sand content decreases with time and goes from 63.40% (reforested banks of 3 years old)

to 49.02% (reforested banks of 6 years old). In high and medium slopes, the sand contents do not show any significant differences.

161 - The clay contents increase significantly with the age of the development. This increase can also be seen

162 in low and top and medium slopes.

### 164 **Table 3: Granulometric composition of the soils of the reforested banks**

### 165

	Clay (%)	Silts (%)	Sands (%)
TS 6 years	17.00a	37.90c	45.10c
MS 6 years DS6 years TS 4 years	17.00a 13.73c 14.38bc	37.90c 37.25c 38.56b	45.10c 49.02c 47.06c
MS4 years DS4 years	15.04bc 12.42cd	38.56b 31.37d	46.41c 56.21b
TS 3 years	12.42cd	43.79a	43.79c
MS3 years	15.69b	33.98d	50.33b
DS3 years Probability	11.76d <.001	24.84e <.001	63.40a <.001

166

167 Student-Newman-Keuls test; Threshold 5%; values are averages of 3 repetitions;

168 Values followed by the same letter in the same column do not differ significantly at the 5% probability threshold; TS =169 top of slope; MS = mean of slope and DS = down of slope

170

172

175

### 171 **3.5. Chemical parameters of soils**

173 The statistical analysis of the chemical parameters of reforested soils (Table 4) shows significant 174 differences between the contents of the chemical elements.

Between the tops of the slopes of the reforested banks of 3 years and those of 6 years, nitrogen levels increased from 0.0283 meq/100g to 0.045 meq/100g. Carbon levels increased from 0.343 mg/g to 0.575 mg/g. The C/N ratio increases from 12 to 13. The available potassium levels increase from 54.12 meq/100g to 68.88 meq/100g. On average and low slope their evolutions are neither function of the time nor the topography.

181

185

182 On the top of slope (TS), water pH values did not differ significantly, except for the 4-year (6.607) and 3-183 year (6.587). The water pH shows a downward trend at the top of the slope. On the mean of slope, the 184 water pH shows a non-significant increase.

186 In high and low slopes, assimilable phosphorus levels are significantly lower in the reforested banks of 3 187 years old and higher in those of 4 years old.

188 The mean slopes have the highest values, while the down of slopes have the lowest ones.

189

### 190 Table 4: Chemical characteristics of reforested soils

191

	Total Carbon (g/kg)	Total nitrogen (g/kg)	C/N	Total organic matter (g/kg)	Water pH	Available phosphorus (meq/100g)	Available Potassium (meq/100g
TS 6 years old	0.575a	0.045a	13.00a	0.992a	6.20b	10.867b	68.88b
MS 6 years old	0.575a	0.045a	13.00a	0.992a	6.20b	10.867b	68.88b

DS E vers eld	0.337c	0.028cd	12.33bc	0.581c	6.13b	7.473c	53.79bc
6 years old							
4 years old	0.405c	0.033c	12.33bc	0.698c	6.61a	12.323a	5937bc
MS							
4 years old	0.450c	0.035bc	12.67b	0.776c	6.12b	10.093b	84.62a
DS							
4 years TS	0,359c	0,028c	12,67b	0,618c	6,03c	8,637c	43,62c
3 years	0,343c	0,028c	12,00c	0,592c	6,59b	4,620d	54,12bc
MS							
3 years	0,567b	0,044b	13,00a	0,978b	6,09b	11,253b	84,95a
DS							
3 years	0,204d	0,019d	11,00d	0,352d	6,22b	5,710cd	53,46bc
Probability	<,001	<,001	<,001	<,001	<,001	<,001	<,001

193 Student-Newman-Keuls test; Threshold 5%; values are averages of 3 repetitions;

values followed by the same letter in the same column do not differ significantly at the 5% probability threshold; TS = Tap of closes MS. Mean of closes and DS. Down of closes

195 Top of slope; MS = Mean of slope and DS = Down of slope 196

### 197 4. DISCUSSION

198

The participatory approach of the reforestation of the banks allowed the revegetation of the banks once degraded and deforested. The reconstruction of the vegetation is more important at the level of the riparian formation. Indeed reforestation was initially done with plants of *Sarcocephalus latifolia* but one finds after a strong population of wild *Mimosa pigra*. The authors [10] and [11] argue that riverbank development improves the regeneration of plant species by trapping seeds and seedlings transported by waterflow and runoff.

The analysis of the specific richness showed that the dominant species are *Nauclea latifolia* and *Mimosa pigra*. These species are typical to flood-prone soils. Aquatic plants by the phenomenon of hydrochory, let the flow disseminate their seeds. It is a common strategy in these plants, such as *Sparganium emersum*, *Nelumbo nucifera*, *Nymphea alba*, *Nuphar luteum* or *Mimosa pigra* [12].

210

In the high and medium slopes there are no species from post-reforestation regeneration despite the large quantities of seeds produced by *Azadirachta indica* and *Senna siamea*. Similarly, there is no regeneration of species other than those reforested or already present on the site before reforestation. This finding shows that assisted natural regeneration is not practiced in high and medium slopes unlike low slopes.

216

217 The small number observed on medium slope of the banks of 4 years (picture 1, appendix) is due to the 218 2007 flood which destroyed almost all the plants planted in 2006 (picture 2, appendix). This flooding could also be the cause of the growth retardation observed in the plants of the 6-year-old reforested banks 219 220 compared to those of the 4-year-old reforested banks. Indeed, the excess of water creates anaerobic 221 conditions that alter the root metabolic functions and can lead to poisoning and cause the deterioration of 222 the physical properties of the soil and make them more vulnerable to degradation. The oxygen required 223 for root breathing, circulates in the interstices of the ground. In case of flood, water occupies all these 224 pores and the roots are placed in an asphyxic condition, which inhibits their growth. Plants can survive 225 short periods of anaerobic condition (up to 0.5% oxygen). Roots require 6-8% oxygen to grow normally. 226 Under these values the leaves become chlorotic, the growth is interrupted, the formation of new roots 227 stops, the stems become dry and the plant dies. Flooding can also reduce sweating by almost 90%, 228 limiting the absorption of water and nutrients and reducing photosynthesis. These anaerobic conditions at

the soil level are combined during insufficient drainage of heavy soils during heavy rains and flooding[13].

231

The mean slopes of the reforested 3-year-old banks are a fallow not replanted by lack of mango trees (Picture 3, appendix), hence the number of plants very low compared to other parts of the banks.

On the soil level, the decline in silt contents at the top of the slope, as well as their increases at the bottom of the slope, show that there is leaching of the silts from the slopes to the bottom of the slope. The higher grades of sand at the bottom of the slope reflect a silting phenomenon. The slope intervenes in the degradation of the land by its importance (declivity), its length and its form; the soil losses increase exponentially with the value of the slope [14]. The increase in the degree of slope results in increased runoff [15].

The low slope plots are in the flood plain of the Nakanbé, where the phenomenon of sandy alluvial deposits is more important (Picture 4, Annex). However, the decline over time of the sand contents at the bottom of the slope shows that reforestation allowed the retreat of silting. Similarly, the increase in clay contents over time indicates that reforestation has increased the fine texture of soils. Clay minerals make the soils heavier and therefore improves the stability of the banks.

247

The other aspect of this study is the participatory approach that brought together all stakeholders from the beginning to agree on the terms of the action so that it is mutually beneficial for all stakeholders. Thus, land owners benefit from the fruit plants offered by the project and firewood that will constitute *Nauclea latifolia*. The project and the Nakanbé water agency benefit from the protection of the bank by fixing species. This approach is a real innovation in that it makes a consensus for the protection of the banks.

# 254 **5. CONCLUSION** 255

256 At the end of this study, we can draw the following conclusions:

The reforestation of the rivers banks initiated in recent years in Sahelian country of Burkina Faso is
beginning to prove this efficiency in the restoration of degraded banks. It accelerates the regeneration of
vegetation, the restoration of soil fertility and texture.

261

• The reforestation of the river banks remains however limited by the availability of the plants for reforestation in the areas and internal villages who need it.

264 and

• Reforestation of the banks also causes a decrease in water pH and have negative effect of soils quality. 266

• These techniques of reforestation would not have any positive impact on regeneration of vegetation without the practice of the assisted natural regeneration.

### 269 270 RECOMMENDATIONS AND PERSPECTIVES

Based on the findings of this study, we propose the following recommendations and perspectives for
 better optimization of participatory reforestation and better management of water, soil and mitigation of
 the effects of climate change in the Nakanbé watershed :

- 274
- Extend the reforestation of the banks in the entire Nakanbé watershed,
- Associate stony barriers and / or grass strips with the reforestation of the banks to increase their durability and efficiency,
- Promote and prioritize participatory environmental management approaches,
- Sensitize people on the practice of the natural assisted regeneration.
- 280 281

ANNEX 282



Photo 1: Area destroyed by the 2007 flood, 4 years after



Photo 3: Average slope of banks of 3 years old





### <u>Photo 2</u>: Result of the flooding of the river bank in 2007 (Source: [15]IUCN)

Photo 4: sandy bank

### 287 **REFERENCES**

288

[1] Albergel J, Lamachère J. Agricultural development of lowlands in the Sahel:
 typology, hydrological functioning, agricultural potential. Final report of a CORAF-R3s
 project, CIRAD, Montpellier, France. 1993, p335 French

[2] DRED / CS, April 2007. Monograph of the province of Bazèga. RGPH / CS. 109P.
 French.

[3] Gouem A. Inventory of the degradation of the banks of Lake Bagré and feasibility
analysis of the management options proposed by the different stakeholder groups ",
IPR / IFRA, 2007, 96p. French

[4] Somé K, Dembélé Y, Somé L and Millogo J. Water pollution in the Nakanbé basin:
the case of Loumbila and Mogtédo reservoirs in Burkina Faso. Rev. Southern Science
and Technology, 2008, No. 16, pp 14-22.

[5] Millogo D, Bazié MM, Koussoubé Y, Gnakambary Z, Zombré PN, Da ECD.
Assessment of agricultural and mining pollutions of waterbodies within the Nakanbé
basin (Burkina Faso): the case of the Goinré, Ziga and Bagré reservoirs. Journal of
Water Resources and Protection (JWARP), in Press

304 [6] Directorate General of Water Resources (DGRE). State of play of the water
 305 resources of the Nakanbé basin. Final Report, Ministry of Agriculture, Hydraulics and
 306 Fisheries Resources, Burkina Faso. 2010, 290p. French

307 [7] Thiombiano LJ. Contribution to the implementation of the concerted action plan for
 308 the protection of the banks of the Bagré dam lake. Institute of Rural Development,
 309 2009, 73p. French

[8] International Union Against Nature (IUCN). Annual Report PAGEV, 2006, 40p.French

[9] Fontes J and Guinko S. Map of the vegetation and land use of Burkina Faso,
National Center for Scientific Research University of Toulouse III, Institute of Rural
Development, Faculty of Science and Technology of the University of Ouagadougou ,
Ministry of French Cooperation, Campus Project 88313101. 1995. French

- [10] Bernard L. Guide for the protection of stream banks in plant techniques. Ministry of
   Regional Planning and Environment, 1994, 30p. French
- [11] Barbry Y., Application of plant techniques for the protection of riverbanks.
   Waterways of France, Guide of application of vegetable techniques. 200,. 24p. French
- [12] Lanzara P. The unknown universe of plants, Brussels, Elsevier-Sequoia, coll.
   "Multiguide Nature", ISBN 2-8003-0203-8, 1977, 254p. French
- [13] Heller R, Esnault R, Lance C. Physiologie Végétale Tome I. Nutrition.Edition
   DUNOD. 2004 323p. French
- [14] Cisse A and Touré IA. Conservation of the environment and natural resources in
   the Sahel. CILSS. Bamako, Mali. 1991, 140 p. French

- [15] Combination A. Erosion and soil conservation (especially in the tropics). Course
- 326 327 328 plan commented and illustrated examples for students of ORSTOM and students of the Postgraduate Course for the management of the natural environment. 1977, 85p.
- 329 French