

# NUTRITIONAL AND SENSORY PROPERTY OF CHIPS MADE FROM POTATO (S. t).

**Abstract:** The purpose of this paper was to summarize nutritional and sensory property of chips made from potato (*Solanum tuberosum* L.). Potatoes have an important role in human nutrition. It is a source of carbohydrates, protein, vitamins and minerals in the diets. Tuber quality is an important attribute for processing, and should be enhanced for increasing farmers' income and livelihood. The acceptability of potatoes for processing as chips is largely dependent on the quality of the end products. Blanching improves chips color, texture, sweetness, and reduce sourness and bitterness thereby increasing its overall acceptability. French fries and potato crisps are the most consumed processed potato products in large cities. High specific gravity is a sign of that the raw potatoes will produce high chip volume due to high dry matter content and has crispy texture. Tuber with low reducing sugar produce light color chips and is more acceptable by consumers.

**Key word:** chips, potato, dry matter, specific gravity.

## INTRODUCTION

Potato is a member of the family Solanaceae and the genus *Solanum* which is one of the most productive and widely grown horticultural food crops in the world. Potato (*Solanum tuberosum* L.) is the fourth most important food crop in the world, after wheat, maize, and rice [1, 2]. The crop plays a significant role in human nutrition worldwide, where more than 320 million tons of potatoes are produced annually on 20 million hectares of land [3, 4]. The worldwide production of potato exceeded 330 million tons/year. Potato contains dietary antioxidants, which may play a part in preventing diseases related to ageing [5]. Since potato contains nearly all important vitamins and nutrients, it supports life better than any other crop when consumed as the sole diet [6]. Although, potato is grown for animal feed, industrial uses and for seed tuber production the main use is still as direct food.

Increasing proportion is processed into snack food. Potato is mainly important for subsistence farmers, but it may also be a cash crop [7]. Processing quality of potato tubers for chip production is determined by its high dry matter, and low reducing sugar and phenol contents [8, 9]. High dry matter content increases chip yield, crispy-consistency, and reduces oil absorption during frying [10, 11]. Low reducing sugars and phenol contents are required to avoid dark color and bitter taste of processed products, which negatively affect consumer acceptance [12, 13]. Dry matter of potato tubers and chip color are genetically controlled and influenced by environmental conditions during growing season and storage temperature [14]. One of the factors affecting the quality of processed product of potato is the physicochemical characteristics. For instance, not all the potato varieties will produce high quality potato chips [15, 16]. Studies have shown significant differences in chip quality as a result of varying chemical composition of the potato tubers [17, 18]. The physical and chemical characteristics of the potato tubers vary from one variety to another and within the same variety depending on growing conditions (e.g. soil temperature and soil moisture), harvesting and handling conditions (storage) [17]. The importance of potato in developing countries could be attributed to two main reasons: Firstly, potato crop produces more edible energy and protein per unit area and time than other food crops, and secondly, potato fits well into multiple cropping systems prevalent in tropical and subtropical agro-climatic conditions [19]. This is further supported by [20] indicating that potato area in Sub Saharan African countries has grown as a result of increasingly emerging market oriented farmers in response to huge demand from growing urban markets. Moreover, because of its bulky in nature, it does not normally enter into international trade unlike major cereals. Only a fraction of potato total production enters foreign trade [21]. The crop serves as a buffer to rising food prices, especially cereals [20]. Potatoes are a principal source of carbohydrates and protein, and also contribute some vitamins and minerals in the diets. Potato (*Solanum tuberosum* L.) is an important food security and cash crop in Ethiopia [22]. With increasing urbanization, the use of the crop not only as fresh tubers but also as processed products such as French fries and crisps is rising in the country [23, 24]. Potato provides not only carbohydrate, proteins, and vitamins but also minerals that are important in the diet for human wellbeing [25], and is important for nutritional security. It is noteworthy that potato absorbs

large quantities of plant nutrients, especially nitrogen (N), potassium (K) and phosphorus (P) from the soil [26]. Potassium and N are found in the largest amounts in a potato plant, followed by Calcium (Ca) and Magnesium (Mg) [27]. According to [28], some potatoes are not suitable for the production of processed products due to low dry matter content. It is important for researchers to recommend farmers to use only those varieties that make good quality products both at harvest and during storage periods. The production of lighter color chips acceptable to the market often requires some pretreatment of the sliced potatoes in the processing plants [29]. Prior to drying most food products are usually subjected to one form of pretreatments, among which blanching is one of the most important techniques [29]. Blanching refers to the process of immersion of raw vegetables in a heated fluid (water, oil or acid) for a period of time. Blanching has a leaching effect on the sugars and serves to even out variations of sugar concentrations at or near the surfaces of the French fry strips. This gives a lighter and more uniform color on frying. It is also used to extract the reducing sugars and asparagines from the surface of potatoes in order to reduce browning, particularly the formation of brown spots [30]. Blanching reduced glucose and asparagines content by average of 76 and 68%, respectively in potatoes [10]. Potato is a high yielding tuber crop with a short cropping cycle of about 3-4 months. This coupled with high potential yield of about 40 tons per hectare, makes the potato a suitable crop for places where land is limited and labor is abundant [5] as in sub-Saharan Africa. Potato is a smallholder cash crop of the future with a potential to improve their livelihoods and reduce poverty. Furthermore, Potato can provide a cheap but nutritionally rich staple food, contributing protein (low in amount, but high in biological value), vitamin C, zinc, and iron. Potato offers employment opportunities to all those involved in potato value chain from production to consumption. The objective of this review is to understand and summarize nutritional importance of chips made from potato (*solanum tuberosum* L.).

### **Description of Chips**

Recently, Potato chips have become a heavily marketed snack foods throughout the world. Potato has high contents of protein (high percentage of essential amino acids: lysine, leucine, phenylalanine, threonine and valine), minerals (potassium, phosphorus and magnesium nutrient salts) several vitamins (B1, B3, B5, B6, folate, pantothenic acid, riboflavin) and large amount of vitamin C. Potato is very low in fat but rich in carbohydrate and several micronutrients. Potato also contains dietary antioxidants, which may play a part in preventing diseases related to ageing [5]. Chips is frying product made from potato by using high temperature. Several oils can be used for frying potato chips, including soybean oil, palm oil, and safflower or groundnut oil.

### **Reducing Sugar 96**

The disaccharide sucrose, and the two monosaccharides, the glucose and fructose, are the three major types of sugars in potato tubers [31]. The sugar content of potatoes varies from 0.2 to 0.6% due to the differences in genotypic characteristic among the varieties of potato [32]. The contents of glucose and sucrose are high in young tubers and decrease with plant maturity. Kimondo also described that the maximum contents of sugars were obtained 1-2 weeks after tuber initiation, and then the sugars decreased along with growth of the tubers and reached a minimum before the end of the growing season. The quality of potatoes is dynamic and continues to change as a result of physiological activity owing to accumulation of reducing sugars and depletion of starch [34]. The reducing sugar content is critical in the quality of processed potatoes, as studies have shown that reducing sugar content negatively influences the color and flavor of the finished product due to Maillard reactions that take place during frying [35]. The Maillard reaction, is initiated at high temperatures (> 120°C) due to the condensation of free amino acids and reducing sugars [36]. Among the sugars, the reducing sugars (glucose and fructose) are of most concern, as they are chemically reactive and involved in the non-enzymatic browning reaction thus determining the eventual fry color of the potatoes [37]. The level of reducing sugars that are generally acceptable for processing of chips is 0.2–0.25% and French fries is 0.3–0.5% [38]. The sugar content of the harvested crop is also important for the fresh market, and sucrose levels above 1% fresh weight (FW) are reported to be unacceptable to the consumers due to its sweetness. This sweetening is more likely to occur after a period of storage, particularly at holding temperatures (below 4°C) that promote low-temperature sweetening.

### **Tuber dry matter content and specific gravity**

The dry matter yield of a potato is the major determinant of texture of the raw or cooked product [39]. It is one of the most important factors that contribute to favorable mouth feel [40].

Potato chips processing requires tubers with dry matter content of greater or equal to 20% and specific gravity of greater or equal to 1.080 [38]. Kabira *et al.* reported that potatoes with a dry matter content of 20 to 24% are ideal for making French fries while those with a dry matter content of up to 24% are ideal for preparing crisps. High specific gravity is an indication that the raw potatoes will produce high chip volume due to high dry matter content.

## **Sensory attributes**

### **Color**

Color in processed products such as potato chips can be affected by several factors including product composition and processing conditions [37] that influence consumer acceptability [41]. For instance, common browning of foods during heating occurs when reducing sugars and a free amino acid or amino group react in the Maillard reaction [41]. Marquez *et al.* studied color development during potato frying and found that both reducing sugars and amino acids are involved in the color development of fried potatoes, reducing sugars being the limiting factor. According to [28], hot water blanching at 65-100°C before frying destroys enzyme activity and leaches out, reducing sugars and other chemical constituents that cause off color and off flavor. Blanching treatments are used to reduce browning of fried products by minimize Maillard reactions which play a predominant role in color and acrylamide formation during frying [10]. Blanching lead to lighter in color potato chips than those of the control after frying at 150°C [10]. Rodriguez *et al.* found that reducing sugars had the biggest influence on lightness, producing the brighter colors when these sugars were absent; however, reducing sugars did not entirely predict color quality when present at low concentrations. Color is one of the most important quality factors that need to be considered during frying [43]. In general, a yellowish brown color [44], uniform light golden [45], and lighter colored are preferred in potato chips.

### **Texture**

In general, chips prepared from tubers with higher dry matter have weak structure compared with those prepared from tubers that have lower dry matter. Moyano *et al.* showed that the texture of potato chips was found to be directly related to specific gravity, total solids, starch content, cell size, and surface area and pectin. Crisps obtained from potatoes rich in dry matter (above 25%) can exhibit hard textures, whereas crisps made of tubers with low dry matter content are characterized by greasy and sticky textures. Texture in food products has dominant contribution to the overall quality and acceptability [47, 48].

### **Bitterness**

Asmamaw *et al.* showed that the loss in taste of chips prepared from tubers stored for extended period may due to the increase in the concentration of glykoalkaloids level of tubers. In all cases, blanched samples get better score by the sensory panel. According to [29], blanching and drying temperature significantly affected the hardness of potato chips positively while the drying method did not show any significance influence on the hardness. Krokida *et al.* pointed out that blanching caused starch gelatinization and softening of structure leading to less hardness of dried starchy products. The authors also reported that unblanched chips had the maximum hardness in all of the cases. Pimpaporn *et al.* found that blanching reduces the hardness and shrinkage of the product. While blanching at low temperatures (55 to 75°C) lead to a firm texture [51], blanching at high temperatures (80 to 100°C) for alternatively long period of times (15 min) leads to the loss of firmness [52]. In vegetables, bitterness is considered as prevention for consumption [53].

### **Flavor**

Potato flavor results from the combination of taste, aroma, and texture. Flavor precursors synthesized by the plant are present in raw potatoes and consist mainly of sugars, amino acids, RNA (Ribo nucleic acid), and lipids. Plant genotype, production environment, and storage environment influence the levels of these compounds and the enzymes that react with flavor precursors produce flavor compounds. During

cooking, flavor precursors react to produce the Maillard reaction compounds and the sugar, lipid and RNA degradation products that contribute to flavor [54]. The flavor of the potato chips is more complex than that of boiled, baked or mashed potatoes whenever the cooking temperature is higher, and the absorbed oil contributes to the overall flavor profile of the product [55]. In fried potato products, flavor compounds are not only inherent in the raw potato but also from the frying oil.

Maillard reaction products with lipid oxidation products [56] the complete composition and understanding of fried potato flavour has not been fully established [57]. Variations in flavor exist between varieties although there is little published research [58]. Ereifej *et al.* reported sensory differences in the aroma and taste of potato chips made from different cultivars that varied in level of dry matter, Sugars, amino acids and lipids. It is generally understood that the bulk of flavor producing compounds in raw potatoes are volatile [57]. The major classes of volatile compounds released by raw potatoes are acid, aldehydes, alcohols, amines, esters furans, hydrocarbons, ketones, pyrazines, pyridines and thiazoles [57]. However, [60] suggested that a potato flavour might be influenced by non-volatile compounds such as amino acids and sugars. Browning become very rapid at temperature higher than 150oC and volatile flavor compounds are produced as secondary products [15]. Lisinska *et al* reported that the desirable flavor of potato chips is limited by the high dry matter content as well as the low sugar concentration. The flavor and odour of light colored chips are less intense than those of dark colored chips [48]. Glycoalkaloid found in potatoes could contribute to bitter off flavors of burning sensation at elevated concentration [60].

### **Crispiness**

For potato chips, a very crispy texture is expected since it is an indicator of freshness and high quality [61]. The crispy structure of potato chips is the result of changes at the cellular and sub- cellular levels in the outermost layers of the product. Blanching causes a permanent modification of the cellular structure in the potato tissue [52]. The heat treatment during blanching affects the typical potato cell by altering the cytoplasmic membrane. Heat destroys the differential permeability of the membrane letting water to enter the cells and intercellular spaces there by expelling gases and other volatiles, causing loss of water soluble nutrients (sugars, vitamins, and minerals) [52]. During pre-treatment, changes occur in the cell membranes that play a key role in the changes that occur within the tissue during further processing [62].

### **Overall acceptability**

Andersson *et al.* reported that in the case of potato processing, blanching is used to inactivate peroxides, to improve the texture, color and, to some extent, the flavor of final product. Potato blanching affect the enzyme activity that lead to quality degradation [63].

### **Conclusion**

Potato is one of the staples of the human diet and is an important raw material in the starch industry as well. Thus, potatoes are a very significant part of the diet in many countries and can make a significant contribution to human nutrition. It is one of the basic crops, which significantly impact nutritive status of the world population. Because it contains nearly all important vitamins and nutrients, so that, it support life better than any other crop when eaten as the single diet. It is a major part of the diet of half a billion consumers in the developing countries. Potato is an important food and cash crop in Eastern and Central Africa, playing a major role in national food security and nutrition, poverty alleviation and income generation, and; provides employment in the production, processing and marketing sub-sectors. The protein content of potato is similar to that of cereals and is very high in comparison with other roots and tubers. In addition, the potato is low in fat and rich in several micronutrients. Generally Potato with high specific gravity and dry matter are preferable for chips making and its frying product has crispy texture. The presence of low reducing sugar content makes the cultivars suitable for chips processing. Chips with light color are more acceptable by consumer and have low level of reducing sugar. As the level of glykoalkaloids increases. the level of bitterness increase. Flavor is one of the important quality factors of potato chips and is affected mainly type of due to the oil used to fry chips, flavor compounds inherent in the raw potatoes, and added.

## References

1. Hamideldin, N. and Hussien, O.S. (2013) Morphological, Physiological and Molecular Changes in *Solanum tuberosum* L. in Response to Pre-Sowing Tuber Irradiation by Gamma Rays. *American Journal of food science and Technology*, 1, 36-41.
2. Nairobi.Kaguongo, W.P., Gildemacher, P., Demo, P., Wagoire, W., Kinyae, P., Andrade, J., Forbes, G., Fuglie, K. and Thiele, G. (2008). Farmer Practices and Adoption of Improved Potato Varieties in Kenya and Uganda. International Potato Center (CIP), Lima. SOCIAL Sciences Working Paper 2008-5, 85 p.
3. Poczai, P., Cernák, I., Gorji, A.M., Nagy, S., Taller, J. and Polgár, Z. (2010). Development of Intron Targeting (IT) Markers for Potato and Cross-Species Amplification in *Solanum nigrum* (Solanaceae). *American Journal of Botany*, 97, e142-e145. <http://intl.amjbot.org/http://dx.doi.org/10.3732/ajb.1000360>
4. FAOSTAT, (2011). *Food and Agricultural Organization Statics*. Agricultural database. <http://a.ostat.fao.org>.
5. FAO, (2008a). *Food and Agriculture Organization*. International Year of the Potato 2008 New Light on a Hidden Treasure. Food and Agriculture Organization of the United Nations, Rome.
6. Reader, J. (2008). Propitious Esculent: The Potato in World History (London: William Heinemann).
7. Struik, P.C. and Wiersema, S.G. (1999). *Seed Potato Technology*. Wageningen Pers, Wageningen, the Netherlands. pp383.
8. Kadam, S. S., Wankier, B. N. and Adusule, N. R. (1991). Potato Production, Processing and Products. Boca Raton: CRC's Press. p.35.
9. Abong, G.O., Okoth, M.W., Karuri, G.E., Kabira, N.J., and Mathooko, M. F. (2009). Levels of reducing sugars in eight Kenyan potato cultivars as influenced by stage of maturity and storage conditions. *J. Anim. and Plant Sci.*, 2: 76-84.
10. Pedreschi, F., Moyano, P., Kaack, K. and Granby, K. (2005). Color changes and acrylamide formation in fried potato slices. *Food Research International*, 38, 1-9.
11. Rommens, C.M., Shakya, R., Heap, M. and Fessenden, K. (2010). Tastier and healthier alternatives to French Fries. *Journal of Food Science*, 75:109-115.
12. Wiltshire, J.J.J. and Cobb, A.H. (1996). A review of the physiology of potato tuber dormancy. *Annals of Applied Biology*, 129:553-569.
13. Wang-Pruski, G. and Nowak, J. (2004). Potato after-cooking darkening. *American Journal of Potato Research*, 81:7-16.
14. Kawchuk, L. M., Lynch, D.R., Vada, R.Y., Bizimungu, B. and Lynn, J. (2008). Marker assisted selection of potato clones that process with light chip color. *American Journal of Potato Research* 85:227-231.
15. Miranda, M. L. and Aguilera, J. M. (2006). Structure band texture properties of fried potato products. *Food Reviews int.*, 22: 173-201.
16. Bennett, R.M. (2001). Managing potato crisp processing. In; Rossell, J. B. (Ed.) *Frying: Improving Quality*. CRC Press: Florida. pp. 215-235.
17. Kumar, D. and Ezekiel, R. (2004). Distribution of Dry Matter and Sugars with in a tuber of potato cultivars grown under short day conditions. *PotatoJ*.31 (3-4):130.
18. Kita, A. (2002). The influence of potato chemical composition on crisp texture. *Food Chem.*, 76: 173-179.
19. Badoni, A. and Chauhan, J.S. (2009). Microtuber: A Source of Germplasm Conservation. *Report and Option*, 1, 69-71.
20. Lemaga, B. (2010). The Potato Value Chain in Sub-Saharan Africa with Case Study on Eastern Africa. In: Cromme, N., Prakash, A.B., Lutaladio, N.B. and Ezeta, F., Eds., *Strengthening Potato Value Chains: Technical and Policy Options for Developing Countries*, The Food and Agriculture Organization of the United Nations and the Common Fund for Commodities, Rome, 43-54.
21. Prakash, A.B. (2010). Introduction: The Role of Potato in Developing Country Food Systems. In: Cromme, N., Prakash, A.B., Lutaladio, N.B. and Ezeta, F., Eds., *Strengthening Potato Value Chains: Technical and Policy Options for Developing Countries*, The Food and

Agriculture Organization of the United Nations and the Common Fund for Commodities, Rome, 14-24.

22. Gildemacher, P., Kaguongo, W., Ortiz, O., Tesfaye, A., Woldegiorgis, G., Wagoire, W., Kakuhenzire, R., Kinyae, P., Nyongesa, M. and Struik, P. (2009). Improving potato production in Kenya, Uganda and Ethiopia: A System diagnosis. *Potato Research* 52: 173-205.

23. Abebe, T., Lemaga, B., Mwakasendo, J.A., Nzohabonayoz, Z., Mutware, J., Wanda, K.Y., Kinyae, P. M., Ortiz, O., Crissman, C. and Thiele, G. (2010). Markets for fresh and frozen potato chips in the ASARECA region and the potential for regional trade: Ethiopia, Tanzania, Rwanda, Kenya, Burundi and Uganda. Working Paper. International Potato Centre (CIP). Lima, Peru.

24. Haverkort, A.J., Koesveld, M.J., van Schepers, H.T.A.M., Wijnands, J.H.M., Wustman, R. and Zhang, X.Y. (2012). Potato Prospects for Ethiopia: on the Road to Value Addition. Lelystad: PPO-AGV, (PPO publication 528).

25. Subramanian, N.K., White, P.J., Broadley, M.R. and Ramsay, G. (2011). The three-dimensional distribution of minerals in potato tubers. *Annals of Botany* 107: 681–691.

26. White, P.J., Bradshaw, J.E., Finlay, M., Dale B. and Ramsay, G. (2009). Relationships between yield and mineral concentrations in potato tubers. *Horticultural Science* 44: 6-11.

27. Westermann, D.T. (2005). Nutritional requirements of potatoes. *American Journal of Potato Research* 82: 301-307.

28. Kabira J, Berga L. (2003). Potato processing quality evaluation procedure for research and food industry applications in east and central Africa. *Kenya Agric. Res. Insti. Nairobi*.

29. Krokida MK, Oreopoulou V, Maroulis ZB, Leeratanarak N, Devahastin S, Chiewchan N (2006). Drying kinetics and quality of potato chips undergoing different drying techniques. *J. Food Eng.*, 77: 635-643.

30. Grob K, Biedermann M, Biedermann B, Noti A, Imhof D, Amrein T, Pfefferle A, Bazzocco D. (2003). French fries with less than 100ug/kg acrylamide. A collaboration between cooks and analyst. *Eur. Food Res. Technol.*, 217: 185-194.

31. Tarn, T. R., Tai, G. C. C. and Liu, Q. (2006). Quality improvement. In: J. Gopal, S. M. Paul Khurana (Eds.), *Handbook of Potato Production, Improvement, and Postharvest Management* (pp. 147–178). Food Products Press, New York, London, Oxford.

32. Tai, G. C. C. and Coleman, W. K. (1999). Genotype x environmental interaction of potato chip colour. *Canadian Journal of Plant Science*, 79: 433–438.

33. Kimondo, N.E. (2007). Sensory quality of deep fat fried potato chips manufactured from potatoes with different physico-chemical characteristics. MSc thesis in food science submitted to university of Pretoria, South Africa.

34. Nourian, F., Ramaswamy, H. S. and Kushalappa, A. C. (2003). Kinetic changes in cooking quality of potatoes stored at different temperatures. *Journal of Food Engineering*, 60: 257–266.

35. Brierley, E.R., Bonner, L. P. and Cobb, H. A. (1996). Factors influencing the free amino acid content of potato (*Solanum tuberosum* L.) tubers during prolonged storage. *J Sci. Food and Agri.*, 70: 515-525.

36. Ames, J.M. (1990). Control of Maillard reaction in food system. *Trends in Food Sci. and Technol.*, 12: 150-154.

37. Rodriguez-Saona L, Wrolstad R (1997). Influence of potato composition on chip color quality. *Am. Potato J.*, 74: 87-106.

38. Kabira, J.N. and Lemaga, B. (2006). Potato Processing: Quality Evaluation Procedures for Research and Food Industries Applications in East and Central Africa. *Kenya Agricultural Research Institute, Nairobi, Kenya*.

39. Thygesen, L. G., Thybo, A. K. and Engelsen, S. B. (2001). Prediction of sensory texture quality of boiled potatoes from low-field <sup>1</sup>H NMR of raw potatoes. The role of chemical constituents. *Lebensmittel-Wissenschaft und-Technologie*, 34: 469–477.

40. Tarn, T. R., Tai, G. C. C., De Jong, H., Murphy, A. M. and Seabrook, J. E. A. (1992). Breeding potatoes for long-day, temperate climates. *Plant Breeding Reviews*, 9: 217–332.

41. Fennema O. (1996). Food Chemistry.3rd edition. New York: Marcel Dekker Fitzpatrick JJ,PorterWL, Houghland VC (1964).Continued studies of the relationship of specific gravity to total solids of potato. Am. Potato J., 46: 120-127.
42. Marquez, G, Anon MC. (1986). Influence of reducing sugars and amino acids in the color development of fried potatoes. J. Food Sci., 51: 157-160.
43. Segnini, S. and Dejemek, P. (1999). Relationship between instrumental and sensory analysis of texture and color of potato chips .J. *Texture Studies*.30:677-690.
44. Burton, W.G., Van Es, A. and Hartmans, J. K. (1992).The Physics and Physiology of Storage.In The Potato Crop 3rd ed;Champan and Hall,London.pp.608-727.
- 45.Stevenson FJ, Akeley RV, Cunningham CE. (1964). The potato- its genetic and Environmental variability. Am. Potato J., 41: 46-53.
46. Moyano PC, Troncoso E, Pedreschi F. (2007). Modeling texture Kinetics during Thermal processing of potato products. J. Food Sci., 72.
47. Kayacier A, Singh R. (2003). Textural properties of baked tortilla chips. Lebensmittel -Wissenschaft und Technol., 36: 463-466.
48. Lisinska G, Eszczynski WL.(1989). Potato Science and Technology. Elsevier Applied Science,New York, NY.
49. Asmamaw Y, Tekalign T., Tilahun S. (2010). Specific gravity, dry matter 332 Afr. J. Food Sci.concentration, pH, and crisp-making potential of Ethiopian potato (*Solanum tuberosum* L. cultivars as influenced by growing environment and length of storage under ambients. Potato Res., 53: 95-109
50. Pimpaporn P, Devahastin S, Chiewchan N. (2007). Effect of combined pretreatments on drying kinetics and quality of potato chips undergoing low-pressure superheated steam drying. J. Food Eng., 81: 318-329.
51. Bartolome LG, Hoff JE. (1972). Firming of potatoes: biochemical effects of pre heating. J. Agric. Food Chem., 20: 266-270
52. Andersson A, Gekas V, Lind L, Oliveira F, Oste R. (1994). Effect of preheating on potato texture Rev. Food Sci. Technol., 34: 229-251.
53. Dinehart M.E., Hayes, J. E., Bartoshuk, L. M., Lanier, S. L. and Duffy, V. B. (2006).Bitter taste markers explain variability in vegetable sweetness, bitterness and intake.Physiol Behav 87:304-13.
54. Duckham S.C, Dodson A.T., Bakker, J. and Ames, J. M. (2001).Volatil flavor components of baked potato flesh: A comparison of eleven potato cvltivars. Food/Nahrung 45:317-23.
- 55.Scanlon, M.( 2003). Comercial potato production: Botany of the potato.*Website:http://www.gov.mb.ca/agricultural/crops/potatoes/bda04s02*.
56. Stier, R. F. (2000). Chemistry of frying and optimization of deep fat fried foods flavor-An introductory review.Eu.J of Lipid Sci and Technology, 102.507-514.
- 57.Gillat, P. (2001). Flavour and aroma development in frying and fried food. In; Rossell, J. B. ( Ed).Frying; Improving quality. CRC Press; florida, 266-327.
58. Dale, M.F.B. and Mackay, G. R.(1994). Inheritance of table and processing quality. In bradshaw, J.E. and Mackay,g.R(Ed) Potato genetics. CABI International.285-315.
- 59.Ereifej, K.I., Shibli, R. A., Ajlouni,M. M. and Hussein,A.(1997). Chemical composition variations of tissues and processing characteristics in ten potato cultivars grown in Jordan. Americ an Journal of potato Research,74 23-30.
60. Maga, J.A. (1994).Potato favour. Food reviews International,101,1-48.
- 61.Moreira RG, Castell-Perez ME, Barrufet MA. (1999). Deep-fat Frying: Fundamentals and Applications. Gatihersburg, MD: Aspen Publishers
62. Taiwo KA, Angersbach A, Ade-Omowaye BIO, Knorr D. (2001). Effect of pre-treatments on the diffusion kinetics and some quality parameters of osmotically dehydrated apple slices. J. Agric. Food Chem., 49: 2804-2811.
63. Moreno-Perez, LF, Gasson-Lara Jh, Ortega-Riuas E. (1996). Effect of low temperature-long time blanching on quality of dried sweet potato. Drying Technol., 14: 1839-1857.