

**Direct and residual effect of phosphorus fertilizer with AM fungi in  
maize- green gram cropping sequence on nutrients content and uptake**

**ASMATULLAH DURANI<sup>1\*</sup>, SONAL TRIPATHI<sup>2</sup>, AMINULLAH  
YOUSAFZAI<sup>3</sup>, HASHMATULLA DURRANI<sup>4</sup> AND S. M BAMBHANEYYA<sup>5</sup>**

<sup>1,2,4,5</sup> Department of Soil Science and Agricultural Chemistry, N. M. College of Agriculture (NMCA),  
Navsari Agricultural University (NAU), Navsari, Gujrat, India.

<sup>3</sup>Department of Horticulture, Faculty of Agriculture, Nangarhar University, Nangarhar, Afghanistan.

**Authors' contributions**

*This work was carried out in collaboration between all authors. Author AD designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript.*

*Authors ST and AY managed the analyses of the study and manuscript writing. Authors HD, and SMB managed the literature references. All authors read and approved the final manuscript.*

---

**ABSTRACT**

The field experiment was conducted at the College Farm, Navsari Agricultural University, Navsari in the year 2015-16 and 2016-17 to study the direct and residual effect of phosphorus fertilizer with AM fungi in maize-green gram cropping sequence on nutrients content and uptake during 2015-16 and 2016-17. Application of phosphorus fertilizer SSP and RP (composted) alone or combined with AM fungi significantly increases the NPK content in maize grain and in straw during both years of the study and in the pooled analysis. The treatment 75%P as RP+AM (290.83, 251.36 and 266.19 %) increased total nitrogen uptake and 333.11, 345.44 and 340.35% total phosphorus uptake by maize (grain + straw) over control T<sub>2</sub> during both the years as well as in pooled analysis, respectively. The total potassium uptake by maize (grain + straw) in treatment 75% P as RP+AM increased 231.1 and 124.3 % over control T<sub>2</sub> during the first year of the study and in pooled analysis, respectively and in the treatment 75% P as SSP+AM increased (92.67 %) of total potassium uptake by maize (grain + straw) over control T<sub>2</sub> during the year 2016-17. Application of treatment 75% P as RP+AM applied to preceding rabi maize increased 425.14%, 320.03 and 358.20 % of total nitrogen uptake 561.54, 377.78 and 450.24% of total phosphorus uptake and 290.21, 147.00 and 191.62% total potassium uptake by green gram (grain + stover) during 2015-16, 2016-17 and in pooled analysis respectively over control T<sub>2</sub>.

**Keywords:** Effect of phosphorus fertilizer, nutrients content and uptake, cropping sequence

\* E-mail address: xyz@abc.com.

## 28 1. INTRODUCTION

29

30 In India, most of the soils are either deficient or marginal in P status.  
31 Adequate P fertilization is thus essential for economic and sustained crop  
32 production. Phosphorus deficient soils require a high dose of phosphatic  
33 fertilizers which are imported and expensive. Also, the phosphorous  
34 fertilizers immediate conversion of water-soluble P due to P fixation results  
35 in low fertilizer use efficiency. Among the different inorganic P sources,  
36 single super phosphate (SSP) is the most widely used phosphatic fertilizers  
37 which supply P in water-soluble form in the immediate vicinity of roots. Its  
38 importance as the most efficient P fertilizer source is well established but it  
39 is very expensive and needs to be imported. It also suffers from the  
40 problem of fixation in the long run. However, India has vast resources of  
41 indigenous rock phosphate (RP), unfortunately, most of the RPs of Indian  
42 origin have the limitation of low  $P_2O_5$  content and low reactivity and perform  
43 poorly when applied directly to the neutral soil and are not suitable for the  
44 manufacture of phosphatic fertilizer. With the discovery of several deposits  
45 of RP in the country, interest in the use of this indigenous material as  
46 alternative phosphatic fertilizers has increased greatly. Although RP can  
47 effectively replace water-soluble phosphates in acid soils, but its efficiency  
48 in neutral, alkaline and calcareous soils is extremely low. To make it  
49 effective in such soils it is being converted into water-soluble form by  
50 mixing with SSP or by partial acidulation with mineral acids, for which  
51 sulphur is being imported.

52 Pulses are integral part of Indian dietary system because of its  
53 richness in proteins and other important nutrients such as Ca, Fe, and  
54 vitamins viz., carotene, thiamine, riboflavin and niacine. Indian population  
55 is predominantly vegetarian and protein requirement for the growth and  
56 development of the human being is mostly met with pulses. Green gram is  
57 an important pulse crop of Indian as it is grown an area of 3.44 million  
58 hectares with total production of 1.4 million tonnes and productivity of 407  
59 kg/ha. In India, major green gram producing states are Odissa, Madhya  
60 Pradesh, Rajasthan, Maharashtra, Gujarat and Bihar. In Gujarat, it is  
61 cultivated in about 2.3 lakh hectares with an annual production of 1.21  
62 lakh tonnes and average productivity of 526 kg/ha [3].

63 [23], reported that sorghum plant inoculated with VAM recorded  
64 higher amount of P, K, Mg, Mn, S, Ca, Fe, Cu and Zn than non-mycorrhizal

65 plants. A part from the fact that phosphorus from rock phosphate is  
 66 solubilized during composting and transformed into available forms,  
 67 enrichment of the compost with rock phosphate also accelerates its rate of  
 68 decomposition [28]. The total P, water soluble P, citrate soluble P, total N  
 69 and NO<sub>3</sub>-N content was also found to increase in the mature phospho-  
 70 compost, [13]. [14], Found that nitrogen and phosphorus uptake by seed and  
 71 stover as well as the total N and P<sub>2</sub>O<sub>5</sub> by maize was found significantly  
 72 superior under the application of 40 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> over 20 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>.

## 73 2. MATERIAL AND METHODS

74 The field experiment was conducted at the College Farm, Navsari  
 75 Agricultural University, Navsari (Gujarat), during 2015-16 and 2016-17.  
 76 Navsari is located 20° 57' N latitude and 72° 54' E longitudes, in the  
 77 tropical region; having an altitude of 10 meters above the mean sea level.  
 78 The campus is located at 3 km away towards west of Navsari and 13 km  
 79 away from the Arabian Sea towards east. The climate of this region is  
 80 characterized by fairly hot summer, moderately cold winter and warm humid  
 81 monsoon with heavy rainfall.

82 The soil of south Gujarat is locally known as “Deep Black Soil”. The  
 83 soil of Navsari campus is classified under the order *Inceptisols* comprising  
 84 of fine *montmorillonitic*, *isohyperthermic*, family of *Vertic Ustrochrepts* and  
 85 soil series Jalalpur by the soil survey officer, Navsari. The important  
 86 physicochemical properties of experimental soil at the initiation were  
 87 presented in Table 1. *Rabi* maize as main plot treatments replicated three  
 88 times in randomized block design with 14 treatment. During summer season  
 89 each main plot treatment was split into two sub plot treatments with two  
 90 level of recommended dose of fertilizers viz., F<sub>1</sub> (75% RDF) and F<sub>2</sub> (100%  
 91 RDF) to green gram resulting in 28 treatment combinations replicated three  
 92 times in split plot design.

93 **Table1.** Important physicochemical properties of experimental soil (0-30  
 94 cm) at the initiation of the experiment.

Sr. No.	Particulars	Values		Methods employed
		2015-16	2016-17	
I	<b>Physical properties</b>			
	Mechanical separates %			
	Fine sand	20.1	20.32	
	Coarse sand	1.76	1.66	
1	Silt	15.95	15.89	International pipette method, [22].
	Clay	61.70	62.13	
	Textural class	Clay	Clay	
2	Bulk density	1.389	1.375	Black, [4].

\* E-mail address: xyz@abc.com.

	(g/cc)			
<b>II</b>	<b>Chemical properties</b>			
1	pH	7.80	7.94	1:2.5 water suspension, [6].
2	EC	0.16	0.43	at 250C (1:2.5) dS/m, [6].
3	Organic carbon %	0.440	0.45	Rapid titration method , [29].
4	Available N kg/ha	206.5	209.3	Alkaline permanganate method, [26].
5	Available P <sub>2</sub> O <sub>5</sub> kg/ha	31.20	38.30	0.5 M Na HCO <sub>3</sub> , pH= 8.5, [18].
6	Available K <sub>2</sub> O kg/ha	323.2	274.9	Neutral ammonium acetate,[15].
<b>III</b>	<b>DTPA extractible micronutrients (mg/kg)</b>			
7	Fe	18.70	19.60	DTPA method, [12].
8	Mn	16.80	19.10	
9	Zn	0.489	0.521	
10	Cu	0.491	0.632	

95

96 **Table 2.** Detail of the treatments evaluated in rabi maize and summer green  
97 gram

Treatment No.	Treatments details	Treatment code
<b>Main plot treatment</b>		
T <sub>1</sub>	Rabi Fallow (No maize crop, absolute control)	<i>Rabi fallow</i>
T <sub>2</sub>	Control (without phosphorus and AM)	control
T <sub>3</sub>	50 percent of phosphorus from rock phosphate (composted)	50% P as RP
T <sub>4</sub>	50 percent of phosphorus from rock phosphate (composted) + Arbuscular mycorrhizae	50% P as RP +AM
T <sub>5</sub>	50 percent of phosphorus from single supper phosphate (composted)	50% P as SSP
T <sub>6</sub>	50 percent of phosphorus from single supper phosphate (composted) + Arbuscular mycorrhizae	50% P as SSP+AM
T <sub>7</sub>	75 percent of phosphorus from rock phosphate (composted)	75% P as RP
T <sub>8</sub>	75 percent of phosphorus from rock phosphate (composted)+ Arbuscular mycorrhizae	75% P as RP+AM
T <sub>9</sub>	75 percent of phosphorus from single supper phosphate (composted)	75% P as SSP
T <sub>10</sub>	75 percent of phosphorus from single supper phosphate (composted)+ Arbuscular mycorrhizae	75% P as SSP+AM
T <sub>11</sub>	100 percent of phosphorus from rock phosphate (composted)	100% P as RP
T <sub>12</sub>	100 percent of phosphorus from rock phosphate (composted)+ Arbuscular mycorrhizae	100% P as RP+AM
T <sub>13</sub>	100 percent of phosphorus from single supper phosphate (composted)	100% P as SSP
T <sub>14</sub>	10 percent of phosphorus from single supper phosphate (composted)+ Arbuscular mycorrhizae	100 % P as SSP+AM
<b>Sub plot treatments</b>		
F <sub>1</sub>	75 percent of recommended dose of fertilizer	75% RDF
F <sub>2</sub>	100 percent of recommended dose of fertilizer	100% RDF

\* E-mail address: xyz@abc.com.

**Note:** Applied fertilizer for *rabi* maize crop 120:60:00 NPK kg/ha with or without of Arbuscular mycorrhizae 250g/ha which have 70 percent raw materials and 30 % VAM 3000 infected propagates/g and two level of recommended dose of fertilizer for summer green gram though 20:40:00 NPK kg/ha.

**Table-3:** Initial properties of the rock phosphate enriched compost and bio-compost

Parameters	Rock phosphate enriched compost		Bio-compost	
Properties	2015-16	2016-17	2015-16	2016-17
pH	7.3	7.1	6.30	6.10
EC dS/m	2.11	2.09	0.491	0.501
Organic carbon %	26.67	29.05	32.66	33.55
Total P %	8.00	8.00	0.34	0.32
Available N %	0.49	0.45	2.42	2.12
Available K %	0.88	0.90	1.45	1.65
Fe mg/kg	143.9	142.4	0.21	0.32
Mn mg/kg	86.00	83.99	98.6	87.5
Zn mg/kg	44.55	33.89	24.4	26.3
Cu mg/kg	18.33	11.33	1.34	1.56

The nitrogen was applied through urea (46% N) whereas phosphorus was applied through single superphosphate (16% P<sub>2</sub>O<sub>5</sub>) and rock phosphate was applied as basal on the base of 8% total phosphorus content for increasing the effectiveness of RP on alkaline soil the it was composted with organic matter (Cowden) in 1:15 ratio along with PSB (*Bacillus megatherium*) for 45 day (Table 3). A common dose of organic manures (bio-compost at @ 15 t/ha) applied to all treatments before sowing of *rabi* maize and evenly spread and mixed in that particular bed. The properties of the bio-compost and rock phosphate enriched compost mentioned in the Table.3. Periodical plant samples were dried at about 60° C and their dry weight was recorded the total PK content in the extract (10HNO<sub>3</sub>: 4HClO<sub>4</sub>) extraction was determined by using Micro plasma-atomic emission spectroscopy (MP-AES) [11]. In case of total N, plant sample was analyzed by micro-kjeldhal assembly according to procedure outlined by [7]. The data on various variables were analyzed by using statistical procedures and pooled analysis of the preceding *rabi* maize analyzed for two years was worked out as per the **standard** method [19].

### 3. RESULTS AND DISCUSSION

#### 3.1 NPK content (%) and uptake by maize

From appraisal of data presented in Table-4, it could be seen that the total nitrogen content in maize grain was found significant due to the

different phosphorus fertilizer treatments. The significantly higher value of nitrogen content in maize grain was recorded 1.095 % under T<sub>11</sub> treatment, which was at par with T<sub>4</sub>, T<sub>5</sub>, T<sub>8</sub>, T<sub>12</sub>, T<sub>13</sub> and T<sub>14</sub> treatments during the 2015-16. In the case of second year 2016-17 and in pooled analysis significantly higher nitrogen (1.105 and 1.070 %) was recorded under treatment T<sub>8</sub> which was at par with all phosphorus fertilizer treatments barring T<sub>2</sub> and T<sub>7</sub> in year 2016-17 and T<sub>2</sub>, T<sub>6</sub>, T<sub>7</sub>, T<sub>9</sub> and T<sub>10</sub> during pooled analysis. The results in Table-5, showed that nitrogen content in maize straw was influenced significantly by different treatments applied to *rabi* maize crop. The nitrogen content in maize straw was significantly higher (0.564, 0.545 and 0.554 %) with application of 75 %P as RP (composted)+AM (T<sub>8</sub>) during both the years as well as in pooled analysis respectively, which was at par with all treatments barring T<sub>2</sub> and T<sub>7</sub> treatments during 2015-16. In the year 2016-17, treatment T<sub>8</sub> was statistically at par among the other phosphorus applied fertilizer treatments except control T<sub>2</sub>, T<sub>7</sub>, T<sub>9</sub>, and T<sub>10</sub> treatments. Similarly in pooled analysis T<sub>8</sub> treatment was at par with all treatments under the study barring T<sub>2</sub>, T<sub>3</sub>, T<sub>7</sub>, T<sub>9</sub> and T<sub>10</sub> treatments.

The data presented in Table-4, the significantly higher total phosphorus registered in maize grain was due to application of 100%P as SSP+AM (T<sub>14</sub>, 0.381%) treatment which was statistically at par with 50% P as RP +AM, 50% P as SSP+AM, 75% P as RP, 75% P as RP+AM, 75% P as SSP, 75% P as SSP+AM, 100% P as RP, 100% P as RP+AM and 100% P as SSP treatments during 2015-16. In the case of second year, significantly higher total phosphorus content in maize grain was recorded in treatment 100% P as RP+AM (T<sub>12</sub>, 0.330 %) which was statistically at par with all phosphorus fertilizer applied treatments barring control and 50% P as SSP+AM. While in pooled analysis, significantly higher phosphorus content in maize grain was found to be 0.346 % in treatment 100 % P as SSP+AM which was statistically at par with all treatments except for control, 50% P as RP, 50% P as RP +AM, 50% P as SSP, 50% P as SSP+AM, 75% P as RP and 75% P as SSP. Significantly lower total phosphorus content in plant was observed 0.118, 0.104 and 0.111 % in control plots and maximum under 100 % P as SSP+AM (T<sub>14</sub>, 0.213, 0.204 and 0.209 %) in the years 2015-16, 2016-17 and in pooled analysis, respectively. Treatment 100 % P as SSP+AM (T<sub>14</sub>) was at par with all phosphorus fertilizer treatments except

treatments control (T<sub>2</sub>, 0.118%) and 50% P as SSP (T<sub>5</sub>, 0.180%) in first year, treatment control (T<sub>2</sub>, 0.104%) in second year and treatments control (T<sub>2</sub>, 0.111%), 50% P as SSP (T<sub>5</sub>, 0.183%), 75% P as SSP (T<sub>9</sub>, 0.190%) and 100% P as SSP (T<sub>13</sub>, 0.191%) in pooled analysis (Table-5).

The potassium content in maize grain 2015-16 and pooled analysis was found to be non significant. In the case of second year, significantly higher potassium (K) content in maize grain was obtained under treatment 75% P as SSP+AM (T<sub>10</sub>, 0.593 %) which was statistically at par with treatment 100% P as SSP (T<sub>13</sub>, 0.507 %) (Table-4). This due to fact that application of phosphorus fertilizer maintained higher phosphorus availability to maize which promotes the root growth and other part of the plant and increased N and P content in maize grain and straw. The beneficial effect of phosphorus fertilizer SSP, RP alone or combination with AM fungi increased NPK content in maize grain and straw over no phosphorus fertilizer these result are in accordance with the finding of [27] and [26], [17], [2] and [16].

**Table 4: Nutrient (NPK) content in maize grain**

Treatment	Nitrogen (%)			Phosphorus (%)			Potassium (%)		
	2015-16	2016-17	Pooled	2015-16	2016-17	Pooled	2015-16	2016-17	Pooled
T2	0.667	0.624	0.646	0.168	0.104	0.136	0.229	0.413	0.321
T3	0.923	0.969	0.946	0.247	0.245	0.246	0.276	0.453	0.365
T4	1.010	0.939	0.975	0.300	0.263	0.281	0.194	0.487	0.340
T5	0.997	1.014	1.005	0.232	0.236	0.234	0.192	0.440	0.316
T6	0.903	0.960	0.932	0.326	0.207	0.267	0.216	0.420	0.318
T7	0.900	0.852	0.876	0.281	0.235	0.258	0.236	0.380	0.308
T8	1.034	1.105	1.070	0.333	0.284	0.309	0.239	0.460	0.350
T9	0.877	0.962	0.920	0.289	0.231	0.260	0.237	0.447	0.342
T10	0.868	0.908	0.888	0.306	0.279	0.292	0.196	0.593	0.395
T11	1.095	1.004	1.050	0.287	0.287	0.287	0.223	0.473	0.348
T12	1.008	1.011	1.010	0.327	0.330	0.329	0.276	0.413	0.345
T13	0.934	0.999	0.967	0.320	0.245	0.283	0.180	0.507	0.343
T14	0.936	1.094	1.015	0.381	0.312	0.346	0.242	0.430	0.334
S.Em.±	0.055	0.069	0.044	0.035	0.037	0.026	0.024	0.033	0.037
C.D. at 5 %	0.163	0.203	0.127	0.102	0.109	0.073	NS	0.098	NS
YXT S.Em.±	—	—	0.017	—	—	0.036	—	—	0.005
C.D. at 5 %	—	—	NS	—	—	NS	—	—	0.014
C.V. %	10.40	12.60	11.60	9.50	6.20	7.70	8.70	9.90	9.00
General mean	0.935	0.957	0.946	0.292	0.251	0.271	0.226	0.455	0.340

T<sub>1</sub>=Rabi Fallow (No maize crop, absolute control).

**Table 5: Nutrient (NPK) content in maize straw**


Treatment	Nitrogen (%)			Phosphorus (%)			Potassium (%)		
	2015-16	2016-17	Pooled	2015-16	2016-17	Pooled	2015-16	2016-17	Pooled
T <sub>2</sub>	0.324	0.296	0.310	0.118	0.104	0.111	0.614	1.140	0.877
T <sub>3</sub>	0.507	0.440	0.474	0.190	0.193	0.192	0.701	1.380	1.040
T <sub>4</sub>	0.504	0.497	0.501	0.197	0.192	0.194	0.805	1.227	1.016
T <sub>5</sub>	0.541	0.482	0.512	0.180	0.186	0.183	0.900	1.087	0.994
T <sub>6</sub>	0.515	0.463	0.489	0.198	0.196	0.197	0.821	1.400	1.110
T <sub>7</sub>	0.460	0.416	0.438	0.195	0.191	0.193	0.621	1.167	0.894
T <sub>8</sub>	0.564	0.545	0.554	0.203	0.197	0.200	1.035	1.113	1.074
T <sub>9</sub>	0.515	0.404	0.460	0.190	0.189	0.190	0.861	1.133	0.997
T <sub>10</sub>	0.478	0.409	0.443	0.195	0.197	0.196	0.757	1.087	0.922
T <sub>11</sub>	0.536	0.521	0.529	0.188	0.198	0.193	0.832	1.147	0.989
T <sub>12</sub>	0.540	0.470	0.505	0.202	0.198	0.200	0.794	1.147	0.970
T <sub>13</sub>	0.522	0.478	0.500	0.190	0.192	0.191	0.774	1.233	1.004
T <sub>14</sub>	0.559	0.532	0.546	0.213	0.204	0.209	0.788	1.313	1.051
S.Em.±	0.035	0.037	0.025	0.0102	0.010	0.006	0.101	0.087	0.067
C.D. at 5 %	0.103	0.108	0.072	0.030	0.019	0.017	NS	NS	NS
YXT S.Em.±	—	—	0.010	—	—	0.007	—	—	0.005
C.D. at 5 %	—	—	NS	—	—	NS	—	—	NS
C.V. %	12.00	9.90	9.00	8.55	9.66	8.92	6.10	12.60	6.40
General mean	0.505	0.451	0.481	0.189	0.189	0.188	0.792	1.190	0.995

187

188 The highest total nitrogen uptake (Fig.1 A) by maize was observed  
 189 under treatment T<sub>8</sub> during the first year of the study and in pooled analysis.  
 190 While in the case of second year of the study the total nitrogen uptake by  
 191 maize was significantly higher under treatment T<sub>8</sub> which was at par with  
 192 treatments T<sub>5</sub> and T<sub>14</sub>. The total nitrogen uptake in the treatment T<sub>8</sub>  
 193 increased (290.83, 251.36 and 266.19 %) over control (T<sub>2</sub>) during both the  
 194 years as well as in pooled analysis, respectively.

195 The application of phosphorus fertilizer alone or along with AM fungi  
 196 increased the total phosphorus uptake in maize crop over control T<sub>2</sub>.  
 197 Significantly higher total phosphorus uptake by maize was observed in  
 198 treatment 75% P as RP+AM (T<sub>8</sub>, 19.36, 25.39 and 22.37 kg/ha), which was  
 199 333.11, 345.44 and 340.35% more than control T<sub>2</sub> during both the year as  
 200 well as in pooled analysis respectively (Fig.1 B). Total potassium uptake by  
 201 maize in treatment 75% P as RP+AM (T<sub>8</sub>) recorded (231.1 and 124.3 %)  
 202 increased over control T<sub>2</sub> during first year of the study and in pooled  
 203 analysis, respectively and in the treatment 75% P as SSP+AM (T<sub>10</sub>, 92.67  
 204 %) increased over control T<sub>2</sub> during 2016-17 year (Fig.1 C). This could be  
 205 attributed to the fact that added phosphorus increased N and P  
 206 concentration in grain and stover by providing balanced nutritional  
 207 environment inside the plant and higher photosynthetic efficiency, which

208 favoured growth and crop yield. Since, the uptake of nutrients is a function  
 209 of dry matter (grain and stover) and nutrient content, the increased grain  
 210 and stover yield together with higher NPK and content resulted in greater  
 211 uptake of these elements. Application of P without or with *Mychorriza*  
 212 inoculation significantly increased the uptake of N, P and K by maize over  
 213 control. A smlir reported by [25], [7] and [2].  
 214



215  
 216  
 217  
 218  
 219  
 220  
 221  
 222  
 223  
 224 **Fig-1: Total NPK uptake by *rabi* maize as influenced by different treatment**  
 225 (A) **Total N uptake, (B) Total P uptake (C) Total K uptake**

226 **3.2 NPK content (%) and uptake by green gram**

227 **3.2.1 Residual effect**

228 The data regarding to NPK content in green gram seeds, stover and  
 229 pod cover presented in Table-6, Table-7, Table-8. Significantly higher  
 230 nitrogen content in green gram seeds was recorded with the application of  
 231 treatment 75%P as RP+AM (T<sub>8</sub>, 2.152, 2.546 and 2.349%) during year 2015-  
 232 16, 2016-17 and in pooled analysis respectively. In the case of green gram  
 233 stover nitrogen content was observed significantly higher under treatment  
 234 75% P as SSP (T<sub>9</sub>, 1.056 and 1.208 %) during first year of the study and in  
 235 pooled analysis, respectively and in the year 2016-17 nitrogen content was  
 236 recorded significantly higher in treatment 50% P as SSP T<sub>5</sub>, 1.418 %

\* E-mail address: xyz@abc.com.

(Table-6).The results might be due to the application of residual effect phosphorus fertilizer to previous *rabi* maize and respective rate of RDF to summer green gram which was higher removal of N and P might be due to better development of root growth which was further increased significantly N content in green gram seeds and stover. Similar results were also reported by [13] in green gram.

**Table.6: Nitrogen content seeds, stover and pod cove of green gram**

Treatment	Nitrogen in seeds (%)			Nitrogen in stover (%)			Nitrogen in pod cover (%)		
	2015-16	2016-17	Pooled	2015-16	2016-17	Pooled	2015-16	2016-17	Pooled
T <sub>1</sub>	1.500	1.447	1.474	0.609	1.422	1.016	0.799	0.915	0.857
T <sub>2</sub>	1.113	1.390	1.252	0.600	0.659	0.630	0.619	0.760	0.689
T <sub>3</sub>	1.780	2.080	1.930	0.958	1.284	1.121	0.770	0.854	0.812
T <sub>4</sub>	1.860	2.292	2.076	0.872	1.196	1.034	0.795	0.960	0.878
T <sub>5</sub>	1.914	2.403	2.159	0.952	1.418	1.185	0.764	0.969	0.866
T <sub>6</sub>	2.021	2.487	2.254	0.891	1.256	1.074	0.874	0.955	0.915
T <sub>7</sub>	1.924	2.411	2.167	0.925	1.268	1.097	0.777	0.852	0.814
T <sub>8</sub>	2.152	2.546	2.349	1.054	1.245	1.150	0.813	0.911	0.862
T <sub>9</sub>	2.049	2.283	2.166	1.056	1.359	1.208	0.800	0.856	0.828
T <sub>10</sub>	1.957	1.895	1.926	0.890	1.336	1.113	0.815	0.949	0.882
T <sub>11</sub>	1.826	2.384	2.105	0.973	1.116	1.045	0.800	0.879	0.840
T <sub>12</sub>	1.971	2.487	2.229	0.922	1.166	1.044	0.783	0.901	0.842
T <sub>13</sub>	1.668	2.303	1.985	0.780	1.196	0.988	0.727	0.894	0.810
T <sub>14</sub>	1.537	2.402	1.970	0.761	1.090	0.926	0.689	0.820	0.755
S.Em.±	0.155	0.201	0.235	0.070	0.103	0.042	0.065	0.07	0.030
C.D. at 5 %	0.451	0.584	0.360	0.230	0.298	0.184	NS	NS	NS
C.V. %	6.08	4.77	5.70	5.40	6.3	4.68	2.43	5.79	5.56
F <sub>1</sub>	1.807	2.219	2.013	0.858	1.191	1.025	0.769	0.907	0.838
F <sub>2</sub>	1.804	2.183	1.993	0.891	1.239	1.065	0.777	0.875	0.826
S.Em.±	0.049	0.030	0.028	0.02	0.026	0.02	0.022	0.015	0.01
C.D. at 5 %	NS	NS	NS	NS	NS	NS	NS	NS	NS
T×F S.Em.±	0.184	0.113	0.15	0.07	0.110	0.07	0.082	0.057	0.05
C.D. at 5 %	NS	NS	NS	NS	NS	NS	NS	NS	NS
C.V. %	4.27	2.99	3.9	3.58	4.35	2.37	1.44	3.1	4.74
General mean	1.803	2.182	2.002	0.891	1.230	1.044	0.777	0.874	0.832

244

245 Significantly higher phosphorus content in green gram seeds was  
 246 recorded with application of 100% P as RP (T<sub>11</sub>) in the first year and  
 247 treatments 75% P as SSP+AM T<sub>(10)</sub> and 75% P as RP+AM (T<sub>8</sub>) during 2016-  
 248 17 and pooled analysis respectively. Similarly significantly higher  
 249 phosphorus content in green gram stover was found (0.423 and 0.395%) in  
 250 the treatment 75% P as RP+AM (T<sub>8</sub>) during 2015-16 and in pooled analysis  
 251 while second year in treatment 100% P as RP (T<sub>11</sub>, 0.379 %). Significantly  
 252 higher phosphorus content in green gram pod cover was recorded under  
 253 treatment 75% P as RP+AM (T<sub>8</sub>, 0.466 and 0.486 %) during fist and second  
 254 years while in pooled analysis result indicated that the significantly highest

255 value of phosphorus content in pod cover was found under T<sub>8</sub> (0.476 %)   
 256 treatment (Table-7). It might be due to the residual effect of different   
 257 phosphorus fertilizers SSP and RP alone and combined with AM to   
 258 preceding *rabi* maize which were more availability of phosphorus in soil   
 259 which had residual fertility status increased significantly phosphorus   
 260 content in green gram grain, stover and pod cover. The results are in   
 261 agreement with the finding [20] and [21].

262 **Table.7: Phosphorus content in seeds, stover and pod cover of green gram**

Treatment	Phosphorus in seeds (%)			Phosphorus in stover (%)			Phosphorus in pod cover (%)		
	2015-16	2016-17	Pooled	2015-16	2016-17	Pooled	2015-16	2016-17	Pooled
T <sub>1</sub>	0.260	0.322	0.291	0.225	0.198	0.211	0.130	0.118	0.124
T <sub>2</sub>	0.263	0.234	0.248	0.138	0.156	0.147	0.110	0.098	0.104
T <sub>3</sub>	0.350	0.435	0.393	0.312	0.327	0.319	0.328	0.307	0.317
T <sub>4</sub>	0.361	0.459	0.410	0.302	0.353	0.328	0.358	0.337	0.348
T <sub>5</sub>	0.367	0.412	0.389	0.288	0.334	0.311	0.340	0.317	0.328
T <sub>6</sub>	0.341	0.398	0.370	0.272	0.320	0.296	0.337	0.313	0.325
T <sub>7</sub>	0.361	0.429	0.395	0.293	0.377	0.335	0.364	0.345	0.355
T <sub>8</sub>	0.436	0.441	0.439	0.423	0.368	0.395	0.466	0.486	0.476
T <sub>9</sub>	0.371	0.447	0.409	0.298	0.330	0.314	0.388	0.408	0.398
T <sub>10</sub>	0.352	0.493	0.422	0.344	0.366	0.355	0.369	0.359	0.364
T <sub>11</sub>	0.443	0.420	0.432	0.347	0.379	0.363	0.376	0.366	0.371
T <sub>12</sub>	0.339	0.373	0.356	0.302	0.343	0.322	0.354	0.366	0.360
T <sub>13</sub>	0.360	0.419	0.390	0.349	0.373	0.361	0.367	0.379	0.373
T <sub>14</sub>	0.361	0.386	0.374	0.261	0.351	0.306	0.371	0.383	0.377
S.E.m.±	0.034	0.038	0.025	0.026	0.042	0.024	0.034	0.034	0.023
C.D. at 5 %	0.098	0.109	0.071	0.075	0.120	0.069	0.098	0.098	0.067
C.V. %	3.6	4.2	3.6	3.9	3.2	2.3	2.4	3.2	2.9
F <sub>1</sub>	0.356	0.412	0.384	0.304	0.328	0.316	0.327	0.322	0.324
F <sub>2</sub>	0.353	0.397	0.375	0.289	0.326	0.308	0.338	0.333	0.336
S.E.m.±	0.013	0.009	0.034	0.008	0.013	0.010	0.011	0.020	0.033
C.D. at 5 %	NS	NS	NS	NS	NS	NS	NS	NS	NS
T×F S.E.m.±	0.049	0.035	0.037	0.028	0.470	0.026	0.041	0.430	0.011
C.D. at 5 %	NS	NS	NS	NS	NS	NS	NS	NS	NS
C.V. %	2.7	2.8	2.4	2.3	2.3	1.9	1.9	2.1	1.6
General mean	0.355	0.405	0.380	0.297	0.327	0.312	0.333	0.327	0.330

263

264 Significantly higher potassium content in green gram seeds was   
 265 recorded with application 75% P as RP+AM (T<sub>8</sub>, 0. 876, 1.040 and 0.958 %)   
 266 during first, second and in pooled analysis respectively, (Table-8).

267 This might be due the adequate supply of potassium supplemented   
 268 by the beneficial residual effects *viz.*, mineralization and slow release of   
 269 nutrients by rock phosphate and synergistic effect of balance P fertilization   
 270 resulting in higher potassium concentration in seeds. [10], reported that rock   
 271 phosphate with organic manure increased the uptake of major nutrients like   
 272 N, P, K, Ca and Mg.

273

**Table.8: Potassium content in seeds, stover and pod cover of green gram**


Treatment	Potassium in seeds (%)			Potassium in stover (%)			Potassium in pod cover (%)		
	2015-16	2016-17	Pooled	2015-16	2016-17	Pooled	2015-16	2016-17	Pooled
T <sub>1</sub>	0.648	0.980	0.814	0.724	1.007	0.866	0.883	0.803	0.843
T <sub>2</sub>	0.538	0.867	0.702	0.656	0.997	0.827	0.923	0.843	0.883
T <sub>3</sub>	0.808	1.017	0.912	0.780	1.030	0.905	0.930	0.850	0.890
T <sub>4</sub>	0.799	0.953	0.876	0.755	0.937	0.846	0.913	0.833	0.873
T <sub>5</sub>	0.782	0.870	0.826	0.744	0.990	0.867	0.920	0.840	0.880
T <sub>6</sub>	0.779	1.013	0.896	0.782	1.097	0.939	0.890	0.810	0.850
T <sub>7</sub>	0.815	0.980	0.898	0.761	0.973	0.867	0.863	0.783	0.823
T <sub>8</sub>	0.876	1.040	0.958	0.846	1.090	0.968	0.877	0.797	0.837
T <sub>9</sub>	0.845	1.010	0.928	0.858	1.103	0.981	0.913	0.833	0.873
T <sub>10</sub>	0.803	1.007	0.905	0.762	0.937	0.849	0.913	0.833	0.873
T <sub>11</sub>	0.831	0.980	0.906	0.751	0.830	0.790	0.943	0.863	0.903
T <sub>12</sub>	0.786	0.963	0.875	0.739	0.833	0.786	0.943	0.863	0.903
T <sub>13</sub>	0.696	1.027	0.862	0.586	0.893	0.740	0.850	0.770	0.810
T <sub>14</sub>	0.707	0.913	0.810	0.673	1.040	0.857	0.883	0.803	0.843
S.Em.±	0.055	0.037	0.032	0.09	0.068	0.055	0.034	0.038	0.017
C.D. at 5 %	0.159	0.107	0.093	NS	NS	NS	NS	NS	NS
C.V. %	7.47	8.26	3.12	6.73	9.87	5.64	5.16	5.15	6.63
F <sub>1</sub>	0.759	0.986	0.873	0.746	0.961	0.854	0.904	0.824	0.864
F <sub>2</sub>	0.771	0.960	0.865	0.742	1.004	0.873	0.902	0.822	0.862
S.Em.±	0.023	0.017	0.014	0.024	0.03	0.02	0.014	0.014	0.013
C.D. at 5 %	NS	NS	NS	NS	NS	NS	NS	NS	NS
T×F S.Em.±	0.085	0.064	0.09	0.088	0.027	0.079	0.051	0.052	0.051
C.D. at 5 %	NS	NS	NS	NS	NS	NS	NS	NS	9.29
C.V. %	4.21	5.44	2.01	2.58	7.10	4.23	3.91	4.87	5.37
General mean	0.771	0.959	0.890	0.769	0.874	0.832	0.904	0.822	0.863

275

276 Application of treatment 75% P as RP+AM (T<sub>8</sub>) to applied preceding  
 277 *rabi* maize increased (425.14%, 320.03 and 358.20 %) of total nitrogen  
 278 uptake, (561.54, 377.78 and 450.24%) of total phosphorus uptake and  
 279 (290.21, 147.00 and 191.62%) total potassium uptake by green gram  
 280 (grain+ stover) during 2015-16, 2016-17 and in pooled analysis respectively  
 281 over control T<sub>2</sub> (Fig-2 A, B and C).

282 This might be due the adequate supply of P supplemented by the  
 283 beneficial residual effects of the phosphorus fertilizer along with AM *viz.*,  
 284 mineralization and slow release of nutrients from organic matter like bio-  
 285 compost and higher organic carbon content and these nominated  
 286 treatments which improved physiochemical properties of soil and  
 287 synergistic effect of phosphorus fertilization resulting in higher NPK  
 288 concentration and uptake by seeds and stover. On the other hand this  
 289 because of more availability of nutrient provided from different phosphorus  
 290 management treatments; higher NPK uptake was obviously due to more  
 291 seeds and stover yield. Similar application of phosphorus fertilizer

increased uptake of NPK by green gram [9], [24] rice-green gram cropping system and [20] reported from *kharif* green gram.



**Fig-2: Total NPK uptake by summer green gram as influenced by different treatment**

(A) Total N uptake, (B) Total P uptake (C) Total K uptake

### 3.2.2 Direct effect

The data regarding NPK content in green gram seeds stover and pod cover were presented in (Table-6, Table-7, Table-8) and NPK uptake (Fig 5.14). NPK content and uptake was non-significant effect between 75% RDF ( $F_1$ ) and 100% RDF ( $F_2$ ). The results were closely related to early findings by [1], [21], [5] and [8]. in green gram they also found that application 75% RDF and 100% RDF among the NPK content and uptake by summer green gram seed and stover at par with each other.

### 4. CONCLUSION

From the current study, it can be concluded that the treatment 75%P as RP+AM (290.83, 251.36 and 266.19 %) increased total nitrogen uptake and (333.11, 345.44 and 340.35%) total phosphorus uptake by maize (grain + straw) over control  $T_2$  during both the years as well as in

\* E-mail address: xyz@abc.com.

320 pooled analysis, respectively. The total potassium uptake increased (231.1  
321 and 124.3 %) over control T<sub>2</sub> during the first year of the study and in pooled  
322 analysis, respectively and in the treatment, 75% P as SSP+AM increased  
323 (92.67 %) of total potassium uptake by maize (grain + straw) over control  
324 T<sub>2</sub> during the 2016-17 year. Application of treatment 75% P as RP+AM to  
325 applied preceding *rabi* maize increased (425.14%, 320.03 and 358.20 %) of  
326 total nitrogen uptake, (561.54, 377.78 and 450.24%) of total phosphorus  
327 uptake and (290.21, 147.00 and 191.62%) total potassium uptake by green  
328 gram (grain+ stover) during 2015-16, 2016-17 and in pooled analysis  
329 respectively over control T<sub>2</sub>. The NPK content and uptake by seeds, stover  
330 and pod of green gram were non-significant under 75% RDF (F<sub>1</sub>) and 100%  
331 RDF (F<sub>2</sub>) to summer green gram.

332 ➤ Application of phosphorus to *rabi* maize at (45kg P<sub>2</sub>O<sub>5</sub> /ha) through  
333 SSP+AM or RP (composted) +AM and application of 75% RDF (15-  
334 30-00kg N-P-K/ha) to summer green gram can achieve higher  
335 productivity of cropping system and higher nutrient content and  
336 uptake by *rabi* maize and summer green gram under South Gujarat  
337 condition.

338 ➤ Rock phosphate should be used along with AM fungi which have  
339 more beneficial role between plants and soil environment.

340

341 **Ethical approval and consent: NA**

342

## 343 REFERENCES

- 344 [1]. Ambhore, A.P. 2004. Response of summer greengram (*Vigna radiata* L.)  
345 to biofertilizers and inorganic fertilizers under south Gujarat  
346 condition. M.sc. (Agri.) thesis submitted to Navsari Agricultural  
347 University, Navsari.
- 348 [2]. Amjad, Ali., Muhammad, Sharif., Fazli, Wahid., Zengqiang, Zhang. and  
349 Syed, Noor. Muhammad Shah. 2014. Effect of composted rock  
350 phosphate with organic materials on yield and phosphorus uptake of  
351 berseem and maize. *American Journal of Plant Sciences*, 12 (5):  
352 975-984.
- 353 [3]. Anonymous .2011. Area, production and productivity of major pulses.  
354 <http://agropedia.iitk.ac.in/node/11677>.
- 355 [4]. Blake, G.R., 1965. Bulk density. In C.A. Black *et al.* (ed.) Methods of  
356 Soil Analysis, Part 1. Agronomy 9;374-390.
- 357 [5]. Jat, R.A., Arvadia, M.K., Tandel, B., Patel, T.U. and Mehta, R.S. 2012.  
358 Response of saline water irrigated green gram (*Vigna radiata*) to  
359 land configuration, fertilizer and farm yard manure in Tapi command  
360 area of South Gujarat. *Indian Journal of Agronomy* 57(3):270-274.
- 361 [6]. Jackson, M. L. 1979. Soil chemical Analysis. Prentice Hall of Englewood  
362 cliffs, New Jersey; USA.

- 363 [7]. Jat, R.S. and Ahlawat I.P.S. 2006. Direct and residual effect of  
364 vermicompost, biofertilizers and phosphorus on soil nutrient  
365 dynamics and productivity of chickpea-fodder maize sequence.  
366 *Journal of Sustainable Agricultural* 28:41-54.
- 367 [8]. Kharadi, Rahulkumar. Ramanbhai. 2014. Yield and chemical composition  
368 of summer green gram and soil properties as influenced by organic  
369 and inorganic fertilization. *M. Sc. (Agri.) Thesis* submitted to Anand.
- 370 [9]. Kuldeep, Singh. R. S., Manohar, Rakesh. Choudhary., Yadav, A.K. and  
371 Sangwan, A. 2015. Response of different sources and levels of  
372 phosphorus on yield, nutrient uptake and net returns on mungbean  
373 under rainfed condition. *Agricultural Science Digest* 35(4): 263-268.
- 374 [10]. Kumari, S. and Ushakumari, K. 2002. Effect of vermicompost enriched  
375 with rock phosphate on the yield and uptake of nutrients in cowpea.  
376 *Journal of Tropical Agriculture* 40: 27-30.
- 377 [11]. Liberato, C.G., Virgilio, A., Raquel, C.M., Nogueira, A.R.A.,  
378 Nobrega, J.A. and Schiavo, D. (2017) Determination of macro and  
379 micronutrients in plant using the Agilent 4200 MP AES. Application  
380 note Agilent Technology (<https://www.agilent.com>> application).
- 381
- 382 [12]. Lindsay, W.L. and Norvell, W.A. 1978. Development of a DTPA soil test  
383 for zinc, iron, manganese, and copper. *Soil Science Society of*  
384 *America Journal* 42:421-428.
- 385 [13]. Meena, Ramawatar. 2017. Response of Greengram (*Vigna radiata*) to  
386 rock phosphate enriched compost on yield, nutrient uptake and soil  
387 fertility in Inceptisol. *International Journal of Chemical Studies*  
388 5(2):513-516.
- 389 [14]. Mehta, Y. K., Shaktawat, M.S. and Singhi, S.M. 2005. Influence of  
390 sulphur, phosphorous and farmyard manure on yield attributes and  
391 yield of maize (*Zea mays* L.) in southern Rajasthan conditions.  
392 *Indian Journal of Agronomy* 50:203-205.
- 393 [15]. Merwin, H. D., and Peech, M. 1951. Exchangeable of soil potassium in  
394 the sand, silt and clay fraction as influenced by the nature of the  
395 complimentary exchangeable cation. *Soil Science Society of*  
396 *America Proceedings* 15:125-28.
- 397 [16]. Naseer, M. and Dost, M. 2014. Direct and residual effect of hazari  
398 rock phosphate (HRP) on wheat and succeeding maize in alkaline  
399 calcareous soil. *Pakistan journal of botany*, 46(5): 1755-1761.
- 400 [17]. Noor, Muhammad. Mashori., Mehrunisa, Memon., Kazi, Suleman.  
401 Memon. and Hidayatullah, Kakar. 2013. Maize dry matter yield and P  
402 uptake as influenced by rock phosphate and single super phosphate  
403 treated with farm manure. *Soil Environ* 32 (2):130-134.
- 404 [18]. Olsen, S. R., Cole, C. V., Watanabe, F. S., and Dean, L. A. 1954.  
405 Estimation of available phosphorus in soils by extraction with sodium  
406 biocarbonate. *U.S.D.A Circ* 93:21-29.
- 407 [19]. Panse, V.G. and Sukhatme, P.V. 1967. Statistical methods for  
408 Agricultural Workers. ICAR, New Delhi, pp. 199-200.
- 409 [20]. Patel, H.F., Maheriya, V.D.S., Attar K and Patel, H.R. 2017. Nutrient  
410 uptake and yield of Kharif green gram as influenced by levels of  
411 sulphur, phosphorus and PSB inoculation. *Legume Research* 23 (10):  
412 1-5.
- 413 [21]. Patel, R.D. 2012. Response of different cultivars of green gram (*Vigna*  
414 *radiata* L.) to integrated nutrient management under South Gujarat  
415 condition. *M.Sc. (Agri.) Thesis* submitted to Navsari Agricultural  
416 University, Navsari.

- 417 [22]. Piper, C.S., 1966. A laboratory manual of methods for the examination  
418 of soils and the determination of the inorganic constituents of plants,  
419 *waite Agriculture Research Institute University of Adelaide* 31; 59-  
420 75.
- 421 [23]. Raju, P.S., Clark, R.B., Ellis, J.R. and Maranville, J. W. 1990. Effects  
422 of species of V AM fungi on growth and mineral uptake of sorghum at  
423 different temperature. *Plant Soil* 12 (5): 165-170.
- 424 [24]. Rama, Lakshmi .C.H.S., Chandrasekhar, Rao .P., Sreelatha,T and  
425 Madhavi, M (2015) Residual effect of organic and inorganic nutrient  
426 sources on macro and micro nutrient status of rabi greengram under  
427 rice-greengram cropping system. *Legume Research* 38 (4):496-502.
- 428 [25]. Rathore, V.P. and Singh, H.P. 1999. Influence of vesicular arbuscular  
429 mucorrhizal fungi and phosphate on maize. *Journal Indian Society of*  
430 *Soil Science* 43 (2): 207-210.
- 431 [26]. Subbiah, B. V., and Asija. 1965. A rapid procedure for the estimation  
432 of available nitrogen in soils. *Current science India* 25;259-60.
- 433 [27]. Tarafdar, J.C. and Marschner, H. 1994. Efficiency of VAM hyphae in  
434 utilization of organic phosphorus by wheat plants. *Soil*  
435 *Science and Plant Nutrition* 40: 593-600.
- 436 [28]. Thakur, S.K. and Sharma, C.R. 1998. Effect of rock phosphate  
437 enrichment and Azotobacter inoculation on the transformation of  
438 nitrogen and phosphorus during composting. *Journal of the Manna*,  
439 M.C., Ghosh, P.K., Ghosh, B.N. and Singh, K.N. 2001. Comparative  
440 effectiveness of phosphate enriched compost and single  
441 superphosphate on yield, uptake of nutrients and soil quality under  
442 soybean-wheat rotation. *Journal of Agricultural Science* 137: 45-54.
- 443 [29]. Walkley, A., and Black, I. A. 1934. An examination of the digtjareff  
444 method for determining the soil organic matter and proposed  
445 modification of the chromic acid titration method. *Soil Sci* 37;29-39.