

## Soil characterization and classification of Gollarahatti-2 watershed, Karnataka, India

### Abstract:

Land resource inventorization is a method to assess the available natural resources for effective utilization. To characterize and classify the soils at large scale (1:7920 scale), this study was carried out in Gollarahatti-2 micro-watershed located in Jagalur taluk of Davanagere district, Karnataka, India. Based on the landform and physiographic units, the Gollarahatti-2 micro-watershed soils were categorized into eleven soil series and twenty two soil phases and mapping units and all the typifying soil pedons representing the study area were sampled. Morphological, Physical, Chemical and Physico-chemical properties of the identified soils were characterized under field and laboratory conditions and the soils were classified into family level as per USDA soil taxonomy. The soils were very shallow or shallow and deep, reddish brown (5 YR4/4 to very husky red (2.5YR2.5/2), slightly acidic to alkaline and non-saline. The texture of the soil was varied into sandy clay, clay loam and clay. The organic carbon ranged between low (<0.5%) to medium (0.5-0.75%). Further, the soils have high base saturation (>60%). Pedon 11 had higher exchangeable sodium percentages (>8%) in subsoil layers. The differentiated soils were grouped under 11 soil series mapped into 22 mapping units and classified into Lithic Ustorthents, Typic Haplustepts, Typic Rhodustalfs, Kanhaplic Rhodustalfs and Rhodic Kanhaplustalfs at sub group level as per USDA soil taxonomy.

**Key words:** Land resource inventory, Soil classification, Land form, Physiographic units and Soil series.

## **Introduction**

In the recent years land resources are under pressure due to degradation of soil and water, which play an important role in human as well as plant life. Soil as a medium, supports the plant growth through supply of essential nutrients and man in-turn depends on plant for food. Soil is a dynamic natural resource developed over a period of thousands of years by weathering of arable lands because of growing population, and competing demands of the various landuses. Indiscriminate use of land resources, in general, leads to their degradation and in-turn decline in productivity (Soil Survey Staff, 1999). Degradation of land resources happening at an alarming rate minimizes productivity and stability of production. Food self-sufficiency is the biggest tasks for most populous nation like India. They need to be used according to their capacity to satisfy the needs of its inhabitants. This can be achieved through proper inventory of land resources and their scientific evaluation. Soil survey provides a valuable resource inventory connected with the survival of life on earth. It provides an accurate and scientific inventory of different soils, their kind and nature and extent of distribution so that one can make prediction about their limitations and potentialities. It also provides adequate information in terms of land form, slope, land use as well as characteristics of soils *viz.*, texture, depth, structure, stoniness, drainage, acidity, salinity *etc.*, which can be utilized for the planning and development. Information of soil and related properties obtained from the soil survey and soil classification can help in better delineation of soil and land suitability for irrigation and efficient irrigation water management. So, depending on the suitability of the mapped agro-ecological units for a set of crops, optimum cropping patterns have to be suggested taking into consideration the present cropping systems and the

socio-economic conditions of the farming community (Sehgal *et al.*, 1996). Sustainable management of land resources is a good option to solve the present-day challenges (GEF council, 2005). Therefore, the knowledge of soil and land resources with respect to their spatial distribution, characteristics, potentials, limitations and their suitability for alternate land use helps in formulating strategies to obtain higher productivity on sustained basis (Vikas, 2016). This calls for systematic and reliable inventory of natural resources like soil, water, land use, *etc.*, at a quicker pace through scientific and modern tools like remote sensing and geographic information system (GIS). Satellite remote sensing data provides information on geology, geomorphology, soil and land use or land cover through synoptic and multispectral coverage of a terrain. The information generated from satellite imageries can be interpreted for various themes *viz.*, land capability, land irrigability and crop suitability *etc.* for better management and conservation of resources on watershed basis. Keeping these facts in view, the detailed soil survey of Gollarahatti-2 micro-watershed, Jagalur taluk, Davanagere district representing Central Dry Zone of Karnataka state, India was carried out with the objective of characterization and classification of Gollarahatti-2 micro-watershed, Jagalur taluk, Davanagere district, Karnataka, India.

## **Methods**

The study area is Gollarahatti-2 micro-watershed in Jagalur taluk, Davanagere district, Karnataka, India and falls under central dry zone (zone no-04) of Karnataka and agro ecological sub region of 8.2 (AESR), which receives its major annual rainfall during *kharif* season (June-September). The length of growing period is 120-150 days. The major crops growing are Ragi (*Eleusine coracana*), Maize (*Zea mays*), Ground nut (*Arachis hypogaea*) and cotton (*Gossypium sp.*) *etc.* (*Carl Linnaeus* binomial naming system). Alfisols occupy major portion of the area. The

study area located at 50 km from Davanagere district. It covers an area of 719 ha, lies between 13° 23' 42" and 31° 25' 39" N latitudes and 77° 33' 36.8" and 77° 33' 54.3"E longitudes. The elevation is in the range of 575 m to 687 m MSL. The dominant geology of the study area is Archean schist with small patches of granite gneiss. *Azadirachta indica*, *Pongamia sp.* *Mimosa pudica* and grasses are the major natural vegetation apart from forest species. Detailed soil survey was carried out by using 1:7920 (scale) cadastral map, Google Earth Image and high resolution satellite imagery of the watershed were used as base map in conjunction with Survey of India toposheet to map the land resources. Physiography soil relationship was established using ground truth data by using satellite imagery of the Gollarahatti-2 micro-watershed. Pedon sites were located in transects along the slope from the upper to lower slopes. Totally in this micro-watershed, 25 profiles were exposed and studied for morphological characteristics as per Soil Survey Manual (Soil Survey Staff, 2014). The representative 11 master profiles of typifying pedons of series identified were selected. Horizon-wise soil samples were collected, air dried and passed through 2 mm sieve and analyzed for particle-size distribution following International Pipette method (Richards, 1954), pH and electrical conductivity (EC) in 1:2.5 soil: water suspension (Piper, 1966). Organic carbon was estimated by Walkley and Black (1934) method. The cation exchange capacity (CEC) and exchangeable cations were determined as described by Jackson (1973). The soils were classified following the USDA system of soil classification (Soil Survey Staff, 2014).

## LOCATION MAP OF GOLLARAHATTI 2 MICRO-WATERSHED





Fig. 1: Location map of study area



**Fig. 2: Cadastral map with profile location of Gollarahatti -2 Micro Watershed, showing plot numbers and soil profile locations.**

## Results and discussion

### *Morphological properties*

The study area has a combination of moderately shallow (3) or shallow/very shallow soils (3) and deep (3) or moderately deep (2) soils. The pedons 6, 8, 9, 10 and 11 were in deep category, remaining all pedons (pedons 1 to 5 & 7) were in shallow group. The depth resulted shallow soils in uplands and deeper soils in lowland physiographic units. The depth of pedons were varied because of manifestation of topography. Similar observations were made by Vinay (2007) in Bhanapur micro-watershed of Koppal. The variation of depth in relation to physiography, mainly because of non-availability of adequate amount of water for prolonged period on upland soils associated with removal of finer particles and their deposition at lower pediplain. The results obtained in the present study are in agreement with the findings of

Ramprakash and Rao (2002). In all soil pedons, hue was 2.5YR-5YR. This colour hue was due to dominance of sesquioxides over silica. The darker colour values in the surface horizons (2.5-3) than sub-surface horizon (2.5-4) was due to the presence of relatively high organic matter content (Tripathi *et al.*, 2006). The sub-surface horizons had comparatively brighter colour chroma (3-6) against 3-4 of surface, which might be due to low organic matter content and higher iron oxide there. Similar kinds of results were observed in the findings of Sidhu *et al.* (1994). This variation in colour is a function of chemical and mineralogical composition, topographic position, textural makeup and moisture regimes of the soils (Thangaswamy *et al.*, 2005). The structure was sub-angular blocky in surface and sub-surface horizons. The consistency was slightly hard to hard when dry and friable to firm when moist.

#### ***Soil physical properties***

The clay content in different pedons in surface horizon ranged from 26.9 to 69.7 per cent. The sub-surface horizons exhibited higher clay content as compared to surface horizons due to the illuviation process occurring during soil development. Similarly, the illuviation process also affected the vertical distribution of silt and sand content. Similar observations were made by Dasog and Patil (2011) in soils of North Karnataka. Silt content ranged from 10.2 to 43.6 per cent. It exhibited an irregular trend with depth. This might be due to variation in weathering of parent material. These results were in agreement with the findings of Naidu and Hunsigi (2001), who observed an irregular trend in silt content with depth in sugarcane growing soils of Karnataka. Similar results were also reported by Kumar *et al.* (2002). Sand content varied from 10.2 to 54.8 per cent it was more in the surface compared to sub-surface horizons. The sand content is much higher than the silt and clay fractions. The coarser fractions dominate in siliceous, granite-gneiss parent material (Dutta *et al.*, 1999). The texture of pedons varied from

clay, clay loam, sandy clay loam to sandy clay. The textural variation might be due to different process of soil formation, in-situ weathering and translocation of clay (Srinivasan *et al.*, 2013). Water holding capacity of various pedons ranged from 36.5 to 63.1 per cent. Irrespective of the pedons, the water holding capacity of sub-soil was higher than surface soil. These differences were due to the variation in clay and organic carbon content of the pedons. Similar results were reported by Singh *et al.* (1999) in soils of Ramganga catchment in Uttar Pradesh and Thangasamy *et al.* (2005) in soils of Sivagiri micro-watershed in Chittoor district of Andhra Pradesh. Bulk density of the pedon samples varied from 1.22 to 1.41 Mg m<sup>-3</sup> (Table 1), followed a common pattern of increasing with increasing depth. It was attributed to the pressure of the overlying horizons and diminishing amounts of organic matter. Similar results were quoted by Marathe *et al.* (2003) in mandarin orchards of Nagpur and in rice soils of Eastern region of Varanasi (Singh and Agrawal, 2005).

#### ***Soil chemical properties***

The pH of red soil pedons ranged from slightly acidic to neutral and alkaline. Iron hydroxide species might have contributed for higher H<sup>+</sup> concentration leading to lower pH values. Similar observations were made by Dasog and Patil (2011) and Satyanarayana and Biswas (1970). In soils of all the pedons, EC ranged from 0.03 to 0.98 dS m<sup>-1</sup> indicating non-saline nature of soils. The soil is non saline having EC less than 1 dSm<sup>-1</sup> which might be due to removal of bases by percolation or by drainage water. These results were in confirmation with the findings of Kumar (2011) and Shivasankaran *et al.* (1993). Organic carbon content in surface horizons ranged from 0.34 to 0.72 per cent and in sub-surface horizon it varied from 0.11 to 0.6 per cent. The lower contents of organic carbon apparently resulted because of high temperature, which induced rapid rate of organic matter oxidation, while the declining trend towards



accumulation of crop residues every year, without substantial downward movement. These observations are line with the-findings of Balpande *et al.* (2007). Similar results were reported by Basavaraju *et al.* (2005) in soils of Chandragiri mandal of Chittoor district in Andhra Pradesh. The exchangeable bases in all the pedons were in order of  $\text{Ca}^{+2} > \text{Mg}^{+2} > \text{Na}^{+} > \text{K}^{+}$  on the exchange complex. From the distribution of  $\text{Ca}^{+2}$  and  $\text{Mg}^{+2}$ , it is evident that  $\text{Ca}^{+2}$  shows the strongest relationship with all the species, comparing these ions ( $\text{Ca}^{+2}$ ,  $\text{Mg}^{+2}$ ,  $\text{K}^{+}$  and  $\text{Na}^{+}$ ) it was clear that  $\text{Mg}^{+2}$  was present in low amount than  $\text{Ca}^{+2}$ . These results were in conformity with findings of Sharma (1996). The low value of exchangeable monovalents as compared to divalents was due to preferential adsorption of divalents than monovalent. These findings were in accordance with the reports of Das and Roy (1979). Cation exchange capacity of the pedons varied both location-wise and depth-wise. The values of cation exchange capacity of soils increased with profile depths and followed the trend of clay content. Similar findings have been reported by Mruthunjaya and Kenchanagowda (1993) and Shadaksharappa *et al.* (1995) in Vanivilas command and Malaprabha command area, respectively. There was a high degree of correlation between clay and CEC in red soils. The ESP ranged from 0.06 to 13.2 percent indicated initiation of the process of sodification in a downward direction. A measure of relative amounts of exchangeable sodium in comparison with the total cations in the soil are dependent on factors such as type of minerals, concentration of electrolytes and status of soluble cations (Sehgal, 1996). The findings were in accordance with the works of Srinath (1979) and Pulakeshi (2010).

The soils in the Gollarahatti-2 micro-watershed were highly base saturated. The base saturation was high in all surface horizons. In most of the soils, the base saturation increased with the depth. The increase of base saturation with the depth is due to the downward movement

of bases along with percolating water from the upper horizon to the lower horizons. Similar results were found by Sitanggang *et al.* (2006) (Table 2).

### ***Soil taxonomy***

Based on morphological characteristics of the pedons, physical, chemical characteristics (Challa, 2000) eleven pedons from the study area were classified into order, suborder, great group and sub-group (Table. 3). Pedons 2, 3, 4, 5, 6, 7, 8 and 9 have argillic sub-surface horizon and do not have plaggan epipedon and spodic or oxic sub-surface horizons above the argillic horizon. Further, the argillic horizon was developed due to clay illuvation and was identified by the presence of clay cutans and the thickness of the horizon is more than 7.5 cm and also more than one-tenth as thick as the sum of the thickness of all the overlying horizons. The base saturation was more than 35 per cent throughout the depth of the argillic horizon. Hence, Pedons 2, 3, 4, 5, 6, 7, 8 and 9 are keyed out as Alfisol at order level. Pedon 1 is classified into Entisols owing to root restricting layer within 25 cm and no diagnostic horizons either on surface or subsurface. Pedons 10 and 11 are classified into Inceptisols due to the absence of any other diagnostic horizons other than colour or texture altered cambic horizon. As the moisture regime is Ustic, Pedons 2, 3, 4, 5, 6, 7, 8 and 9 were classified as Ustalfs at sub-order level. Pedon 1 classified at sub-order level as Orthents as they are better drained than Aquents, non-fluviatile. Pedon 10 and 11 were classified as Ustepts. Pedon 5 did not have either Duripan or Calcic horizon and the base saturation was more than 60 percent at a depth between 0.2 to 0.7 m from the soil surface. These characters indicated that these pedons confirmed to the central concept of Ustalfs. So, this pedon grouped under Haplustalfs at great group level. Similarly, the pedons 10 and 11 were keyed out as Haplustepts, as they do not have Duripan, Kandic and Petrocalcic horizons. Pedon 2, 3, 5, 6, 7, 8, 9 keyed out as Rhodustalfs at great group level as they have within upper 100 cm or the

entire argillic horizon more than 50 per cent 2.5YR or redder, and values (moist)  $\leq 3$  and dry values are no more than 1 unit higher than moist values. Pedon 1 classified as Ustorthents as they have Ustic moisture regime. At the sub-group level, pedon 5 do not exhibit inter-gradation with other taxa or an extra-gradation from the central concept, hence keyed out as Typic Haplustalfs. Pedons 2, 6, 7, 9 keyed out as Typic Rhodustalfs. Pedon 10 and 11 as Typic Haplustepts, Where as pedon 1 was classified as Lithic Ustorthents due to lithic contact within 100 cm of mineral soil surface. Pedon 3 and 8 were classified as Kanhaplic Rhodustalfs, owing to a lower CEC per kg clay of less than 24 cmol(p+) kg<sup>-1</sup> in the argillic horizon. Pedon 4 as Rhodic Kanhaplustalfs, owing to the presence of kandic horizon with very low CEC per kg clay of less than 24 cmol (p+) kg<sup>-1</sup> in the argillic horizon. CEC per kg clay of less than 16 cmol (p+) kg<sup>-1</sup> in the kandic horizon with a hue redder than or equal to 2.5 YR in at least half of the depth of kandic horizon (Soil Survey Staff, 2014).

## Conclusions

Gollarahatti-2 micro-watershed soils are grouped under eleven soil series and they were characterized and mapped into 22 mapping units. These soils come under Entisol, Inceptisol, and Alfisol soil orders. Based on base saturation, organic carbon content and clay content of the soil, the soils of the study area are classified as Lithic Ustorthents, Typic Haplustepts, Typic Rhodustalfs, Kanhaplic Rhodustalfs, Rhodic Kanhaplustalfs at sub-group level. The major crops cultivated in this watershed are in the order of short duration and rainfed in a combination of pulse crop adjusting monsoon, main cereal or millet crop, followed by a very short duration oil seed crop (*Sesamum* (*Sesamum indicum*), Ground nut (*Arachis hypogaea*) or mustard (*Brassica sp.*) or coriander (*Coriandrum sativum*), utilizing the residual moisture and all based on rainfall probability. The climate is highly responsible for the crop selection. Since, the probable length of

growing period is 120-150 days, the farmers can go for deep ploughing before first showers, harrowing to keep land ready to receive and accept water reaching through rainfall and to provide crops, two subsequent short duration crops (Maize- *Zea Mays*, Sorghum- *Sorghum bicolor*, Ragi-*Eleusine coracana*) to reap higher economic benefits.

## References

- Balpande HS, Challa O, Jagdish Prasad. Characterization and classification of Grape-growing soils in Nasik district, Maharashtra. Journal of Indian Society of Soil Science.2007;55:80-83.
- Basavaraju D, Naidu MVS, Ramavatharam N, Venkaiah K, Rama Rao G, Reddy KS. Characterization, classification and evaluation of soils in Chandragiri mandal of Chittoor district Andhra Pradesh. Agropedology.2005;15:55–62.
- Challa O, Bhaskar BB, Anatwar SG, Gaikwad MS. Characterization and classification of some problematic Vertisols in semi-arid ecosystem of Maharashtra Plateau. Journal of Indian Society of Soil Science.2000;48(1):139-145.
- Das SN, Roy BB. Characterization of a catenary soil. Indian J. Agric. Chem.,1979;12:43-51.
- Dasog GS, Patil PL. Genesis and classification of black, red and lateritic soils of north Karnataka. In: Special publ. on Soil Science Research in north Karnataka, 76<sup>th</sup> Annual Convention of ISSS, Dharwad Chapter of ISSS; 2011.
- Dutta D, Sah KD, Sarkar D, Reddy RS. Quantitative evaluation of soil development in some Alfisols of Andhra Pradesh. Journal of Indian Society of Soil Science. 1999;47:311-315.
- Jackson ML. ‘Soil Chemical Analysis’. Prentice Hall of India Pvt. Ltd., New Delhi; 1973.
- Kumar AV, Natarajan S, Sivasamy R. Characterization and classification of soils of lower palanimanimuthar watershed of Tamil Nadu. Agropedology, 2002;12:97-103.

250 Kumar MD. Characterization and classification of soils of a micro-watershed on basalt parent  
 251 rock in Northern transition zone of Karnataka. M. Sc. (Agri.) Thesis, University of  
 252 Agricultural Sciences, Dharwad, Karnataka (India); 2011.

253 Marathe RA, Mohanty S, Shyam Singh. Soil Characterization in relation to growth and yield of  
 254 Nagpur mandarin (*Citrus reticulata* Blanco). Journal of Indian Society of Soil  
 255 Science.2003;51:70-73.

256 Mruthunjaya S, Kenchanagowda SK. Physico-chemical properties of some salt affected soils of  
 257 Vanivilas command area of Karnataka. Mysore Journal of Agricultural  
 258 Sciences.1993;27:349-355.

259 Naidu LGK, Hunsigi G. Application and validation of FAO-framework and soil potential rating  
 260 for land suitability evaluation of sugarcane soils of Karnataka.  
 261 Agropedology.2001;11:91-100.

262 Piper CS. 'Soil Plant Analysis'. The University of Adelaide, Australia; 1966.

263 Pulakeshi HBP, Patil PL, Dasog GS. Characterization and classification of soil resources of soil  
 264 resources derived from chlorite schist in northern transition zone of Karnataka. Karnataka  
 265 Journal of Agricultural Sciences.2014;27(1):14-21.

266 Pulakeshi HB. Characterization and classification of soil resources of Mantagani village in  
 267 Haveri district. M. Sc. (Agri.) Thesis, University of Agricultural Sciences., Dharwad,  
 268 Karnataka (India); 2010.

269 Ramprakash T, Rao SM. Characterization and classification of some soils in a part of Krishna  
 270 district Andhra Pradesh. The Andhra Agricultural Journal.2002;49:228-236.

271 Richards LA. Diagnosis and improvement of saline and alkali soils. USDA, Hand Book No 60,  
 272 101;1954.

273 Satyanarayana T, Biswas TD. Chemical and mineralogical studies of associated black and red  
 274 soils. Mysore Journal of Agricultural Sciences.1970;8:253-264.

275 Sehgal JL. Pedology – Concepts and Application. Kalyani Publishers, New Delhi; 1996.

276 Shadaksharappa GS, Patil CV, Hebsur NS. Irrigation induced changes in soil properties of  
 277 Malaprabha command area. J. Maharashtra Agricultural Universities.1995;20: 283-284.

278 Sharma JP, Landey RJ, Kalbande AR, Mandal C. Morphological, physical and chemical  
 279 characteristics and land use plans of different landforms in Pratapgarh region of Southern  
 280 Rajasthan. Agropedology.1996;9:76-83.

281 Shivasankaran K, Mithyantha MS, Natesan S, Subbarayappa CT. Physico-chemical properties  
 282 and nutrient management of red and lateritic soils under plantation crops in Southern  
 283 India, NBSS Publication, 1993;37: 280.

284 Sidhu PS, Raj Kumar, Sharma BD. Characterization and classification of Entisols in different  
 285 soil moisture regimes of Punjab. Journal of Indian Society of Soil Science.1994;42:633-  
 286 640.

287 Singh HN, Sharma AK, Om Prakash. Characterization and classification of some cultivated soils  
 288 of Ramganga catchment in the soils of Uttar Pradesh. Agropedology.1999;9:41-46.

289 Singh IS, Agrawal HP. Characterization, genesis and classification of some rice soils of Eastern  
 290 region of Varanasi, Uttar Pradesh. Agropedology.2005;15(1):29-38.

291 Sitanggang M, Rao YS, Nayan Ahmed, Mahapatra SK. Characterization and classification of  
 292 soils in watershed area of Shikohpur, Gurgaon district, Haryana. Journal of Indian  
 293 Society of Soil Science.2006;54:106–110.

294 Soil Survey Staff. Soil Taxonomy – Keys to Soil Taxonomy (Twelfth edition). United States,  
 295 Department of Agriculture, Washington DC, USA; 2014.

296 Srinath K. A study of Physico-chemical characteristics of some salt affected soils of  
 297 Tungabhadra river valley project left bank canal area. *M. Sc. (Agri.) Thesis*, University of  
 298 Agricultural Sciences Bangalore, Karnataka (India);1979.

299 Srinivasan R, Natarajan A, Anil Kumar KS, Kalaivanan D. Characterisation of major cashew-  
 300 growing soils of Dakshina Kannada district of Karnataka. *Agropedology*. 2013;23(2);59-  
 301 64.

302 Thangasamy A, Naidu MVS, Ramavatharam N, Raghava Reddy C. Characterization,  
 303 classification and evaluation of soil resources in Sivagiri micro-watershed of Chittoor  
 304 district in Andhra Pradesh for sustainable land use planning. *Journal of Indian Society of*  
 305 *Soil Science*. 2005;53:11–21.

306 Tripathi PN, Verma JR, Patil KS, Singh K. Characteristics, classification and suitability of soils  
 307 for major crops of Kiar-Nagali micro-watershed in North-West Himalayas. *Journal of*  
 308 *Indian Society of Soil Science*. 2006;54(2):131-136.

309 Vikas NK. Land resources assessment of Gollarahatti-2 micro-watershed, Jagalur taluk,  
 310 Davanagere district using remote sensing and GIS techniques. *M. Sc. (Agri.) Thesis*,  
 311 University of Agricultural Sciences, Bangalore, Karnataka (India), 2016.

312 Vinay L. Characterization and classification of soil resource of Bhanapur micro-watershed  
 313 (Koppal district) for land evaluation. *M. Sc. (Agri.) Thesis*, University of Agricultural  
 314 Sciences, Dharwad, Karnataka (India), 2007.

315 Walkley A, Black IA. An examination of the Degtjareff method for determining organic carbon  
 316 in soils: Effect of variations in digestion conditions and of inorganic soil constituents. *Soil*  
 317 *Science*. 1934;63:251-263.

318

319 **Table 1: Physical properties of Gollarahatti-2 micro-watershed pedons**

Horizon	Depth (cm)	Colour	Coarse sand (2-0.25 mm)	Fine sand (0.25-0.05 mm)	Total sand (2.0-0.05 mm)	Silt (0.05-0.002 mm)	Clay (<0.002 mm)	Texture	B.D Mg m <sup>-3</sup>	WHC (%)
			------%-----							
	Pedon 1									
Ap	0-21	5 YR 3/4	31.30	14.40	45.70	16.60	37.70	sc	1.34	39.88
	Pedon 2									
Ap	0-15	2.5 YR 2.5/4	37.60	12.10	49.70	16.60	33.70	sc	1.31	36.14
Bt1	15-30	2.5 YR 2.5/4	24.25	16.75	44.00	22.02	33.98	cl	1.35	51.00
Bt2	30-41	2.5 YR 2.5/4	32.25	9.50	41.75	23.75	34.50	cl	1.39	53.02
BC	41-50	2.5 YR 2.5/4	32.50	9.25	41.75	21.75	36.50	cl	1.41	52.16
	Pedon 3									
Ap	0-15	5 YR 3/4	37.10	12.60	49.70	12.50	37.80	sc	1.31	39.88
Bt	15-32	2.5 YR 2.5/4	12.75	21.50	34.25	22.50	43.00	c	1.34	56.33
BC	32-50	2.5 YR 2.5/4	23.20	11.30	34.50	24.65	41.10	c	1.36	59.18
	Pedon 4									
Ap	0-22	2.5 YR 2.5/4	38.20	15.60	53.80	10.40	35.80	sc	1.26	37.27
Bt1	22-32	2.5 YR 2.5/2	22.50	16.50	39.00	25.00	36.00	cl	1.32	39.76
Bt2	32-47	2.5 YR 3/6	22.50	15.40	37.90	22.60	39.50	cl	1.35	52.15
Bt3	47-60	2.5 YR 2.5/3	26.26	10.15	36.40	22.46	41.14	cl	1.36	55.45
BC	60-74	2.5 YR 3/4	25.50	10.50	36.00	23.50	40.50	cl	1.36	53.02
	Pedon 5									



Ap	0-19	5 YR 3/4	25.50	18.75	44.25	26.00	29.75	scl	1.31	33.63
Bt1	19-38	5 YR 4/4	8.82	11.75	20.57	43.23	36.20	cl	1.34	58.18
Bt2	38-54	5 YR 3/2	28.09	8.31	19.40	42.48	38.12	cl	1.36	57.51
Pedon 6										
Ap	0-24	2.5 YR 2.5/3	21.43	18.57	43.00	27.50	29.50	scl	1.27	35.24
Bt1	24-34	2.5 YR 3/6	13.75	10.50	24.25	16.25	59.50	c	1.28	59.20
Bt2	34-51	2.5 YR 2.5/4	8.75	5.75	14.50	16.75	68.75	c	1.34	61.52
Bt3	51-69	2.5 YR 2.5/4	8.75	5.25	14.00	17.25	68.75	c	1.34	62.76
BC	69-81	2.5 YR 3/4	7.75	6.25	14.00	16.28	69.72	c	1.35	63.15
Pedon 7										
Ap	0-22	2.5 YR 3/4	34.4	11.20	45.60	18.70	35.70	sc	1.22	51.96
Bt1	22-48	2.5 YR 2.5/2	6.00	12.50	18.50	23.25	58.25	c	1.29	39.09
Bt2	48-60	2.5 YR 2.5/3	5.75	6.25	12.00	18.25	69.75	c	1.34	61.52
BC	60-74	2.5 YR 2.5/3	6.00	7.61	13.61	33.00	53.39	c	1.38	62.76
Pedon 8										
Ap	0-20	2.5 YR 3/4	39.20	15.60	54.80	18.30	26.90	scl	1.26	31.02
Bt1	20-47	2.5 YR 4/6	12.10	13.77	25.87	31.79	42.34	c	1.31	57.24
Bt2	47-66	2.5 YR 2.5/3	5.87	8.74	14.61	32.06	53.33	c	1.33	60.67
Bt3	66-76	2.5 YR 2.5/4	14.09	7.52	21.61	33.32	45.07	c	1.35	56.79
Pedon 9										
Ap	0-17	2.5 YR 2.5/4	27.75	18.75	46.50	26.15	27.35	scl	1.30	33.56
Bt1	17-32	2.5 YR 2.5/4	27.50	17.00	44.50	27.25	28.25	scl	1.28	36.53
Bt2	32-55	2.5 YR 2.5/4	28.50	16.50	45.00	24.25	30.75	scl	1.31	36.98
Bt3	55-80	2.5 YR 2.5/4	38.20	15.60	53.80	10.40	35.80	sc	1.31	37.28

Bt4	80-104	2.5 YR 3/6	27.75	17.25	45.00	28.75	36.25	cl	1.33	50.13
Pedon 10										
Ap	0-30	5 YR 3/3	37.00	15.80	52.80	12.10	35.10	sc	1.29	35.47
Bw1	30-70	5 YR 3/3	22.50	17.50	40.00	24.50	35.50	cl	1.32	51.00
Bw2	70-87	5 YR 3/3	21.50	17.00	38.50	25.25	36.25	cl	1.32	55.02
Bw3	87-107	5 YR 3/3	29.50	9.75	39.25	22.25	39.50	cl	1.35	59.16
Bw4	107-142	5 YR 3/3	5.65	5.00	10.25	31.50	58.25	c	1.41	54.56
Pedon 11										
Ap	0-21	5 YR 3/4	30.1	13.40	43.50	12.40	44.10	c	1.27	52.44
Bw1	21-46	5 YR 2.5/2	35.50	5.70	41.20	10.10	48.70	c	1.31	58.14
Bw2	46-71	5 YR 3/4	23.40	15.20	38.60	8.10	53.30	c	1.32	59.65
Bw3	71-102	5 YR 3/4	1.48	17.30	18.78	43.61	37.61	cl	1.34	57.63
Bw4	102-140	5 YR 3/4	1.05	25.50	26.55	37.85	35.60	cl	1.34	55.98

320

321 **Table 2: Chemical properties of Gollarahatti-2 micro-watershed pedons**

Horizons	Depth (cm)	pH (1:2.5) Water	EC (1:25) (dS m <sup>-1</sup> )	O.C. (%)	Exch.Ca	Exch.Mg	Exch. Na	Exch. K	CEC	BS	ESP
					----- cmol (p+)kg <sup>-1</sup> -----						----%---
Pedon 1											
Ap	0-21	6.88	0.38	0.53	11.14	5.78	0.16	0.18	19.92	86.64	0.80
Pedon 2											
Ap	0-15	6.65	0.08	0.50	11.20	4.40	0.61	0.31	19.02	86.85	3.20
Bt1	15-30	7.05	0.09	0.38	12.60	3.40	0.79	0.38	21.37	80.35	3.70
Bt2	30-41	7.30	0.08	0.33	9.60	1.90	0.35	0.09	14.87	80.29	2.35
BC	41-50	7.33	0.08	0.31	12.50	0.80	0.29	0.10	15.89	86.16	1.82
Pedon 3											
Ap	0-15	6.81	0.08	0.34	11.12	2.80	0.86	0.33	19.31	78.25	4.45
Bt	15-32	7.24	0.06	0.30	12.60	3.40	0.79	0.38	21.37	80.35	3.70
BC	32-50	7.43	0.06	0.11	10.26	3.60	0.68	0.23	18.51	79.80	3.67
Pedon 4											
Ap	0-22	6.71	0.11	0.51	5.80	3.80	0.57	0.23	14.00	74.29	4.07
Bt1	22-32	6.72	0.10	0.48	8.40	5.20	0.76	0.41	17.94	82.17	4.24
Bt2	32-47	6.71	0.17	0.45	10.00	3.00	0.81	0.33	16.72	84.45	4.85
Bt3	47-60	6.75	0.14	0.39	11.00	1.20	0.74	0.28	15.55	81.99	4.76
BC	60-74	6.90	0.14	0.32	12.45	4.01	0.30	0.33	19.40	88.14	1.57
Pedon 5											

Horizons	Depth (cm)	pH (1:2.5) Water	EC (1:25) (dS m <sup>-1</sup> )	O.C. (%)	Exch.Ca	Exch.Mg	Exch. Na	Exch. K	CEC	BS	ESP
					----- cmol (p+)kg <sup>-1</sup> -----					----%----	
Ap	0-19	6.19	0.15	0.54	6.85	3.10	0.09	0.03	13.00	75.38	0.69
Bt1	19-38	6.45	0.08	0.43	7.01	3.45	0.10	0.02	15.50	68.25	0.64
Bt2	38-54	6.94	0.05	0.35	6.98	3.47	0.13	0.01	14.60	72.53	0.89
Pedon 6											
Ap	0-24	6.46	0.06	0.57	7.46	3.00	0.10	0.02	12.50	84.64	0.80
Bt1	24-34	6.27	0.09	0.55	8.00	3.40	0.21	0.20	13.85	85.27	1.51
Bt2	34-51	6.76	0.06	0.51	10.46	4.10	0.28	0.09	16.95	88.08	1.65
Bt3	51-69	7.10	0.06	0.45	11.20	4.56	0.23	0.18	18.90	85.55	1.21
BC	69-81	7.14	0.05	0.32	11.22	5.40	0.13	0.19	19.15	88.45	0.67
Pedon 7											
Ap	0-22	6.58	0.05	0.62	8.30	3.40	0.10	0.21	12.73	83.50	0.78
Bt1	22-48	6.56	0.04	0.57	8.50	2.30	0.02	0.01	15.10	71.72	0.13
Bt2	48-60	6.61	0.05	0.51	10.10	4.40	0.01	0.01	16.60	87.57	0.06
BC	60-74	6.64	0.03	0.40	10.60	3.40	0.02	0.01	17.02	87.07	0.11
Pedon 8											
Ap	0-20	6.65	0.07	0.63	10.23	3.80	0.35	0.29	17.09	84.24	1.96
Bt1	20-47	7.16	0.07	0.51	11.20	4.40	0.61	0.31	19.02	86.85	3.20
Bt2	47-66	7.90	0.15	0.51	12.60	2.60	0.48	0.31	17.90	89.30	2.67
Bt3	66-76	8.11	0.11	0.43	7.40	2.60	0.48	0.36	14.08	76.98	3.40
Pedon 9											
Ap	0-17	6.36	0.04	0.56	4.81	2.40	0.15	0.11	9.60	77.81	1.56

Horizons	Depth (cm)	pH (1:2.5) Water	EC (1:25) (dS m <sup>-1</sup> )	O.C. (%)	Exch.Ca	Exch.Mg	Exch. Na	Exch. K	CEC	BS	ESP
					----- cmol (p+)kg <sup>-1</sup> -----					----%---	
Bt1	17-32	6.45	0.04	0.53	5.20	2.50	0.26	0.09	11.01	73.11	2.36
Bt2	32-55	6.47	0.03	0.52	7.11	3.40	0.37	0.10	14.10	77.23	2.62
Bt3	55-80	6.55	0.05	0.48	8.00	3.40	0.28	0.09	14.29	83.20	1.95
Bt4	80-104	6.61	0.03	0.45	8.50	4.30	0.41	0.09	15.58	87.22	2.63
Pedon 10											
Ap	0-30	7.93	0.25	0.72	11.50	3.30	0.48	0.39	16.42	83.25	2.92
Bw1	30-70	7.87	0.20	0.69	13.53	2.50	0.58	0.31	19.89	85.06	2.91
Bw2	70-87	8.03	0.21	0.64	11.80	1.50	0.43	0.25	16.08	86.94	2.60
Bw3	87-107	8.05	0.20	0.41	12.40	5.70	0.58	0.31	22.50	88.84	2.57
Bw4	107-142	8.09	0.22	0.40	15.60	7.20	0.45	0.36	25.05	88.84	1.79
Pedon 11											
Ap	0-21	7.74	0.11	0.51	7.45	3.67	0.18	0.04	12.70	89.29	1.41
Bw1	21-46	8.13	0.55	0.43	9.18	5.32	1.44	0.04	17.10	93.45	8.41
Bw2	46-71	8.11	0.96	0.39	11.56	5.35	1.97	0.04	19.40	97.52	10.15
Bw3	71-102	8.12	0.98	0.35	9.67	4.30	2.43	0.08	18.30	90.05	13.27
Bw4	102-140	8.01	0.49	0.19	10.43	4.24	1.57	0.10	18.50	88.32	8.48

323 Table 3. Taxonomic classification of identified soil series

Sl. No	Pedon number	Order	Sub-order	Great group	Sub-group	Sub group level taxonomic classification
1.	1	Entisols	Orthents	Ustic	Lithic	Lithic Ustorthents
2.	5	Alfisols	Ustalfs	Haplic	Typic	Typic Haplustalfs
3.	2,6,7,9	Alfisols	Ustalfs	Rhodic	Typic	Typic Rhodustalfs
4.	3,8	Alfisols	Ustalfs	Rhodic	Kanhaplic	KanhaplicRhodustalfs
5.	4	Alfisols	Ustalfs	Kanhaplic	Rhodic	RhodicKanhaplustalfs
6.	10, 11	Inceptisols	Ustepts	Haplic	Typic	Typic Haplustepts

324