

Original Research Article

SENSORY EVALUATION ASSESSMENT OF BREAD PRODUCED WITH COMPOSITE FLOUR FERMENTED BY BAKER'S YEAST IN AKURE, NIGERIA

ABSTRACT

This study investigated the functional properties of bread produced from composite flour (wheat, air potato and cassava) and comparatively evaluated the sensory acceptability of bread products fermented using Baker's yeast amongst prospective consumers at the Federal University of Technology (F.U.T), Akure, Nigeria. Different blend ratios (A-F) of composite flour containing wheat, cassava and air potato flours respectively were formulated for production of bread fermented using Baker's yeast. Proximate and mineral analysis of cassava, wheat and air potato flours were carried out according to standard procedures set by A.O.A.C., 2012. A descriptive sensory evaluation was carried out to assess bread products (A-F) produced from the different composite flour blends as sensory evaluators were selected by cross sectional simple random sampling techniques. The organoleptic parameters accessed for the bread products include taste, aroma/texture, appeal/appearance and overall consumer acceptability index respectively. Bread products A-C were adjudged standard controls for Air potato flour (A), Wheat flour (B) and Cassava flour (C) while products D-F were the test bread products of varying formulation blends of the composite flours. All the bread products (A-F) were evaluated and compared statistically at $p \leq 0.05$ level of significance. Bread product F had the highest overall acceptability ($99.90 \pm 0.10\%$) followed by E ($88.89 \pm 1.11\%$) and D at ($55.56 \pm 1.44\%$) respectively; the findings of this research

revealed realistic potentials of air potato flour supplemented with wheat and cassava flours respectively to produce widely accepted bread products in Nigeria.

Keywords: Air potato, Composite flour, Baker's yeast, Sensory evaluation, overall acceptability.

INTRODUCTION

Bread is a staple food in Nigeria and all over the world, with an exponential increase in consumption over recent years [1-3]. The rising cost of bread products in Nigeria is chief due to the importation of wheat flour, one of the major ingredients used in local bread production [2-4]. Wheat flour is suitable for bread production because it contains high levels of sugar and fiber ~~contents~~, high content of gluten: a protein which aids excellent formation of bread dough during fermentation, and moderate mineral elements constituents [2-5]. However, many locally made Nigerian flours from tuber crop products such as cassava and cocoyam lacks adequate levels of gluten protein to enable them compete effectively as suitable alternatives to wheat flour [4-6]. There have been therefore lots of research efforts centered on discovering suitable alternatives to wheat flour that can be used to bake bread, thereby reducing the demand for wheat and also improve the economic value of locally made flour products in Nigeria [3-6]. While research efforts in the past were originally tailored towards total substitution of wheat flour by several locally made flours from starch derived Nigerian crops suitable for bread production, efforts are now channeled towards supplementing wheat flour with these same locally derived flours in different blends and formulations in a bid to improve the nutritional value and consumer acceptability of these bread products in Nigeria [6-9].

Cassava (*Manihot esculenta* Crantz) a common root tuber is consumed extensively in Nigeria [5-9]. Nutritionally, cassava is a major source of dietary energy for low income citizens in

many parts of Nigeria because of its cheap availability [7-9]. Despite being a cheap source of food, cassava is nutritionally deficient in essential amino acids but rich in arginine [8, 9]. Its use in the production of bread as composite flour has been reported ~~in~~ [1, 4-6, 8], although the bread products obtained generally had low acceptability [1, 6, 8]. Air potato (*Dioscorea bulbivora*) ~~however~~ is a root tuber crops in the family Dioscoreacea found in many Nigerian settlements but hardly consumed due to several cultural considerations and customs despite its immense health benefits [8, 10]. Moreover, this yam specie has attracted scientific interests owing to its numerous therapeutic applications which have been attributed to its unique phytochemistry in many research findings [6, 10]. ~~Discorea~~ *bulbivora* has been reported to be a good source of protein (gluten), lipids and crude fibers; making it an alternative suitable flour source for bread production [4-6, 8, 10].


In Nigeria, baked products are fermented by Baker's yeast (*Saccharomyces cerevisiae*) as different findings have shown its effectiveness in qualitative fermentation of baked products for consumption [1-3, 4-8]. The use of composite flour containing flour blends of cassava, air potato and wheat for production of bread products fermented by Baker's yeast have the potential to conserve the foreign exchange spent on wheat importation and also add value to indigenous crops like air potato and cassava locally grown in Nigeria [1, 4-10]. Hence, this study investigated the functional properties of bread produced from composite flour (wheat, air potato and cassava) and comparatively evaluated the sensory acceptability of bread products fermented using Baker's yeast amongst prospective consumers at the Federal University of Technology (F.U.T), Akure, Nigeria.

68

69 MATERIALS AND METHODS

70 Production of Composite flour

71 The methods described in the findings of [11-14] were adopted for preparing dried
72 unfermented cassava and air potato flours as indicated in figure 1 and 2 below. Already prepared
73 dried wheat flour was obtained from commercial vendors at the main market of the Akure
74 metropolis.



83 **Figure 1:** Flowchart for production of Cassava Flour [13]

84 Proximate analysis of composite flour

The protocols described by the Association of Official Analytical Chemist (A.O.A.C, 2012) were adopted to determine the proximate and mineral compositions of the wheat flour, cassava flour and air potato flour respectively [1, 4, 15]. The moisture content, ash content, crude fiber content, crude protein content, fat content, carbohydrate content, water and oil absorption capacity and swelling index were all determined using the A.O.A.C, 2012 protocols as described in [14, 15]. The three flour types were also analyzed for mineral elements compositions (calcium, iron, potassium, manganese, sodium and zinc contents) respectively [1, 8, 12-14].




Figure 2: Flowchart for production of Air Potato Flour [14]

Formulation of Composite Flour Blends

Modifications of the methods described in [4, 5, 8, 10, 14] were adopted as different composite flour blends were obtained using simple ratios of Air Potato flour, Cassava flour and Wheat flour respectively to obtain different composite flour blends. The six flour ratios obtained were 100% Air Potato Flour (A), 100 % Wheat flour (B), 100 % Cassava flour (C), Air Potato flour/Wheat flour (50%/50%) (D), Air Potato flour/Cassava flour (50%/50%) (E) and Air Potato flour/Wheat flour/Cassava flour (40%/40%/20%) (F) respectively. The blend ratios are: 200g of Air potato flour for blend A, 200g of Cassava flour for blend B, 200g of Wheat flour for blend C, 100g each of Air potato flour and Wheat flour for blend D, 100g each of Air potato flour and Cassava flour for blend E and 80g of Air potato flour, 80g of Wheat flour and 40g of Cassava flour for blend F respectively.

Table 1: Composite flour combinations of different bread types

Bread Type	Flour Composition (percentage by mass)
A	100% Air potato flour (200g)
B	100% Wheat flour (200g)
C	100% Cassava flour (200g)
D	50% Air potato flour (100g) + 50% Wheat flour (100g)
E	50% Air potato flour (100g) + 50% Cassava flour (100g)
F	40% Air potato flour (80g) + 40% Wheat flour (80g) + 20% Cassava flour (40g)

Production of Bread from different composite flour combinations

The different composite flour combinations ~~A, B, C, D, E and F~~ were all baked into bread products by yeasts isolated from both palm wine and stale bread following standard procedures of

dough mixing, fermentation, punching, scaling, moulding, proofing, baking, cooling and de-panning respectively as ~~described separately in the findings of~~ [1, 2, 4-6, 13, 16]. The products were labeled (A–F) and lyophilized Baker’s yeast (*Saccharomyces cerevisiae*) was obtained from commercial Bakeries in Akure metropolis for fermentation of the flour blends ~~(A–F)~~.

Description of Study Area used for Sensory Evaluation

The Federal University of Technology, Akure is located in Ondo State, Nigeria with coordinates 7°16’ N 7°18’ N/ 5°9’ E 5°11’ E ~~[17]~~. It is located at the extreme southern region of the Akure South Local Government Area of Ondo state, Nigeria [17].

Sensory Evaluation of Bread samples

Simple random sampling techniques were adopted for obtaining 18 students as evaluators in the study area (Federal University of Technology, Akure Campus) to assess the organoleptic properties and the overall consumer acceptability of the different bread samples produced ~~(A–F)~~. The organoleptic properties assessed include the taste, the aroma/texture, appeal/appearance of the products and overall acceptability by the consumers [1-8, 13, 14, 16, 18]. The evaluators assessed the bread products separately at different time intervals to avoid bias; ~~a method described in the findings of [4-8]~~. Each evaluator rated the bread products independently on a hedonic scale ranging from 7 (moderately liked) to 9 (extremely liked) [13, 14, 16]. The bread samples A, B, and C served as controls while bread products D, E, and F were the test products of interest. All bread products were nevertheless rated by the evaluators. The raw scores were expressed in percentages and analyzed statistically ~~using the methods described in~~ [4-8, 18].

Data analysis

Analyzed bread products were in triplicates; data means obtained for evaluation ratings of the organoleptic properties of the bread products were subjected to a 2-way analysis of variance and the means were separated using Duncan's New Multiple Range test at $P \leq 0.05$ level of significance [4-8, 16, 18].

RESULTS

The comparative proximate compositions (Carbohydrate, Ash, Fat, Fiber, Moisture and Protein contents) and mineral compositions (Calcium, Iron, Potassium, Manganese, Sodium and Zinc contents) of the different flours (Air potato flour, Wheat flour and Cassava flour) used in composite flour formulations were analyzed separately and reported at $P \leq 0.05$ levels of significance as represented in figures 3 and 4. Air potato flour has the highest carbohydrate content ($90.35 \pm 1.58\%$) while Cassava flour has the lowest content ($37.58 \pm 1.64\%$); The Ash contents of Air potato flour was also the highest with ($2.51 \pm 0.28\%$) while the ash contents of wheat and cassava flours respectively are not significantly different at the specified level of confidence. While the fat contents of the three flours are however not significantly different at $P \leq 0.05$ level of significance, wheat flour has the highest fiber content ($2.50 \pm 0.25\%$) while the fat contents of air potato and cassava flours are not significantly different at the specified level of significance, Cassava flour has the highest moisture content ($61.40 \pm 2.20\%$) while wheat flour has the lowest moisture content ($3.67 \pm 1.10\%$); alternatively, wheat flour has the highest protein content of $11.37 \pm 1.21\%$ while cassava flour is very low in protein content with just $1.97 \pm 0.28\%$ concentration.

Moreover, the comparative mineral compositions of the flour samples also gave an insight into their nutritive suitability for bread production. Air potato was discovered to have the highest

calcium content at $52.30 \pm 2.20\%$ while Cassava has the least concentration at $20.00 \pm 1.50\%$. The iron, zinc and manganese concentrations of all the three flours were not significantly different at the specified level of significance while wheat flour has the highest potassium concentration at $87.50 \pm 2.50\%$ and cassava has the lowest at $30.2 \pm 1.80\%$. On the other hand however, the sodium concentration of air potato flour was the highest ($89.40 \pm 1.69\%$) while wheat has a comparatively low concentration at $1.73 \pm 0.24\%$. The results above signified an important edge that air potato flour possess as a suitable alternative for bread production compared to wheat flour while cassava flour is comparatively a poor alternative for wheat flour.

Furthermore, the sensory evaluation results gave useful information on the preferences of consumers to both the control bread products (A-C) and the test bread products (D-F). The evaluation indexes of the bread products (A –F) fermented by Baker’s yeast are represented in Tables 2 with respect to the organoleptic properties (taste, appeal/appearance, texture/aroma) and the overall acceptability of the bread products are represented in Table 3 by adopting the methods described in [4-6, 8].

The evaluation scores of the test bread products represented in Table 2 are expressed in percentages of the 18 evaluators at $p \leq 0.05$ level of significance and the evaluation scores of the hedonic scale 7-9 represents the acceptability of the bread products by the evaluators; a standard described in [4-6, 8, 10-14]. Products F has the highest evaluation score $88.89 \pm 1.11\%$ for taste of the test bread products followed by E at $77.87 \pm 1.23\%$, and D at $66.67 \pm 1.33\%$ respectively (Table 2). The evaluation scores for the aroma/texture of the bread products were also similar to that of the taste as products F has an evaluation score of $99.90 \pm 0.10\%$ followed by E at $66.67 \pm 1.33\%$ and D has a moderate evaluation score of $55.56 \pm 1.44\%$ (Table 2 and 3).

Conversely, the evaluation scores of the appeal/appearance of the test bread products also followed the same pattern as bread product F had the highest evaluation score in this category at 99.90±0.10% while D had a score of 66.67±1.33%, while E had a score of 44.45±1.54% respectively (Table 2). The overall acceptability of the test bread products were also estimated by the evaluators and analyzed at $p \leq 0.05$ levels of significance as bread product F had excellent overall acceptability at a score of 99.90±0.10% while the test product E had an overall acceptability at 88.89±1.11% and product D had its overall acceptability index at 55.56±1.44% respectively as represented in Table 3.




Figure 3: Comparative proximate composition of Air potato flour, Wheat flour and Cassava Flour used in composite bread production




Figure 4: Comparative Mineral composition of Air potato flour, Wheat flour and Cassava Flour used in composite bread production

212 Table 2: Sensory evaluation scores of the bread products A- F fermented by Baker's yeast

H.Scal	Sensory Evaluation Scores of Bread Products for Taste (%)					
e	A	B	C	D	E	F
7	00±00 ^a	33.33±1.69 ^d	55.56±1.44 ^f	11.11±1.89 ^b	44.45±1.55 ^e	11.11±1.89 ^b
8	33.33±1.69 ^d	00±00 ^a	22.22±1.78 ^c	44.45±1.55 ^e	33.33±1.69 ^d	11.11±1.89 ^b
9	00±00 ^a	00±00 ^a	00±00 ^a	11.11±1.89 ^b	00±00 ^a	66.67±1.33 ^g
H.Scal	Sensory Evaluation Scores of Bread Products for Appeal/Appearance (%)					
e	A	B	C	D	E	F
7	22.22±1.78 ^c	11.11±1.89 ^b	11.11±1.89 ^b	00±00 ^a	66.67±1.33 ^g	33.33±1.69 ^d
8	22.22±1.78 ^c	00±00 ^a	44.45±1.55 ^e	44.45±1.55 ^e	00±00 ^a	33.33±1.69 ^d
9	00±00 ^a	00±00 ^a	00±00 ^a	11.11±1.89 ^b	00±00 ^a	33.33±1.69 ^d
H. Scale	Sensory Evaluation Scores of Bread Products for Texture/Aroma (%)					
	A	B	C	D	E	F
7	11.11±1.89 ^b	22.22±1.78 ^c	22.22±1.78 ^c	11.11±1.89 ^b	44.45±1.55 ^e	33.33±1.69 ^d
8	22.22±1.78 ^c	11.11±1.89 ^b	33.33±1.69 ^d	55.56±1.44 ^f	00±00 ^a	33.33±1.69 ^d
9	22.22±1.78 ^c	11.11±1.89 ^b	33.33±1.69 ^d	00±00 ^a	00±00 ^a	33.33±1.69 ^d

213

214 Keys: H. Scale- Hedonic scale, 7- moderately liked, 8- like very much, 9- liked extremely, A-

215 100% air potato flour fermented by Baker's yeasts, B- 100% wheat flour fermented by Baker's

216 yeasts, C- 100% cassava flour fermented by Baker's yeasts,, D- 50% air potato flour + 50% wheat

flour fermented by Baker's yeasts, E- 50% air potato and 50% cassava flour fermented by Baker's yeasts, F- 40% air potato flour + 40% wheat flour + 20% cassava flour fermented by Baker's yeasts, values with the same superscript have no significant difference at $p \leq 0.05$ level of significance. Bread A- C is control bread products respectively while D- F are test bread products

Table 3: Overall acceptability of bread products A to F

~~Overall acceptability Bread Products fermented by Baker's yeast (%)~~					
A	B	C	D	E	F
11.11 $\pm$ 1.33 ^a	11.11 $\pm$ 1.33 ^a	77.78 $\pm$ 1.22 ^c	55.56 $\pm$ 1.44 ^b	88.89 $\pm$ 1.11 ^d	99.90 $\pm$ 0.10 ^e

Keys: A- 100% air potato flour fermented by Baker's yeasts, B- 100% wheat flour fermented by Baker's yeasts, C- 100% cassava flour fermented by Baker's yeasts,, D- 50% air potato flour + 50% wheat flour fermented by Baker's yeasts, E- 50% air potato and 50% cassava flour fermented by Baker's yeasts, F- 40% air potato flour + 40% wheat flour + 20% cassava flour fermented by Baker's yeasts, values with the same superscript have no significant difference at $p \leq 0.05$ level of significance. Bread A- C is control bread products respectively while D- F are test bread products

DISCUSSION

It was generally observed that variations in the nutritional constituents of the different flour blends used had a direct relationship on the quality of bread products produced since all the bread products were fermented by ~~Baker's yeast~~ (*Saccharomyces cerevisiae*); this was also similar in the findings of [4-8]. Several recent Nigerian literature texts ~~from research findings in [1, 4, 6, 9, 10, 14]~~ have revealed that different species of *Saccharomyces* ferment different carbohydrate substrates and produce varying degrees in textures, appeal and aroma of the products fermented.

~~The findings of [2, 4, 13, 19]~~ suggests that products fermented by the Baker's yeast (isolates of *Saccharomyces carlsbergensis* and *Saccharomyces cerevisiae*) generally possess better dough qualities, better impacted taste flavors and favorable overall acceptability indexes; this mainly because these yeast species are highly osmophilic and hydrophilic.

Consequently, the bread products F and E were the most acceptable of the test bread products and a closer observation at the composite flour blends of these two products (composite blends E and F) reveals that the two blends (E and F) contains high sugar rich contents with blends E (50% Air potato/50% Cassava) and F (40% Air potato, 40% wheat, 20% cassava) having high combined levels of sugar, fibers and protein levels; hence encouraging optimal dough formation of the bread products F and E by the ~~Baker's yeast (*Saccharomyces cerevisiae*)~~; as justified in the findings of [10, 14, 21-23].

Similarly, the product D also had a moderate overall acceptability index, the composite blend D is composed of 50% air potato and 50% wheat flour. However, since wheat flour has moderate levels of some essential mineral elements and fiber [24-27], and since air potato is supplemented in the blend, a moderate dough quality is justified as reflected in the result. Interestingly, the composite blend C (100% Cassava flour) which is rich in high mineral element and fiber content levels as reflected in the proximate and mineral elements evaluation in the results section had also had a high overall acceptability index compared to flour blends A and B respectively; this further highlights the nutritional preferences of the Baker's yeast used in fermenting the bread products; ~~this was also described in the findings of [1, 2, 4, 6, 24-26].~~

CONCLUSION

The high acceptability indexes obtained from the different bread products containing air potato and cassava flours in the findings of this research consolidated recent efforts directed at discovering other locally processed flours that can compete favorably with wheat flour in the Nigerian baking industry. The findings of this research however revealed realistic potentials of air potato flour supplemented with wheat and cassava flours respectively to produce widely accepted bread products in Nigeria.

REFERENCES

1. Udofia, P., Udoudo, J., and Eyen N. Sensory evaluation of wheat-cassava-soybean composite flour (WCS) bread by the mixture experiment design. *African Journal of Food Science*, 2013; 7(10): 368-374.
2. Eddy, N., Udofia, P., Eyo, D. Sensory evaluation of wheat/cassava composite bread and effect of label information on acceptance and preference in Benin, Nigeria. *African Journal of Biotechnology*, 2007; 6(16):123-134.
3. Dendy, D. Review of composite flour technology in the context of Tanzania. National Resources Institute, Chatham, U. K. 2013; 1-23.
4. Akanbi, T., Nazamid, S., Adebawale, A., Farooq, A., and Olaoye, A. Breadfruit starch-wheat flour noodles: preparation, proximate compositions and culinary properties. *International Food Research Journal*, 2011; 18: 1283-1287.
5. Trejo-González, S., Loyo-González, A., and Munguía-Mazariegos, R. Evaluation of bread made from composite wheat-sweet potato flours. *International Food Research Journal*, 2014; 21(4): 1683-1688.

6. Aniedu, C., and Agugo, U. Acceptability of bread produced from Hausa-potato and sweet potato composite flours in Lagos, Nigeria. *Journal of Agriculture and Social Research*, 2010; 10(2): 162-166.
7. Hathorn, S., Biswas, M., Gichuhi, P., and BovellBenjamin, A. Comparison of chemical, physical, micro-structural, and microbial properties of breads supplemented with sweet potato flour and high gluten dough enhancers. *Lebensmittel Wissenschaft und Technologie*, 2008; 41(5): 803-815.
8. FIIRO 2006. Cassava production, processing and utilization in Nigeria. Federal Institute of Industrial Research, Oshodi, Lagos, Nigeria.
9. Osundahunsi, O., Fagbemi, T., Kesselman, E., and Shimoni, E. Comparison of physicochemical properties and pasting characteristics of flour and starch from sweet and air potato cultivars in Nigeria. *Journal of Agricultural and Food Chemistry*, 2003; 51(8): 2232-2236.
10. Oviahon, I., Yusuf, S., Akinlade, J., and Balogun, O. Determinants of Bread Consumers' Willingness to Pay for Safety Labels in Oredo Local Government Area, Edo State, Nigeria. *New York Science Journal*, 2011; 4(9): 18-24.
11. Eduardo, M., Svanberg, U., Oliveira, J., and Ahrne, L. Effect of cassava flour characteristics on properties of cassava-wheat-maize composite bread types. *International Journal of Food Science*, 2013; 2:1-10.
12. Dhingra, S., and Jood, S. Physico-chemical and nutritional properties of cereal-pulse blends for bread making. *Nutritional Health*, 2002; 16(3): 183-94.

13. Igbabul B., Num, G., and Amove, J. Quality Evaluation of Composite Bread Produced from Wheat, Maize and Orange Fleshed Sweet Potato Flours in Makurdi, Nigeria. *American Journal of Food Science and Technology*, 2014; 2(4):109-115.
14. Nwosu, J., Owuamanam, C., Omeire, G., and Eke, C. Quality parameters of bread produced from substitution of Wheat flour with Cassava and Soybean as improvers. *American Journal of Research Communication*, 2014; 2(3): 99-118.
15. A.O.A.C. Official Methods of Analysis for food crops, Association of Analytical Chemists, Adingbon, Virginia. 2012, 1(3): 8-45.
16. Abdelghafor, F., Mustafa, I., Ibrahim, H., and Krishnan, G. Quality of bread from composite flour of sorghum and hard white winter wheat. *Advance Journal of Food Science and Technology*, 2011; 3: 9-15.
17. Agbelade, A., and Akindele, S. Land Use Mapping and Tree species diversity of Federal University of Technology (F.U.T.), Akure. *American International Journal of Contemporary Research*, 2013; 3(2): 104-113.
18. Olaoye, O., Onilude, A., and Idowu, O. Quality characteristics of bread produced from composite flours of wheat, plantain and soybeans. *African Journal of Biotechnology*, 2006; 11: 1102-1106.
19. Noorfarahzilah, M., Lee, J., Sharifudin, S., Mohd Fadzelly, B. and Hasmadi, M. Applications of composite flour in development of food products. *International Food Research Journal*, 2014; 21(6): 2061-2074.
20. Hasmadi, M., Siti Faridah, A., Salwa, I., Matanjun, P., Abdul Hamid, M. and Rameli, A. The effect of seaweed composite flour on the textural properties of dough and bread. *Journal of Applied Phycology*, 2014; 26:1057–1062.

21. Husniati, D., and Anastasia, F. Effect of the addition of glucomannan to the quality of composite noodle prepared from wheat and fermented cassava flours. *Journal Basic Application Science Resources*, 2013; 3(1): 1-4.
22. Oni, O., Oladele, O. and Inedia, O. Consumer willingness to pay for Food Safety labels in Nigeria: A Case study of Potassium Bromate in bread in Ibadan, Nigeria. *Journal of Central European Agriculture*, 2005; 6(3): 382-387.
23. Greene, J., and Bovell-Benjamin, A. Macroscopic and sensory evaluation of bread supplemented with sweet-potato flour. *Journal of Food Science*, 2004; 69(4): 167-173.
24. Odebode, S., Egeonu, N., and Akoroda, M. Promotion of Sweet and Air potato for the food industry in Nigeria. *Bulgarian Journal of Agricultural Science*, 2008; 14: 300-308.
25. Adenaike, O., Ameh, J., and Whong, C. Comparative studies of the fermentative capacity of Baker's yeast and local yeast strains isolated from fermented beverages and stale bread; *Nigerian Mycological Society Journal*, 2006; 5(1): 18-29.
26. Ezenronye, O., and Okerentugba, P. Genetic and Physiological variants of yeast selectively identified from local palm wine in Nsukka, Nigeria. *Mycopathologia*, 2001; 15(2): 85-99.
27. Malomo, S., Eleyinmi, A., and Fashakin, J. Chemical composition, rheological properties and bread making potentials of composite flours from breadfruit, breadnut and wheat in Akure, Nigeria. *African Journal of Food Science*, 2011; 5(7): 400 – 410.