

Original Research Article

Response of weaner rabbits fed toasted sickle pod (*Senna occidentalis*) seed meal

Abstract

This study was conducted to determine the effect of toasted sickle pod seed meal (TSSM) on the growth performance and carcass characteristics of weaner rabbits. Forty weaner rabbits were allotted to the five dietary treatments of eight rabbits per treatment and replicated four times with 2 rabbits per replicate in a Completely Randomized Design (CRD). Five experimental diets were compounded using TSSM at 0%, 2.5%, 5% 7.5% and 10% inclusion levels designated as treatments 1, 2, 3, 4 and 5, respectively. Results showed that all the growth parameters measured were not significant ($P>0.05$) across the dietary treatments as the level TSSM increased in the diet. Total Feed intake (TFI) ranged from 3201.20 to 3235.40g/rabbit while the total weight gain (TWG) varied from 930.00 g in T3 to 936.67g in T4. Better feed conversion ratio was observed in the rabbit fed experimental diets. There was no significant difference ($P>0.05$) in carcass characteristics and internal weight organs evaluated. It was concluded that toasted sickle pod seed meal (TSSM) can be included in rabbit diet up to 10% without negative effect on growth performance and carcass characteristics.

Keywords: Rabbit, sickle pod seed, processing and carcass

1.0 INTRODUCTION

There is global awareness on the shortage of animal protein supply in most developing countries [1]. Long production intervals, feed shortage, poor genetic make-up and disease incidence among other factors affecting livestock production in the tropics [2]. There is therefore an urgent need to search for fast maturing animal such as rabbit. Rabbit has immense potentials and good attributes which include high growth rate, high efficiency in converting feeds to meat, short gestation period, high prolificacy and relatively low cost of production. It's production is becoming more popular in most developing countries [3]. However, high cost feedstuff most especially the conventional protein ingredients like soybean meal and groundnut cake feeding are among other factors rabbit production in the tropics [4]. A possible solution to this problem is to explore alternative protein sources, which are cheap and locally available. One of the alternative sources considered in this study is sickle pod (*Senna occidentalis*) seed.

Senna occidentalis belong to the leguminosae family, and it is distributed throughout the tropical and subtropical regions of the world [5]. It mostly found in an open pastures and in fields cultivated with cereals such as soybean, corn, sorghum among crops [6]. The nutrient composition showed that the seed has crude protein of 18.64% and 29.54% reported by [7] and [8] respectively

which showed it has potential as a protein source in livestock feedstuffs. Studies conducted on the use of raw and soaked sickle pod seed resulted in decrease in feed intake, total weight gain and impaired nutrient utilization in rabbits and broiler chicken. There is scanty information on the use of toasted sickle pod seed meal in rabbit production. Therefore, this study was carried out to determine the effects of toasting sickle pods on the performance of weaner.

2.0 MATERIALS AND METHODS

2.1 Study location

The study was conducted at the Rabbit Research Unit of the Department of Animal Production and Technology, College of Agriculture, Ganye Local Government Area, Adamawa State. The study area lies between Latitude 8.26' and 11.98' North of the Equator and Longitude 12° 3' East of the Greenwich Meridian. The area falls within the Northern Guinea Savannah Zone and has a tropical wet and dry climate [9].

2.2 Collection and Processing of Sickle pod meal

Sickle pod seeds were sourced within the study area, cleaned and subjected to toasting. Toasting was achieved by constant stirring the seed for 30minute in a metallic frying pan to maintain uniform heating until the seed turns to light brown. The seeds were allowed to cool then mill and tagged as toasted sickle pod seed meal (TSSM)

2.3 Experimental animals, design and management

Forty weaner rabbits with an initial average weight of 709 ± 1.89 g between 6-7 weeks old procured within Yola metropolis, Adamawa State Nigeria. The weaner rabbits were allotted to five dietary treatments. Each treatment was replicated four times with two rabbits per replicate in completely randomized design. One week before commencement of the experiment, rabbit house was fumigated, cages, feeders and drinkers were thoroughly washed, disinfected and were tightly fitted to the cages to prevent spillage and feed wastage. The experimental animals weretreated against ectoparasites and endoparasites using Ivomectin.

2.4 Experimental diets and Treatments

The milled TSSM was used to compound five diets. Diet1 served as control with 0% TSSM while diets, 2,3,4 and 5 contains 0%, 2.5%, 5% 7.5% and 10% designated T1, T2, T3, T4 and T5 respectively. The ingredient composition of the experimental diets is shown in Table1.

2.5 Collection of data

The experiment lasted for 56 days during which the weaner rabbits were offered experimental diets and water *ad libitum*. The rabbits were weighed at the beginning of the experiment and subsequently on a weekly basis. Parameters measured include total weight gain, total feed intake, feed conversion ratio (FCR) and mortality.

2.6 Carcass and Organs Evaluation

A total of 4 rabbits from each treatment (one per replicate) were randomly selected, weighed, starved overnight to clear the gut and then slaughtered. After slaughtering, external offal such as the pelt head and feet were removed and weighed. During evisceration, the internal organs and other gut contents were carefully removed and weighed. Carcass was weighed; organ weights were expressed as percentage live weight. Dressing percentage was calculated thus: Dressing percentage (D %)

$$\frac{\text{dressed weight}}{\text{Live weight}} \times 100$$

2.7 Laboratory analysis

The proximate composition of the raw and toasted sickle pod seed meals, experimental diets and anti-nutritional factors were determined using the standard procedure of analysis described by [10].

2.8 Statistical analysis

Data collected were subjected to one way analysis of variance using completely randomized design [11] and means were separated using Duncan's multiple range test at the $p < 0.05$ level [12].

Table 1 composition of experimental diets

Ingredient	Inclusion levels of toasted sickle seed meal				
	T1	T2	T3	T4	T5
Maize	41.01	39.34	37.69	36.05	34.40
Soybean meal	20.00	19.16	18.31	17.45	16.60
TSSM	0.00	2.50	5.00	7.50	10.00
Cowpea Husk	10.00	10.00	10.00	10.00	10.00
Maize offal	23.00	23.00	23.00	23.00	23.00
Fish meal	2.00	2.00	2.00	2.00	2.00
Bone meal	3.00	3.00	3.00	3.00	3.00
Salt	0.50	0.50	0.50	0.50	0.50
Premix*	0.50	0.50	0.50	0.50	0.50
Total	100	100	100	100	100
Determined analysis					
Dry matter	90.95	90.92	90.90	91.00	90.95
Crude protein	18.77	18.52	18.33	18.90	18.20
Crude fibre	8.02	8.24	8.60	8.53	8.78
Ether extracts	7.86	7.17	7.65	7.01	7.50
Ash	7.91	7.23	7.76	7.41	7.88

NFE	57.44	58.84	57.66	58.15	57.64
*ME(Kcal/Kg)	3370.27	3410.72	3344.79	3293.63	3327.12

Vitamin-mineral premix provider per kg the following: Vit. A 1500 IU; Vit.D₃ 3000 IU; Vit.E 30 IU; Vit.K 2.5mg; Thiamine B₁ 3mg; Riboflavin B₂ 6mg; Pyrodoxine B₆ 4mg; Niacin 40 mg; Vit. B₁₂ 0.02mg; Pantothenic acid 10mg;Folic acid 1mg; Biotin 0.08mg; Chloride 0.125mg; Mn 0.0956g; Antioxidant 0.125g; Fe 0.024g; Cu 0.006g; 10.0014g; Se 0.24g; Co 0.240g

*Metabolizable Energy = ME (kcal/kg) = 37 x % CP + 81 x % EE + 35.5 x % NFE.
Calculated according to the formula of [13]

3.0 RESULTS AND DISCUSSION

3.1 Proximate and anti-nutritional factors of raw and toasted sickle seed meal

The proximate composition of the raw and toasted sicklepod seed are presented in Table 2. The results showed higher Crude Protein (CP), Crude Fibre (CF) and ether Extract (EE) and ash values for toasted sickle pod seeds meal while nitrogen free extracts (NFE) and metabolisable energy were not improved by toasting. This indicates that the nutrient of sickle podseed is affected by heat treatment. The CP values of both raw and toasted sicklepod seedhowever differs from the value of 18.46% reported by [7] and (2011) and 29.54% reported by [8].The crude fibre (CF) and ether extracts(EE) values ranged from 10.18-17.10% and 7.50-8.00% respectively The values of CF and ash are higher than2.45% and 9.00% reported by [7and14]. Nitrogen free extract for raw and toasted sickle seed meal were 46.77% and 33.25% respectively. Differences in the nutrient composition could be as a result of laboratory analysis, processing methods, edaphic factors and climatic factors.All the values for anti- nutritional factors(ANFs) determined decreased as a result of the toasting. The higher Crude Protein (CP) level of the toasted sickle seed may be attributed to the processing methods employed. This agreed with the findings of [15 and16] who reported reduction in anti-nutritional factors when they subjected sorrel seed to toasting. Reduction in the contents of the anti-nutritional factors suggests the ability of toasting as an effective method of detoxification sickle pod seed.

Table 2 Proximate and anti-nutritional composition sickle seed meal (%DM)

Parameters	Raw sickle pod seed meal	Toasted sickle pod meal
Dry matter	94.80	90.80
Crude protein	21.45	30.15
Crude fibre	10.18	17.10
Ether extracts	7.50	8.00
Ash	4.16	11.00
Nitrogen free extracts	46.77	33.25
MEkcal/kg [*]	3071.41	2988.94
Anti-nutritional factors mg/100g		
Saponin	185.00	40.12
Phenols	137.85	46.35
Tannins	191.14	43.45
Oxalates	83.00	21.64

Phytate	240.00	26.97
Cyanides	6.98	0.63

Metabolizable Energy = ME (kcal/kg) = 37 x % CP + 81 x % EE + 35.5 x % NFE.
Calculated according to the formula of [13]

3.1 Growth performance of weaner rabbit fed toasted sickle seed meal (TSSM).

The growth performance parameters of weaner rabbits fed graded levels of toasted sickle seed meal are presented in Table 3. Result showed that total feed intake (TFI) of rabbits on T1 diet was not significantly ($P>0.05$) different from those on T2, T3, T4 and T5 diets. The TFI values ranged from 3201.20g for rabbits on T3 diet to 3235.40g for those on T2 diet. There was no significant ($p>0.05$) difference in the total weight gain (TWG) of rabbits. The TWG varied from 930.00g for rabbits fed T1 and T3 diets to 936.67g for those on T4 diet. However, final weights and Feed conversion ratio (FCR) were not influenced by dietary inclusion of toasted sickle seed meals. Final weights were within the range of 1649.00 – 1650.00g, while FCR ranged of 3.43 in rabbits fed T2 and T4 to 3.47 in rabbits fed T1 diet.

The result is in consonant with that of [17 and 18], who fed roasted pigeon pea meal to broilers and reported no significant difference on the broilers performance. Ref [19] also made similar observation when he fed graded levels of toasted pigeon pea meal to weaner rabbits. However, the result negates the finding of [20 and 21] who reported significant decreased in growth performance of broiler chicken and rabbits fed up to 10% inclusion of cassia *obtusifolia* seed Meal. Variation in the result could be attributed to the differences in species of animal used in the two studies and processing methods. The result of average daily weight gain (ADWG) of weaner rabbits fed graded levels of TSPM was higher the range of 12.27 to 15.70g reported by [19] and that of the range of 13.49–15.44g reported by [22]. The non-significant difference of parameter measured the result obtained for final weight, total weight gain, feed conversion ratio and total feed intake agreed with the findings of [23] who reported rabbits fed diets containing heat processed seed meal were not significantly different from the control diet.

Table 3 Growth performance of weaner rabbit fed toasted sickle seed meal (TSSM)

Parameter	Inclusion levels of toasted sickle seed meal					SEM
	T1	T2	T3	T4	T5	
Initial weight (g)	710.67	708.00	710.00	708.33	709.67	7.09 ^{ns}
Final weight (g)	1646.70	1650.00	1640.00	1645.00	1642.30	12.30 ^{ns}
Total weight gain (g)	930.03	942.00	930.00	936.67	932.63	6.32 ^{ns}
Average daily weight gain(g)	16.60	16.82	16.60	16.72	16.65	8.21 ^{ns}
Total feed intake (g)	3228.10	3235.40	3201.20	3213.70	3219.70	10.12 ^{ns}
Average daily feed intake (g)	57.67	57.78	57.16	57.38	57.49	5.12 ^{ns}
Feed conversion ratio	3.47	3.43	3.44	3.43	3.45	1.32 ^{ns}

NS= not significantly different ($P<0.05$), SEM = Standard error mean

3.2 Carcass characteristics and organ weights of weaner rabbits fed toasted sickle seed meal (TSSM)

The result of carcass characteristics and internal weight organs of weaner rabbits are presented on Table 4. The result showed non- Significant differences ($P > 0.05$) on the live weight, dressed weight and dressing percentage among the dietary treatment groups. The live weight values obtained in this study are within the range 985.08- 1960.08 recorded by [24]. The dressed weight value range of 550 - 820g obtained in the present study are lower than the value 499.0- 592.00g recorded by [3]. The dressing percentage varied from 51.29 – 53.11%

The result of internal organ weights were not significant ($P > 0.05$) influenced by inclusion levels of TSSM in the diets. It is evident therefore, feeding weaner rabbit with vary levels of TSSM did not illicit any toxic response as internal organs such as liver and kidney are the major organs of detoxification did not undergo any hypertrophy [25, 26].

Table 4 Carcass characteristics and organ weights of weaner rabbits fed toasted sickle seed meal (TSSM)

	Inclusion levels of toasted sickle seed meal.					
	T1	T2	T3	T4	T5	SEM
Live weight (g)	1619.38	1621.00	1619.21	1605.00	1603.00	16.13 ^{ns}
Slaughter weight	1600.00	15900.00	1603.30	1525.00	1545.00	18.86 ^{ns}
dressed weight (g)	841.08	860.99	835.34	823.53	812.20	8.36 ^{ns}
Dressing (%)	51.93	53.11	51.58	51.30	51.29	0.51 ^{ns}
Small intestine length (cm)	213.33	212.00	216.67	212.33	210.00	2.12 ^{ns}
Large intestine length (cm)	101.67	106.70	103.87	106.88	106.77	4.40 ^{ns}
Internal organs (% live weight)						
Small intestine weight (g)	3.25	3.24	3.25	3.38	3.92	0.03 ^{ns}
L intestine weight (g)	1.09	1.22	1.46	1.63	1.67	0.22 ^{ns}
Caecal	6.62	6.15	6.12	6.21	6.53	0.06 ^{ns}
Liver	2.81	2.50	2.75	2.78	2.83	0.03 ^{ns}
Heart	0.20	0.19	0.20	0.20	0.22	0.02 ^{ns}
Lungs	0.77	0.76	0.77	0.81	0.90	0.01 ^{ns}
Kidney	0.56	0.61	0.68	0.72	0.79	0.40 ^{ns}
Stomach	3.32	3.03	3.32	3.59	3.57	0.18 ^{ns}

NS= not significantly different (P<0.05), SEM = Standard error mean

4.0 Conclusion

From the results of this study it can be concluded that inclusion of toasted sickle pod seed meal (TSSM) up to 10% in the diet of weaner rabbits did not affect growth performance and carcass characteristics. Therefore, TSSM can be included up to 10% in rabbits' diets

References

- [1] Yashim, S.M., Anene, D., Gadzama, I.U. and Makama, S. R (2016). Effect of Different Processing Methods of *Crotalaria retusa* Leaves on Growth Performance of Weaner Rabbits. Proc. 21st Ann. Con. Animal Science Association of Nigeria 18 – 22, 2016/ Port Harcourt .814 -814pp
- [2] Igwebuike, J. U., Medugu, C. I., Kwari, I. D. and Dauda, A. (2013). Growth Performance and Nutrient Digestibility of Growing Rabbits fed two varieties of Sorghum as replacement for Maize as energy source in Tropical Environment. Global Journal of Bio-Science and Biotechnology, 2 (2): 167 – 172.
- [3] Wafar, R.J and Tarimbuka, L.I (2016) effects of substituting groundnut cake with supported by water spinach (*ipomoea aquatica*) leaf meal on performance, carcass yield and blood profile of weaner rabbit. FUW Trends in Science & Technology Journal Vol. 1 No. 1 pp 238-242
- [4] Ani, A. O. and Ugwuowo, L. C. (2011). Response of weaner rabbits to diets containing graded levels of processed velvet beans (*Mucunapruriens*). African Journal of Biotechnology Vol. 10(66), pp. 14984-14989
- [5] Lar J, Gupta PC. Anthraquinone glycosides from the seeds of *Cassia occidentalis* Linn. *Experientia*. (1973);29:142–3.
- [6] Ibrahim, M.A; Aliyu, A.B and T. S. Umar *Senna occidentalis* leaf extract possesses antitrypanosomal activity and ameliorates the trypanosome-induced anemia and organ damage. *Pharmacology Research* 2(30). 175- 180
- [7] Ismaila, Y. S., Denban, M. K., Emmanuel, K.M and Augustine, C., (2011), Nutritional and phytochemical screening of *Senna obtusifolia* Indigenous to Mubi, Nigeria. *Advances in Applied Science Research*, p 2011, 2(3): 431-437
- [8] Ingweye, J.N. G.A. Kalio, J.A Ubua and E.P. Umoren, (2010). Nutritional evaluation of wild sicklepod (*Senna obtusifolia*) seeds from Obanlikusouth Nigeria. *Am. J. Food Technol.*, 5: 1-12.
- [9] Adebayo, A. A. (1999). Climate II. In: Adamawa State in Maps. Paraclete publishers, Yola, Nigeria Nigeria Pp.112
- [10] AOAC, 2000. Association of Analytical Chemist. Official Methods of Analysis 17th Edition AOAC Inc, Arlington, Virginia, USA.
- [11] Steel RGD, Torrie JH (1980). Principles and Procedures of Statistics. A Biometric approach 2nd ed. McGraw-Hill Publishers, New York.
- [12] Duncan DB (1955). New Multiple Range and Multiple F. Tests. *Biometrics*, 11: 1-42.
- [13] Pauzenga, U. (1985). Feeding parent stock. Zootech characteristics of weanling rabbits fed graded levels *International*, 34: 22-25
- [14] Essiett, U. A and Bassey, I. E., (2013), Comparative Phytochemical Screening and Nutritional Potentials of the Flowers (petals) of *Senna alata* (L) roxb, *Senna hirsuta* (L.) Irwin and barneby, and

- Sennaobtusifolia (L.) Irwin and Barneby (Fabaceae). Journal of Applied Pharmaceutical Science Vol. 3 (08), pp. 097-101
- [15] Bressani, R. Elias, L.G. and Braham, J.E. (1982) Reduction of digestibility of legume proteins tannins. J. Plants. Foods: 4: 43-55.
- [16] John, A. B., Yahaya M. S., Julius, N. (2010). Effect of Sorrel Seed (Hibiscus Sabdariffa) Supplementation on the Performance of Yankasa Rams Fed Brachiaria Decumbens as Basal Diet. J. Agri. Vet. Sciences., 2: 8-16.
- [17] Amaefule, K. U., & Nwaokoro, C. C. (2002). The effect of graded levels of raw pigeon pea (Cajanus cajan) seed meal on the performance of weaner rabbits. In V. A. Aletor & G. E. Onibi, (Eds.), Increasing household protein consumption through improved livestock production (pp. 113-115). Proceedings of the 27th Annual Conference of Nigerian Society for Animal Production (NSAP), March 17-21, Akure.
- [18] Ani, A. O., & Okeke, G. C. (2003). The substitution of pigeon pea (Cajanus cajan) seed meal for soyabean in broiler finisher ration. In E. A. Olatunji, B. A. Ayanwale, E. L. Shiwoya, & A. Aremu (Eds.), Sustainable livestock productivity and national development: The holistic approach (pp. 10-12). Proceedings of the 8th Annual Conference of Animal Science Association of Nigeria (ASAN), September 16-18, Federal University of Technology Minna, Niger State, Nigeria.
- [19] Kemi E. Akande (2015) Dietary Effects of Increasing Levels of Pigeon Pea Meal on Rabbit Performance. Journal of Agricultural Science; Vol. 7, No. 7; 156-162 Pp
- [20] Hebert, C.D. and W. Flory, 2002. Plant food for human nutrition Springer Netherlands, pp: 57.
- [21] Augustine, C.; Igwebuike, J.; Salome, S. Midau, Ja'afar-Furo, A. M.R. Mojaba, D.I. and Dazala, I.U. (2010) Evaluation of Economic Performance of Broiler Chickens Fed Graded Levels of Processed Cassia obtusifolia Seed Meal. International Journal of Sustainable Agriculture 2 (3): 47-50,
- [22] Iorgyer, M. I., Carew, S. N., & Ayoade, J. A. (2008). The replacement value of pigeon pea (Cajanus cajan) for maize in weaner rabbit diets. Repositioning Animal Agriculture for the Realization of National Vision 2020 (pp. 456-459). Proceedings of the 13th Annual Conference of the Animal Science Association of Nigeria, September 15-19, Zaria, Nigeria.
- [23] Ihekweumere, F. C., Onyekwere, M. U., & Egu, U. E. (2008). Growth, carcass and reproductive characteristics of male rabbits (bucks) fed raw and boiled pigeon pea seed (Cajanus cajan) meal. Pakistan Journal of Nutrition, 7, 17-20. <http://dx.doi.org/10.3923/pjn.2008.17.20>
- [24] Omoikhoje S.O, Obasoyo, D.O and Obazee, S (2016) Growth performance of weaner rabbits fed graded levels of bread waste meal supplemented with guinea grass (Panicum maximum) pp836-840
- [25] Bone, F.J. (1979). Anatomy and Physiology of Farm Animals. 2nd Edn., Reston Publishing Comp, Inc Virginia, USA, pp: 560.
- [26] Carew, L. B., Hardy, D., Gernat, A.G. and Zakrzewska, E. I. (2003). Heating raw velvet beans (Mucuna pruriens) reverses some antinutritional effects on organ growth, blood chemistry and organ histology in growing chickens. Journal Tropical and Subtropical Agroecosystems 1 (2-3): 267 – 275