

Nutritonal properties of composite flour (blends of rice (*Oryza sativa*), acha (*Digitaria exilis*) and soybeans (*Glycine max*) and sensory properties of noodles produced from the flour

ABSTRACT

This study was carried out to evaluate the nutritional properties of composite flour made from rice, acha, soy beans and wheat, and to assess the sensory acceptability of the noodles made from the composite flour. Different ratios of rice, acha, soybeans and wheat flour blends 100W (wheat) which is the control sample; 75: 15:10, (wheat, rice and soybean) sample, 70:20:10 (wheat, rice and soybean) sample, 65:20:15 (wheat, rice and soybean) sample, 60:25:15 (wheat, rice and soybean) sample, 50:30:20 (wheat, rice and soybean) sample, 40:40:20 (wheat, rice and soybean) sample; 80:15:5 (wheat, acha, soybean) sample, 70:20:10 (wheat, acha, soybean) sample, 65: 20: 15 (wheat, acha, soybean) sample, 60:25:15 (wheat, acha, soybean) samples, 50: 30:20 (wheat, acha, soybean) sample, 40:40:20 (wheat, acha, soybean) sample. Proximate composition data for protein, fat, crude fibre, ash, carbohydrate and energy ranged from (5.26-18.82%, 7.02- 21.01%, 0.33- 0.79%, 0.40- 1.21%, 57.63- 74.94%, 1568.91- 1945.29%, respectively. There was a significant difference ($P<0.05$) in the mineral content of samples. Samples with rice blends had the highest mineral content and blends with acha compared favorably with the control. The result of this study shows that vitamin A and vitamin C were significantly different ($P<0.05$) in the sample. The result of the sensory evaluation of the noodles prepared from the composites shows that most of the noodles were generally accepted. The samples that had rice in their blends were more accepted than the samples that had acha and they all compared favorably with the control. The results shows that supplementing wheat with cereals (rice, acha) and legumes (soybean) to producing a good quality composite flour and noodles is achievable and this will increase the health benefits of noodles and promote the consumption and utilization of our local foods that will improve nutrients in the meals of children.

Keywords: Noodles, composite flour, Acha, nutritional, sensory

1. INTRODUCTION

Noodles have become one of the most staple and popular foods in many parts of the world including Nigeria. Available in an amazing range of shapes and flavours, it is incredibly versatile, and can be served in scores of different ways. Modern instant noodle was invented in Osaka Japan in 1958 by Taiwanese/Japanese businessman, Momofuku Ando, the founder of Nissin foods, one of the biggest manufacturers of noodles in the world (Miskelly, 1993). Approximately 85 billion servings of instant noodles are eaten worldwide every year (BBC News, 2005). Of this number China alone consumes 44 billions packs, being the largest producer and consumer of noodles (WINA, 2008). Per capita noodle consumption in Nigeria was estimated at 650 million packs in 2005, 700 million packs in 2006 and one billion packs in 2007 (WINA, 2008). Hence noodles and pasta market in Nigeria increased between 2000 - 2008 at an annual rate of 3.6% (WINA, 2008).

Rice is the most common and important cereal for human consumption and calorie intake. It is the staple food for over 3 billion people, constituting over half of the world's population providing more than 5th of the

calories consumed worldwide (Cantral and Reeves, 2002). Rice is grown in all the ecological and dietary zones of Nigeria, with different varieties processing adaptation traits for each ecology (Sanni *et al.*, 2005). Rice is an economic crop, which is important in household food security, ceremonies, nutritional diversification, income generation and employment (Perez *et al.*, 1987). It is utilized mostly at the household level, where it is consumed as boiled or fried or ground rice with stew or soup. Rice is cooked by washing and boiling in water which leads to loss of some nutrients (Ihekeronye and Ngoddy, 1985; Perez *et al.*, 1987).

Acha (*Digitariaexillis*) is a cereal grain in the family of *gramineae* and commonly referred to as *folio* or hungry rice (Ayo *et al.*, 2014). Nigerian people of Plateau, Bauchi and Kaduna states burn straw and ash extracted with water through perforated basket. The filtrates are used as potash for cooking indigenous delicacies. Acha grain can also be grounded into flour to produce biscuit (Obafunmi *et al.*, 2001). Acha are also classified on the color and sizes of the grain. Acha is one of the most nutritious of all grains. Its seed is rich in methionine and cystine, they are vital to human health and these amino acids are deficient to major cereals. Like wheat, rice, maize, sorghum, barley and rye (Jideani and Akingbala, 1993).

Soybean is one of the protein rich legumes consumed around the world and used as enrichment in many food formulations especially in Africa. Soybean contains about 40% protein; it is higher than other legumes in protein. The protein of soybean yields all the essential amino acids in adequate amount except methionine and cystine which are deficient (Srilakshmi, 2009). Soybean is a rich source of vitamins like thiamine, riboflavin, niacin, folic acid. It is also a rich source of iron. It contains high amount of phosphorus, magnesium and potassium, and fair source of carotene and vitamin D. It is a poor source of calcium (Iwe, 2003). Soybean is nutritionally valued for their unsaturated fatty acids, protein, and fibre content. The oil contains about 85% unsaturated fatty acid and 15% saturated fatty acids. It contains 30-32% carbohydrate (soluble and insoluble). The soluble carbohydrate includes sucrose and oligosaccharide, starchyose and raffinose. Soybean is rich in lecithin and linoleic acid (Srilakshmi, 2009).

Despite the wide consumption of noodles in Nigeria the products are regarded as unhealthy (junk) food (Gunathilake and Abeyrathne, 2008). This is orchestrated by the fact that the conventional instant noodles are high in carbohydrate (<70%) and low in protein, fibre, vitamins and minerals.

There is also the problem of gluten allergenicity where some individuals are intolerant to glutens of wheat and the related cereals oats, rye, and barley. This intolerance, celiac disease, seriously impairs intestinal absorption and can lead to severe malnutrition (Ciclitira and Ellis, 1987; Davison and Bridges, 1987). To worsen the case instant noodles are typically fried as part of the manufacturing process resulting in high levels of saturated fat and trans-fat. Conventionally, noodles are made from the dough of unleavened durum wheat flour (Hoseney, 1986) which unfortunately is not a good source of fibre and protein. The unique ability of wheat flour to form cohesive, elastic extensive dough as a result of its gluten protein makes it an ideal material for noodles production and baked product such as breads, cakes, biscuits and cookies. In bakery gluten forms, as glutenin molecules, cross-link to make a sub-microscopic network and associates with gliadin, which contributes with viscosity and extensibility to the mix. Gluten content was implicated as a stalling factor of bread, because it binds water by hydration. The development of gluten affects the texture of the baked goods (Alvarenga *et al.*, 2011).

Enrichment of cereal based food with other protein source such as legumes has received considerable attention. The use of cereal-legume based food has long been advocated as alternative protein and energy source for infant and young children food products (Mensah and Tunkins, 2003). It is evident that when cereals and legumes are judiciously selected and combined, desirable pattern of essential amino acids of high biological value is obtained (Nnam, 2001).

This study is one of the efforts to promote the use of composite flours in which flour from locally grown crops and soybean with high protein content shall be used to produce protein-enriched composite flour. Thus, the aim of this work is to produce and evaluate the nutritional, optimum proportion of rice flour, acha flour and soybean flour for production of composite flour and sensory evaluation of noodles produced compared with wheat flour (as control sample).

2. MATERIAL AND METHODS

2.1 Sample collection

The raw materials for this study were rice (*Oryza sativa*), acha (*Digiteria exillis*) and soybean (*Glycine max*). The raw materials were purchased from Ubani market in Umuahia, Abia state.

2.2 Sample Preparation

The rice grains were sorted and washed five times with water, sun dried, and then dried in the oven for 12 h at 60^o C. It was then milled and sieved with 0.05 mm screen and then packaged in a plastic container.

Acha grains were sorted and washed five times with water. After washing, they were sun dried for 12 h, oven dried for 30 min at 65^oC and milled with hammer mill to a fine texture. The flour was sieved with 0.05 mm screen to obtain fine flour and stored in an air tight container.

The soybean seeds were sorted and soaked for 12 h, then dehulled. The seeds were boiled for 30 min at 100^oC, drained and oven dried for 12 h at 65^oC, milled and sieved with 0.05mm screen to fine flour.

2.3 Composite flour formulations

The flours were mixed in the following ratios 75:15:10, 70:20:10, 65:20:15, 60:25:15, 50:30:20, 40:40:20 for wheat, rice and soybean while, wheat, acha and soybean were blended in the following ratios 80:15:5, 70:20:10, 65:20:15, 60:25:15, 50:30:20, 40:40:20 and 100 % wheat flour served as the control.

2.4 Noodle production

Noodles were produced by blending separately the flour samples with warm water (40^oC) and 5% CMC according to the method described by Omeire *et al.*, (2014). The mixtures were thoroughly worked to form dough. The formed dough was allowed to rest for 20 min, then kneaded and rolled with a rolling pin to form sheets. The sheets were extruded using a cold extruder (Eurosonic, Globe 150) model. The noodle strands were put in cleaned aluminium trays and oven dried at 60^oC. The process is as shown in Figure 1.




Figure 1: Flow diagram showing the production of Noodles

2.5 Chemical analysis

The proximate compositions of the samples were determined using Association of Official Analytical Methods (2000). The samples were analyzed for moisture content, ash, crude protein, lipids and crude fibre. Carbohydrates content was determined by difference and energy value was calculated by method

describe by Onwuka (2005). The minerals (sodium, calcium, magnesium and phosphorus) contents were determined using standard method according to the Official Method of Analysis (1995). The vitamins compositions of the samples were determined using Barakat titration method for Ascorbic acid and vitamin A according to the Official Method of Analysis (2000).

2.6 Sensory evaluation

Five variables were used for this study to test for acceptability of the samples. The variables were taste, appearance, texture, aroma and general acceptability. The panel comprised of a 20 man semi-trained judges who were asked to rank the samples using 9 point Hedonic scale where, 1 represents dislike extremely and 9 like extremely.

2.7 Statistical analysis

Data obtained from nutritional and sensory evaluation of the noodles were recorded in triplicates and subjected to One-way analysis of variance (ANOVA) using Statistical Program for Social Science (SPSS) version 21.0 (IBM SPSS inc., Chicago, IL). Where difference existed, LSD was used to separate means and accepted at 5% significance level (Adeboye *et al.*, 2015).

3. RESULTS AND DISCUSSION

3.1 Proximate composition of the flour samples

Table 1 shows the results of the proximate composition of the composite flour samples and the control. There was significant difference ($P>0.05$) observed between the samples and the control for moisture, protein, fat, crude ash, carbohydrate and energy. In terms of moisture content the composite flour samples and the control values ranges from 8.32 to 10.82%. The moisture content values of the samples and the control were within the recommended moisture content of 15% acceptability value for long term storage of grains and acceptable limit for dry product (Okaka, 2005). The relatively low moisture content of this flour samples is important because, it will help in extending the shelf life if adequately protected, by inhibiting the development of contaminating microorganism whose growth and activities are favoured by presence of moisture (Okafor and Ugwu, 2014).

It is believed that materials such as flour and starch containing more than 12% moisture have less storage stability than those with lower moisture content, for this reason water content of 10% is generally specified for flour and related products (Opeifa *et al.*, 2015). According to Haard and Chism (1996), high

162 quality noodles should have an adequate shelf life without any microbiological deterioration and therefore
163 the low moisture content of these flour samples will be seen to extend the shelf life of the final product.

164 **Table 1: Proximate composition of rice, acha, soybean and wheat composite flour at different**
165 **proportions**

Samples	Moisture %	Protein %	Fat %	Crude fibre %	Ash %	Carbohydrate %	Energy %
75W 15R 10S	9.96 ^e ±0.12	9.63 ^f ±0.00	14.51 ^e ±0.12	0.42 ^{cd} ±0.06	0.61 ^d ±0.12	65.19 ^f ±0.21	1808.91 ^f ± 0.18
70W 20R 10S	10.32 ^b ±0.12	6.13 ^j ±0.15	19.52 ^b ±0.15	0.40 ^c ±0.06	0.80 ^c ±0.00	63.03 ^h ±0.06	1897.84 ^b ± 0.47
65W 20R 15S	9.33 ^e ±0.12	9.17 ^g ±0.20	21.01 ^a ±0.17	0.39 ^{de} ±0.06	0.80 ^c ±0.00	59.59 ⁱ ±0.06	1946.29 ^a ± 0.50
60W 25R 15S	9.06 ^g ±0.00	7.87 ⁱ ±0.12	7.02 ^k ±0.15	0.37 ^{ef} ±0.06	1.00 ^b ±0.00	74.94 ^a ±0.06	1667.28 ⁱ ± 0.31
50W 30R 20S	8.02 ^j ±0.12	13.57 ^c ±0.12	16.51 ^c ±0.10	0.35 ^{fg} ±0.12	1.00 ^b ±0.00	60.69 ^j ±0.38	1873.27 ^c ± 0.19
40W 40R 20S	8.38 ^k ±0.12	14.02 ^b ±0.12	11.01 ^h ±0.17	0.33 ^h ±0.06	1.21 ^a ±0.12	65.82 ^e ±8.7	1755.93 ^j ± 0.40
80W 15A 5S	10.82 ^a ±0.12	5.26 ^m ±0.12	9.49 ⁱ ±0.12	0.46 ^b ±0.06	0.40 ^e ±0.00	74.11 ^b ±0.20	1698.21 ^k ± 0.94
70W 20A 10S	9.41 ^d ±0.23	7.44 ^j ±0.00	12.02 ^g ±0.20	0.39 ^{de} ±0.12	0.40 ^e ±0.00	70.65 ^c ±0.31	1772.21 ^g ± 0.56
65W 20A 15S	8.47 ^j ±0.10	18.82 ^a ±0.20	14.01 ^f ±0.17	0.38 ^{de} ±0.23	1.00 ^b ±0.00	57.63 ^k ±0.15	1818.20 ^e ± 0.51
60W 25A 15S	9.23 ^f ±0.15	13.12 ^d ±0.12	12.01 ^g ±0.12	0.38 ^{de} ±0.12	1.00 ^b ±0.00	64.51 ^g ±0.35	1764.20 ^h ± 0.51
50W 30A 20S	8.64 ⁱ ±0.00	6.57 ^k ±0.12	9.52 ⁱ ±0.15	0.35 ^{fg} ±0.06	0.81 ^c ±0.12	74.41 ^b ±0.15	1728.61 ^j ± 0.14
40W 40A 20S	8.68 ^h ±0.21	8.76 ^h ±0.00	16.02 ^d ±0.15	0.39 ^{de} ±0.12	1.21 ^a ±0.12	65.16 ^f ±0.15	1849.20 ^d ± 0.66
100W (control)	10.82 ^a ±0.12	10.51 ^e ±0.06	8.51 ^j ±0.12	0.79 ^a ±0.12	1.00 ^b ±0.00	69.06 ^d ±0.21	1667.55 ⁱ ± 0.04
LSD	0.25	0.02	0.03	0.02	0.01	0.45	2.05

166 Values are means of data of triplicate determinations. Values in the same column having the same
167 superscript are not significantly difference (P>0.05). W- Wheat, R- Rice, A –Acha, S- Soybean

168
169 The crude protein content of the formulated samples and the control ranges from 5.26 to 18.82%. In this
170 case, there was significant different (P<0.05) between the samples and control. The protein content of the
171 control 100W reported in this study was found to be lower than the 14.70% (Adebawale *et al.*, 2005) and
172 12.86% (Morris *et al.*, 1999) for wheat flour but close to the values of 10.23% reported by Oppong *et al.*,
173 (2015). This could be attributed to the geographical location, since soil with high nitrogen levels can
174 influence protein level in the grain (Brown, 1991). Wheat variety play more important role in determining
175 the protein content of wheat. The variety of wheat used was the soft red winter grown in countries like
176 America, Europe and Australia. The protein content of this variety ranges from 8.00-11.0% (Okaka,
177 2005).

Sample 65W20A15S had the highest crude protein value, while the rest of the samples had reasonable amount of crude protein in them and they compared favorably with the control but sample 80W15A5S had the least value of 5.26%. The protein content of these flour blends with rice, acha and wheat in this study suggests that they may be useful in food formulation systems (Oppong *et al.*, 2015). The protein content of most of the samples and the control were above the standard value of 8-12% and may definitely contribute to the nutritional needs of children and adults (Srilakshmi, 2008).

The higher protein values of these wheat, rice, soybean and wheat, acha, soybean composites will be of nutritional importance in most developing countries like Nigeria where many people can hardly afford high proteinous food because of cost (Falola *et al.*, 2013). With these results high protein content of ready to eat noodles will be achieved. The presence of soybean in the formulation also was contributed to the high crude protein recorded in this study.

The fat content of the flour samples and the control 100W ranges from 7.02 to 21.01%. blend sample 65W 20R 15S had the highest value while the least value was recorded by sample 60W 25R 15S .The high fat content observed in this sample blends will be attributed to the presence of soybean as reported by Iwe (2000), that malted soybean have high fat content. Since all the samples had soybean so their fat content were high. The results obtained in this study were similar to the 12.02-18.34% fat content that was obtained by Ayo *et al.*, (2014) for the production and evaluation of malted soybean – acha composite flour bread and biscuit.

The increase in the fat content could be a good source of energy to the human body. Fats and oils provide more than twice as much energy as the carbohydrate on a weight-weight basis (Iwe, 2000). One gram of fat or oil will yield about 368k/cal of energy when oxidized in the body. High fat content of the product may not provide a conducive environment for microbial growth and activities hence improving the shelf life of the product (Okaka, 2005). High lipid content can reduce clarity of starch paste and hinder the swelling of starch granules because of high rate formation of amylose lipid complex (Chez, 2003). Nevertheless the protein and lipid played important role in retention of amylose in starch noodles during cooking in minimizing cooking loss (Kim *et al.*, 1996).

The crude fibre values of the samples and the control ranges from 0.33 to 0.79%. The control 100W had the highest fibre content with the value of 0.79%, this value was close to the 0.85% value of wheat

reported by Ayo et al., (2007). The high crude fibre is most likely from the bran of the whole wheat and the hull of soybean represents variable fraction of dietary fibre which include mostly the lignin, cellulose and hemicellulose (Manny and Shadaksharaswany, 2005).

The rest of the samples also had relatively high fibre content compared to the control this could be attributed to the presence of wheat and soybean due to the fact that malted soybean are relatively high in crude fibre (Iwe, 2000), which are important in bowel movement and reduces constipation. The crude fibre content obtained in this study is lower than the range of 0.47 – 1.04% obtained for malted soybean-acha composite flour bread and biscuit by Ayo *et al*, (2014). Dietary fibre results in reduction of the risk of bowel disorders and fight constipation (Osaretin *et al.*, 2007). It is expected that the fibre content of the samples should be higher, since acha and wheat grains are said to have high fiber content. The results of the samples were generally low. This could be attributed to the separation processes carried out from milling to obtaining fine flour used for the blend with wheat and acha.

The ash content of the samples and the control ranges from 0.40 to 1.21%. Samples 40W40R20S and 40W40A20S had the highest ash content value of 1.21%. The samples with the highest portion of soybean were observed to have the highest ash content value. The increase could be attributed to the fact that soybean are high in mineral (Omeire *et al.*, 2014). Ash is a non organic compound containing mineral content of food and nutritionally it aids in the metabolism of other compounds. The high ash content of the samples could be a source of minerals which apart from its nutritional value are good for good skin and strong bones (Ayo *et al.*, 2014). The ash content of the samples was averagely high which indicates that they have good mineral content. The ash content represents the total mineral content in foods and serves as available tool for nutritional evaluation (Lienel, 2002).

The carbohydrate content of the samples and the control varied from 57.63 to 74.94%. Sample 60W25R15S had the highest value of carbohydrate content with value of 74.94% and the least value is sample 50W30R20S with 57.63% value. The differences in the carbohydrate contents of the samples and control could be attributed to the differences in their protein and fat content values of the samples.

The high carbohydrate content of these samples suggests that the flours would be a very cheap source of energy for the body (Omeire *et al.*, 2015). The high carbohydrate content of the samples is attributed to the high carbohydrate content in the cereals that are the principle ingredients in the samples. A marginal

difference was observed when the results are compared with the values reported by Kanu *et al.*, (2009) for breakfast cereal based porridge.

The energy content of the samples and control ranges from 1568.91% – 1945.29%. Sample 65W20R15S had the highest energy value content of 1945.29KJ and the least value was recorded by sample 75W15R10S. This result could be attributed to the values of protein and fat content of the samples. This is highly desired especially in famine and war locations where the next meal is not easy to come by. High energy foods have shown to have protective effect in the optimal utilization of other nutrients (Wardlaw, 2004). Energy was observed to be high for all the samples, so energy content is a parameter used to determine the quality of food especially for formulation designed for adults with high energy requirements.

3.2 Mineral and Vitamin compositions of the flour samples.

The result of mineral (sodium, calcium, phosphorus and magnesium) and Vitamin (vitamin A and vitamin C) values of rice, acha, soybeans and wheat composite flours at different proportions is shown in Table 2.

There was significant difference ($P<0.05$) that existed among the composite flour samples and the control for the minerals and vitamins analyzed. In terms of sodium (Na) content samples 50W30R20S and 40W40R20S had the highest Na values of 4.02 and 4.06mg/100g respectively and the control 100W had the least value of 3.05mg/100g. This result could be attributed to the fact that all the other samples had soybean substitution. Soybean seeds have been reported to contain appreciable amount of mineral and fat (Plahar *et al.*, 2003). Sodium intake needs to be monitored as it can become a major dietary problem where high blood pressure problems are concerned. The low level of sodium observed in this study will make the product suitable for use in sodium restricted diets (Iyang and Ekop, 2015).

In terms of calcium content sample 50W30R20S had the highest Calcium value of 45.07mg/100g and sample 80W15A5S had the least value of 29.63mg/100g, this could be attributed to the fact that it had the least amount of soybean. All the samples compared favorably with the control (100W) which had a value of 35.13mg/100g.

Calcium is by far the most important mineral that the body requires and its deficiency is more prevalent than many other minerals, it helps in the formation of strong bone and teeth (Ishitani *et al.*, 1999). High calcium of flour samples and control shows that products from them would be nutritionally beneficial for

both children and elderly people who require high calcium intake for strong bone and body development (Bolarinwa *et al.*, 2015).

Table 2: Mineral and Vitamin compositions of rice, acha, soybean and wheat flour at different proportions

Samples	Sodium mg/100g	Calcium mg/100g	Magnesium mg/100g	Phosphorus mg/100g	Vitamin A IU	Vitamin C mg
75W 15R 10S	3.37 ^d ±0.81	39.20 ^c ±6.95	128.60 ^d ±1.21	363.33 ^a ±1.15	1505.33 ^c ± 1.16	6.33 ^d ±0.06
70W 20R 10S	3.15 ^f ±0.12	38.00 ^d ±5.12	117.40 ^f ±5.1962	311.00 ^h ±1.73	1508.33 ^b ± 1.16	6.36 ^d ±0.06
65W 20R 15S	3.51 ^c ±0.12	39.13 ^c ±0.58	131.57 ^c ±6.33	337.33 ^d ±1.15	1508.33 ^b ± 0.58	6.57 ^{bc} ±0.06
60W 25R 15S	3.52 ^c ±0.12	39.23 ^c ±0.58	131.60 ^c ±1.21	334.33 ^e ±5.77	1505.33 ^c ± 0.58	6.46 ^{bcd} ±0.06
50W 30R 20S	4.05 ^a ±0.06	45.07 ^a ±0.58	136.17 ^a ±0.58	342.33 ^c ±5.77	1506.33 ^c ± 0.58	6.41 ^{cd} ±0.06
40W 40R 20S	4.02 ^a ±0.06	42.68 ^b ±0.58	134.10 ^b ±1.73	336.67 ^d ±5.77	1600.67 ^a ± 0.58	6.59 ^b ±0.06
80W 15A 5S	3.64 ^b ±0.23	29.63 ⁱ ±0.58	115.90 ^g ±5.20	325.67 ^f ±1.15	1350.00 ^e ± 0.00	5.67 ^e ±0.06
70W 20A 10S	3.46 ^c ±0.06	32.47 ^f ±1.16	110.80 ⁱ ±1.21	315.33 ^g ±1.15	1300.33 ^h ± 0.58	5.13 ^f ±0.06
65W 20A 15S	3.26 ^e ±0.06	30.37 ^h ±0.58	114.47 ^{gh} ±4.62	326.33 ⁱ ±5.77	1321.67 ^g ± 0.58	5.07 ^f ±0.06
60W 25A 15S	3.31 ^e ±0.06	32.47 ^f ±1.16	116.40 ⁱ ±6.93	297.67 ^j ±5.77	1330.33 ⁱ ± 0.58	4.19 ^g ±0.06
50W 30A 20S	3.28 ^e ±0.12	31.33 ^g ±1.16	113.53 ^h ±5.78	302.67 ⁱ ±5.77	1281.33 ^j ± 1.16	5.06 ^f ±0.06
40W 40A 20S	3.13 ^f ±0.17	31.27 ^g ±0.58	115.83 ^g ±6.35	317.33 ^g ±5.77	1170.33 ^j ± 0.58	5.13 ^f ±0.06
100W (control)	3.05 ^g ±0.01	35.13 ^e ±0.58	123.33 ^e ±5.76	348.67 ^b ±1.15	1400.00 ^d ± 0.00	7.03 ^a ±2.31
LSD	0.05	0.45	1.32	1.71	0.30	0.01

Values are means of data of triplicate determinations. Values in the same column having the same superscript are not significantly difference (P>0.05). W- Wheat, R- Rice, A –Acha, S- Soybean

In terms of magnesium content, the magnesium content values of the flour samples and the control ranges from 110.80 – 136.17mg/100g, where sample 50W30R20S had the highest value and sample 70W20A10S had the least value. The blends with acha had slightly lower values than the blends with rice but all the samples compared favorably with the control (100W).

For phosphorus content, the phosphorus content of the samples and the control varied from 297.67 to 363.33 mg/100, where sample 75W15R10S had the highest value while the least value was recorded by sample 60W25A15S.

The samples with acha had slightly lower values of phosphorus than the samples with rice but all samples still compared favorably with the control (100W). The phosphorus content of the samples and control obtained in this study were higher than the phosphorus content 183.10mg/100g of wheat soy flour reported by Anna *et al* (2011). The relatively high phosphorus content of the samples and the control is an indication that the flour products will help in the formation of teeth and bones in children and in their proper development (Bolarinwa *et al.*, 2015). The result from this study shows that the noodles would contribute substantially to the recommended dietary requirement for mineral (Wardlaw, 2004).

Vitamin A is very essential for growth, reproduction, good vision, healthy skin, hair and nail and to balance energy level in the human body. The deficiency of vitamin A in the body causes keratomalacia (night blindness) (Ojimelukwe *et al.*, 2005). The vitamin A content in this study ranges from 1176.33 to 1600.67IU, where sample 40W40R20S had the highest value while sample 40W40A20S had the least value. The samples with acha had values lower than the control but they still compared favorably with control. Soybean is a rich source of vitamins like thiamine, riboflavin, niacin, folic acid. It is also a rich source of iron. It contains high phosphorus, magnesium and potassium and fair sources of carotene and vitamin D and is a poor source of calcium (Iwe, 2003).

In terms of vitamin C content, the control 100W had the highest vitamin C value of 7.03mg and the least value of 4.19mg was recorded by sample 60W25A15S. Vitamin C is required by the body for maintenance of health, gum, healing of wounds, mopping excess of oxygen from the system and is a powerful antioxidant. Deficiency of vitamin C in the body will cause sore gum and scurvy (Ojimelukwe *et al.*, 2005).

3.3 Sensory evaluation of noodles

The result of the sensory evaluation of the formulated samples and the control (100W) into noodles are represented in Table 3.

In terms of taste, there was no significant difference ($P>0.05$) between the samples and the control except for samples 65W20A15S, 60W25A15S and 40W40A20S. The control (100W) had the highest value of 6.77 but the rest of the samples compared favorably with it except for the samples that were significantly different from the control. Taste is an important parameter when evaluating sensory attribute of food. The product might be appealing and having high energy density without good taste, such a

product is likely to be unacceptable (Muhimbula *et al.*, 2011). The samples with rice flour were more liked than the samples with acha flour, it was observed that as the acha flour increased in the mix the less its taste was liked.

In the case of texture, there was no significant difference ($P>0.05$) between the noodles formed from samples and the control except for samples 60W25A15S, 50W30A20S, 40W40W10S, that there was significant difference ($P<0.05$). The panelist preferred the texture of the noodles made from samples with rice flour to the control and the blends with the acha flour was slightly lower to the control, though the noodles of the samples still compared favorably with the control. Texture of staple food 'tuwo' is an important quality attribute that usually influence consumers acceptability of the product (Aboubacar *et al.*, 1999). It is the prevailing textural characteristics of the product at the point of consumption, that usually determine whether such food is swallowable or chewable (Szezesniak, 2002). Sanni *et al.*, (2007) and Omeire *et al.*, (2014) reported similar results for noodles produced from cassava/wheat flour and from wheat, acha, soybean respectively.

In terms of aroma, there was no significant difference ($P>0.05$) between the samples and the control except sample 40W40A20S. The control (100W) and sample 70W20R10S had the highest values of 6.53 and the rest of the noodles made from the other samples compared favorably with them. Smell is an integral part of taste and general acceptance of the food before it is put in the mouth. It is therefore an important parameter when testing acceptability of formulated food (Muhimbula *et al.*, 2011). From this study the noodles produced from rice samples performed slightly better than the samples with acha.

Table 3: Sensory evaluations of noodles from rice, acha, and soybean flour at different proportions

Samples	Taste	Texture	Aroma	Appearance	General acceptability
75W 15R 10S	6.57 ^{ab} ±1.63	6.33 ^{ab} ±1.67	6.07 ^{ab} ±1.87	6.70 ^a ±1.82	6.80 ^a ±1.81
70W 20R 10S	6.23 ^{abc} ±2.01	6.43 ^{ab} ±1.94	6.53 ^a ±1.89	6.47 ^{ab} ±1.68	6.50 ^a ±2.00
65W 20R 15S	6.10 ^{abc} ±2.07	5.87 ^{abc} ±1.96	5.73 ^{ab} ±1.86	6.37 ^{ab} ±1.94	6.43 ^a ±1.36
60W 25R 15S	6.77 ^a ±1.57	6.57 ^a ±1.50	6.13 ^{ab} ±1.85	6.17 ^{ab} ±2.07	6.53 ^a ±1.91
50W 30R 20S	6.37 ^{ab} ±1.61	6.17 ^{abc} ±1.50	6.30 ^{ab} ±1.86	6.53 ^a ±1.53	6.47 ^a ±1.30

40W 40R 20S	6.40 ^{ab} ±1.16	5.77 ^{abc} ±1.70	6.23 ^{ab} ±1.74	6.30 ^{ab} ±1.97	6.27 ^a ±1.41
80W 15A 5S	6.37 ^{ab} ±1.96	6.20 ^{abc} ±2.02	6.27 ^{ab} ±1.76	6.63 ^a ±1.79	6.53 ^a ±1.36
70W 20A 10S	5.97 ^{abc} ±2.25	5.70 ^{abc} ±2.42	5.57 ^{ab} ±2.34	5.67 ^{ab} ±2.15	5.97 ^{ab} ±1.99
65W 20A 15S	5.07 ^{bcd} ±2.38	5.47 ^{abc} ±0.06	5.27 ^{ab} ±2.13	4.80 ^{bc} ±2.11	5.60 ^{abc} ±2.06
60W 25A 15S	4.30 ^d ±2.09	4.83 ^{bc} ±1.98	4.87 ^{ab} ±2.26	3.43 ^c ±1.91	4.33 ^c ±1.75
50W 30A 20S	5.37 ^{abcd} ±2.36	4.77 ^{bc} ±2.25	5.47 ^{ab} ±1.94	4.80 ^{bc} ±2.11	5.57 ^{abc} ±2.03
40W 40A 20S	4.67 ^{cd} ±2.37	4.60 ^c ±2.24	4.77 ^b ±2.11	5.73 ^{ab} ±2.60	4.70 ^{bc} ±2.56
100W (control)	6.77 ^a ±1.59	5.80 ^{abc} ±2.19	6.53 ^a ±1.83	6.10 ^{ab} ±2.37	6.93 ^a ±1.62
LSD	1.10	1.12	1.10	1.14	1.03

Values are means of data of triplicate determinations. Values in the same column having the same superscript are not significantly difference (P>0.05). W- Wheat, R- Rice, A –Acha, S- Soybean

In terms of appearance, there was no significant difference (P>0.05) between the samples and the control except for samples 65W20A15S, 60W25A15S, 50W30A20S, which were significantly different (P<0.05) from them. Appearance is an important attribute in food choice and acceptance (Muhimbula *et al.*, 2011). Outcome of sensory evaluation indicates that some samples were similar in appearance while others differed significantly. The noodles made from the samples that had rice were more appreciated in terms of appearance this could be attributed to the presence of rice that would have brightened its color a little. The noodles made from acha did not compare favorably with the control especially for sample 60W25A15S.

In terms of general acceptability, there was no significant difference (P>0.05) between noodles made from the samples blends and control except for those noodles made from samples 60W25A15S, 40W40A20S that were significantly different (P<0.05) from them. Although all the noodles made from the samples compared favorably with control (100W) which has already gained popularity for acceptability. This study shows that all the flour samples could be used for noodle and products will have good organoleptic acceptability.

4. CONCLUSION

Through this study, there have been a lot of interesting findings that noodles can be produced from blends of soybean, acha, rice and wheat flour. The use of composite flour in noodles production has

showed good potential improved product quality in terms of nutritional values as were observed in the proximate, mineral components, vitamins. At the same time, development and consumption of such blends will not only improve the nutritional status of the general population but also helps those suffering from degenerative diseases associated with today's changing lifestyles and environment. The sensory evaluation results indicated that, noodles produced from the blends of soybean, acha, rice and wheat flours had a good sensory attributes and were able to compare favourably with the control sample.

REFERENCES

- A.O.A.C. (1995). Official Methods of Analysis of the Association of Official Analytical Chemists, 15th ed., Association of Official Analytical Chemists, Arlington VA, pp. 1058-1059.
- A.O.A.C. (2000). Official Methods of Analysis (15thedn), Association of Official Analytical Chemists, Washington, DC.
- Aboubacar, A., Kirleis, A.W. and Oumarou, M. (1999). Important sensory attributes affecting consumers acceptance of sorghum porridge in West Africa as related to quality tests. *Journals Cereals Science*, 30:183-187.
- Adebowale, A. A., Sanni, L. O. and Awonorin, S. O. (2005). Effect of texture modifiers on the physico-chemical and sensory properties of dried "fufu". *Food Science Technology*, 11:373-385.
- Alvarenga, N. B., Lidon, F. C., Belga, E., Motrena, P., Guerreiro, S., Carvalho, M. J. and Canada J. (2011). Characterization of gluten-free bread prepared from maize, rice and tapioca flours using hydrocolloid seaweed agar-agar. *Recent Research in Science and Technology* 3(8): 64-68.
- Anna, W. M. and Malgorzata, K. (2011). Evaluation of the mineral composition of breadstuff and frequency of its consumption. *Acta Scientia Polonorum Technologia Alimentaria* 10: 487-495.
- Ayo, J. A., Ayo, V. A., Nkama, I. and Adewori, R. (2007). Physicochemical, In-Vitro Digestibility and Organoleptic Evaluation of "Acha" Wheat Biscuit supplemented with soybean flour. *Nigerian Food Journal*, 25(1): 77-89.
- Ayo, J. A., Ayo, V. A., Popoola, C, Omosebi, M. and Joseph, L. (2014). Production and evaluation of malted soybean-acha composite flour bread and biscuit. *African Journal of Food Science and Technology*. Vol 5(1). Pp 21-28.
- Bolarinwa, I. F., Olaniyan, S. A., Adebayo, L. O. and Ademola, A. A. (2015). Malted Sorghum-Soy Composite Flour: Preparation, Chemical and Physico-Chemical Properties. *Journal of Food Processing Technology*, 6:8
- Cantral, R.P, and Reeves, T.G (2002) .The cereal of the World's Poor takes center stage. *Science*.296: 53.Chemist Inc. U.S.A Pp.277-279.
- Ciclitira, P. J. and Ellis, H. J. (1987). Investigation of cereal toxicity in celiac disease. *Postgraduate Medicine Journal* 63: 767-775.
- Chen, Z. (2003). Physicochemical properties of sweet potato starches and their application in noodle products. Ph.D dissertation. No 90-5808-887-1. Wageningen University, The Netherlands.

- 385 Davison, A. G. F. and Bridges, M. (1987). Celiac Disease. *ClinicaChimicaActa* 163: 1-40.
- 386 Falola, A. O., Olatidoye, O. P., Balogun, I. O. and Opeifa, A. O. (2013). Evaluation of nutritional,
387 physicochemical properties and acceptability of dehulled ofada rice and soybean flour blends. *Journals*
388 *of Agriculture and Veterinary Sciences*, 5(1):118-125.
- 389 Gunathilake, K. D. P. P and Abeyrathne, Y. R. M. K. (2008). Incorporation of coconut flour into wheat flour
390 noodles and evaluation of its rheological, nutritional and sensory characteristics. *Journal of Food*
391 *processing and Preservation*, 32(1):133-142.
- 392 Haard, N. F. and Chism, G. W. (1996). "Characteristics of edible plant tissues, Food Chemistry", New
393 York: Marcel Dekker, Inc. pp. 943–1001.
- 394 Hosene, C.R (1986). Principles of Cereal Science and Technology.American Association of Cereal
395 <http://www.instantramen.org.jp/english/data/data02.html>.Retrieved May 1, 2008.
- 396 Ihekoronye, A. I. and Ngoddy, P. O. (1985).Integrated Food Science and Technology for the tropics.
397 Macmillan Publishers Ltd., London and Basingstoke, pp. 386.
- 398 Inyang, U. E. and Ekop, V. O. (2015). Physico-Chemical Properties and Anti-nutrient Contents ofUnripe
399 Banana and African Yam Bean Flour Blends. *International Journal of Nutrition and Food Sciences*, 4(5):
400 549-554.
- 401 Ishitani, K. E., Hakura, S. G and Esash, T. (1999). Calcium absorption from the ingestion of cereal
402 derived by calcium by human. *Journal of Nutrition Science Vitmanol* (Tokyo), 45(5):9-10.
- 403 Iwe, M. O. (2000). Effects of extrusion cooking on some functionalproperties of soy-sweet potato mixture:
404 A response surface Analysis. *Plant Foods for Human Nutrition*. 80: 169 – 184.
- 405 Iwe, M. O. (2003). The science and Technology of soybean.Rejoint communication services Ltd, Enugu.
406 Pp 43.
- 407 Jideani, A. I, and Akingbala J. A. (1993). Some physiochemical properties of acha(*Digiteriaexillis*) (acha)
408 *digiteriaiburua*. *Review Plant Food for Human Nutrition*, 54:362–373.
- 409 Kanu, P. J., Sandy, E. H., Kandeh, B. A., Behsoon, J. Z. and Huiming, Z. (2009). Production and
410 evaluation of breakfast cereal-based porridge mixed with sesame and pigeon peas for adults. *Pakistan*
411 *Journal of Nutrition*, 8(9):1335-1343.
- 412 Kim, Y. S., Wiesenborn, D. P., Lorenzen, J. H. and Berglund, P. (1996). Suitability of edible bean and
413 potato starches for starch noodles. *Cereal Chemistry*, 73: 302-308.
- 414 Lienel, H. H.(2002). Ash analysis. In: Introduction in Chemical Analysis of Food. Nielson, S. S. (Ed). CBS
415 Publishers, New Delhi. p 972.
- 416 Mannay, S. and Shadaksharaswamy, C. M. (2005). Foods: Facts and Principles. (2nd ed.).New Age
417 International Ltd. Publishers. New Delhi, India.
- 418 Mensah, P. and Tunkins, A. (2003). Household level technologies to improve the availability and
419 preparation of adequate protein and safe complementary foods In; *Food and Nutrition Bulletin*, 24:104-
420 125.

- 421 Miskelly, D. M. (1993). Noodles; a new look at an old food. *Food Australia* 45: 496-500
- 422 Muhimbula, H. S., Issa-Zacharia, A. and Kinabo, J. (2011). Formulation and sensory evaluation of
423 complementary foods from local, cheap and readily available cereals and legumes in Iringa, Tanzania.
424 *African Journal of Food Science*, 5(1):26 – 31
- 425 Nnam, N. M. (2001). Comparison of the Protein nutrition value of food blends based on sorghum,
426 bambara groundnut and sweet potatoes In: *International Journal of Food Sciences and Nutrition*, 52:25-
427 29.
- 428 Obafunmi, M. O, Balami Y. A, and Christiana B. N. O (2009). Acha as a Potential Substitute raw material
429 for the Production of Malt drink. In: Presented at the international Seminar on Raw Materials Research
430 Development and Utilization held at the Raw Materials Research and Development Council (RMRDC)
431 Headquarters, Maitama, Abuja: 10-13.
- 432 Ojmelukwe, P. C., Asumugha, V. C. and Omeire, G. C. (2005). Fundaments of Food Science and
433 Nutrition. Bel's Book Publisher Nigeria.
- 434 Okafor, G. I. and Ugwu, F. C. (2014). Production and evaluation of cold extruded and baked ready-
435 to-eat snacks from blends of breadfruit (*Treculia africana*), cashewnut (*Anacardium occidentale*) and
436 coconut (*Cocos nucifera*). *Food Science and Quality Management*, 23.
- 437 Okaka, J. C. (2005). Handling, Storage and processing food plant. OCJ academic publishers Enugu.
- 438 Omeire, G. C., Kabuo, N. O., Nwosu, J. N. Peter-Ikechukwu, A. and Nwosu, M. O. (2015). Enrichment of
439 Wheat/Cassava Noodles with Partially Defatted Protein-Rich Flours. *Journal of Environmental Science,*
440 *Toxicology and Food Technology*, 9(5): 121-125.
- 441 Omeire, G. C., Umeji, O. F., and Obasi, N. E. (2014). Acceptability of Noodles Produced from Blends of
442 Wheat, Acha and Soybean Composite Flours. *Nigeria Food Journal* 32(1): 31-37.
- 443 Onwuka, G. I. (2005). Food Analysis. Naphtali Publishers. Lagos Nigeria.
- 444 Opeifa, A. O., Olatidoye, O.P., Adesala, S. O. and Fayomi, M. J. (2015). Production and Quality
445 Evaluation of Ogi Produced from Fermented Maize and Horse Eye Bean (*Mucuna urens*). *Pakistan Journal*
446 *of Nutrition* 14 (7): 417-425.
- 447 Oppong, D., Eric A., Samuel O. K., Eric B. and Patrick S. (2015). Proximate Composition and Some
448 Functional Properties of Soft Wheat Flour. *International Journal of Innovative Research in Science,*
449 *Engineering and Technology*.
- 450 Osaretin, A., Ebuehi, T. and Oyewole, A. C. (2007). Effect of cooking and soaking on physical
451 characteristics, nutrient composition and sensory evaluation of indigenous and foreign rice varieties in
452 Nigeria. *African Journal of Biotechnology*, 6(8): 1016-1020.
- 453 Perez, C. M, Juliano, B. O, Pascual, C. G, and Novenario, V. G. (1987). Extracted lipids and
454 carbohydrates during washing and boiling of milled rice. *Starch* 39: 386-390. Presented at the
455 International Seminar on Raw Materials Research Development and Utilization held at Raw Materials
456 Research and Development Council (RMRDC) Headquarters, Maitama, Abuja: 10 – 13.
- 457 Plahar, W. A., Okezie, B. O. and Gyato, C. K. (2003). Development of a high protein weaning food by
458 extrusion cooking using peanuts, maize and soybeans. *Plant Foods for Human Nutrition*, 58: 1-12.
- 459 Sanni, L.O., Bamgbose, C.A., Babajide, J.M. and Sanni, S.A. (2007). Production of instant cassava
460 noodles. In: Proceedings of the 13th Triennial Symposium of the International Society for Tropical Root
461 Crops (ed. Kapinga et al.), 10 – 14 Nov, 2003, Arusha, Tanzania, pp. 466 – 472.

- 462 Sanni, S. A, Okeleye, K. A., Soyode, A. F., Taiwo, O. C. (2005). Physio-chemical properties of early and
463 medium maturing and Nigerian rice varieties. *Nigeria Food Journal*, 23: 148-152.
- 464 Srilakshmi, B. (2009). Food Science. New Age International Ltd, publishers, New Delhi. Pp 88 – 90, 115.
- 465 Szezesniak, A. S. (2002). Texture is a sensory property. *Food Quality Preference*, 13:215-225.
- 466 Wardlaw, G. M. (2004). Perspectives in Nutrition. (6th ed.). McGraw Hill Companies, New York, U.S.A.
- 467 WINA. (2008). Instant Ramen facts worldwide (World Instant Noodle Association).