

Original Research Article

Gene encoding bacterial wilt related protein as a target for stress resistant transgenic plants

ABSTRACT

Background & Aim: Bacterial Wilt is an important and crucial disease of tomato causing decrease in crop yield per year. In order to produce wilt resistant varieties exploration of defense mechanism against bacterial wilt is important. Aim of study is to confirm the expression of some of suspected defense genes reported to be produced in bacterial wilt resistant varieties in comparison to susceptible varieties.

Place and Duration of Study: Study was done in Department of Biochemistry and Molecular Biology, University of Gujrat Pakistan for one year for fulfillment of masters of Philosophy degree of research student.

Study design: Salicylic acid was deliberated on tomato var. Roma (medium resistant) and Riogrande (resistant variety) for three consecutive days. Expression analysis studies of iron ATP-binding cassette (ABC) transporters, ATP synthase and chaperonin using RT-PCR along with measurement of antioxidants activities.

Results: Iron ABC transporters was expressed in control and within 1st day of SA treatment in both varieties, while chaperonin was expressed after two days in Roma and after three days in Riogrande of SA treatment. ATP synthase was expressed after three days in Riogrande while started from 1st day in Roma after SA treatment. Peroxidase and catalase activity increased in both varieties after one day; with little decrease after two days of SA treatment.

Conclusion: Expression of iron ABC transporters, ATP synthase and chaperonin in response to SA application signifies importance of these genes in response to bacterial wilt in tomato.

Keywords: Salicylic acid, Tomato, ATP synthase, Iron ABC transporter, Chaperonin.

1. INTRODUCTION

Tomato (*Solanum lycopersicum*) is second world most significant vegetable crop, and for last 25 years total world production and consumption of this important crop has raised moderately. Small genome, short life cycle similar to many commercially vital plants (potato, eggplant, peppers and tobacco), made it important as studies conducted on tomatoes are equally valid to these plants [1; 2]. It is challenged by various biotic stresses and abiotic stresses; became chief reason for tomato loss worldwide [3]. Tomato respond to biotic/abiotic stresses by multiple defense mechanisms including cellular reactions generated by the pathogens leads to signaling pathways [4; 5]. Plant immune systems dependent on resistant/susceptible variety and their defense mechanisms [6].

Main objective of studying plant pathogen interactions is to reveal the signaling molecules that lead to defense related pathways [7]. Signaling molecules/hormones including auxins, cytokinins, salicylic acid (SA), abscisic acid (ABA), gibberellic acid (GA), ethylene (ET), jasmonic acid (JA), secreted by plants in response to environmental stresses have important functions in production of protein related to defense, growth and development [8; 9]. SA has an important role in plant defense against biotic and abiotic stresses, along with regulation of various physiological and biochemical processes (Plant growth, photosynthesis, respiration, thermotolerance, stomatal responses, nodulation, nitrate metabolism, flowering), and ET biosynthesis [10; 11; 12]. Increased SA in plant cells deliberately/naturally is also coupled with augmentation of pathogenesis related proteins (PR) proteins along with defense and metabolism related proteins resulted in systemic acquired resistance (SAR) [13; 14; 15]. It had been reported for its critical role in defense aided with organization of SAR after biotrophic and hemi-biotrophic pathogens infections specifically in model plants [16]. SAR is a potent form of innate immunity which is efficient and a long-term resistance effect against a wide variety of pathogens [17]. SA had been also reported to produce oxidative stress in plants at high concentration but at low concentrations enhanced antioxidative capability help to cope with stresses [18]. Antioxidant enzymes such as polyphenol oxidase, peroxidase (POD), catalase (CAT) were reported to increase 4 days after SA treatment (War et al. 2011). Antioxidant activity after SA treatment can be a measure of resistance or susceptibility of the varieties.

Taking into account the mechanisms which plants employ to defend themselves against pathogens may enable the researchers to design novel strategies to improve disease resistance in crop plants. ATP synthase, iron ATP-binding cassette (ABC) transporters and chaperonin related to protein destination, storage synthesis and photosynthesis expression were some of the important proteins expressed in plants resistant to biotic stress in response to SA treatment [19; 20]. Previously SA induces proteins were explored in susceptible and resistant varieties of tomato (Roma and Pant Bahr)[15]. In current study ATP synthase, iron ABC transporters and chaperonin were confirmed using RTPCR analysis in response to SA in Riogrande (resistant) and Roma (medium resistant) to bacterial wilt. Along with these protein encoding genes, antioxidant enzymatic assays (POD, CAT) were carried out after one, two and three days of SA treatment.

2. MATERIALS AND METHODS

2.1. Plant Material

Tomato (*Solanum lycopersicum*) Riogrande (resistant) and Roma (medium resistant) cultivars were used. The seeds of Riogrande and Roma were obtained from National Agriculture Research Centre (NARC), Islamabad. Seeds were surface sterilized in 0.8 % (v/v) "Clorox" bleach (Sodium hypochlorite) as reported [15]. Seedlings were planted in pots containing sand and soil, grown in growth chamber under white fluorescent light ($600 \text{ mol m}^{-2} \text{ s}^{-1}$; 16 h light/8 h dark) at 25°C and 70% relative humidity.

2.2. SA Treatment, RNA Isolation and cDNA Synthesis

SA (Wako) was used at 1 mM concentration prepared in water. Three-week-old seedlings of the Roma and Riogrande were treated with SA for 1, 2 and 3 days and RNA extraction was done for 3 consecutive days. Total RNA was isolated from leaf cells of controlled and treated tomato plants using the TRI Reagent (Cat. No. TRI. 118; Lot No. 4221). RNA was precipitated by mixing it with isopropanol, centrifuged (12000 g) for 8 min at 4-25° C. RNA pellet was washed with 75% ethanol, centrifuged, dissolved in nano-pure water and stored. cDNA was prepared from total RNA samples using cDNA synthesis kit (Invitrogen, USA).

2.3. RT-PCR Analysis

Accession number of five proteins(given in Table 1) were analyzed using bioinformatics tools, and relatively suitable length of aligned region of PCR primers were designed using Primer3 online software. Forward and reverse primers were designed for Iron ABC transporter, ATP synthase and 60-KDa chaperonin. cDNA was amplified with gene-specific primers using PCR. Confirmation of the desirable sequences in extracted cDNA from all species will be done by running PCR reaction with five set of primers under specific conditions (Table 1). 20 µl PCR reaction mixtures contains 1.5 of the template cDNA, with 2 µl Taq buffer, 1 µl MgCl₂, 0.5 µl dNTP mixture, 0.5 µl of each forward and reverse primer, 1 µl Taq polymerase and 13 µl of nanopure water. Initial denaturation step for PCR is restricted to (95°C) for 4 minutes, followed by 40 cycles of denaturation step at (95°C) for 30 seconds, annealing temperature (55°C for ATP synthase, 53°C 60-kDa chaperonin and iron ABC transporter) for 1.5 minutes and extension (72°C) for 30 sec, and a final extension cycle of 7 min (72°C). The amplified PCR products were analyzed by 1 % (w/v) agarose gel electrophoresis.

2.4. Enzymatic Assays

2.4.1. POD activity

POD activity was determined following the dehydrogenation of guaiacol as a substrate [21]. Plant material (2g) is grinded in pre-chilled mortar with 0.1 M phosphate buffer (pH 7). Homogenate is centrifuged at 18000 g for 15 min at 5°C. Supernatant is used as enzyme source within 2-4 h. 3 ml of the buffer solution is pipetted along with 0.05 ml guaiacol solution (20 mM), 0.1 ml enzyme extract and 0.03 ml hydrogen peroxide solution (12.3 mM) in cuvette. Absorbance was taken on spectrophotometer at 750 nm.

2.4.2. CAT Assay

CAT was assayed by monitoring the decrease of absorbance as described by Aebi[22]. The reaction mixture consisted of 50 µL of enzyme, 2.5 mL potassium phosphate buffer(50 mM; pH 7.6), 200 µl hydrogen peroxide (12.5 mM) and homogenate sample (50 µl), along with 250 µl of distilled water were taken in a cuvette. Absorbance was measured at 240 nm three times for each sample with a time interval of 30 seconds. One unit activity of CAT is described as an absorbance change of 0.01 as unit per minute.

3. Results and Discussion

Bacterial wilt is one of the important bacterial stresses faced by tomato, restraining its production [23]. Different genes had been identified which are expressed in tomato varieties resistant and susceptible to bacterial wilt [24; 25]. SA mediated defense pathway play a key role in response to biotic stresses[26]. RTPCR is widely used for confirmation of gene or protein expression was used for confirmation of Iron ABC transporter, ATP synthase and chaperonin 60 in Riogrande and Roma [27]. RNA extraction was done with 3-W-Old seedlings leaves, followed by cDNA synthesis and RTPCR of Iron ABC transporter, ATP synthase and chaperonin 60 in resistant and medium resistant varieties of tomato (Fig S1A; S1B).

3.1. Morphological variation

Leaves became light green after 1, 2 and 3 days of SA treatment in comparison to control in Roma (Fig 1A, 1B, 1C, 1D). Riogrande leaves almost remain same as control (Fig 1E, 1F, 1G, 1H). Reduction in chlorophyll in semi-resistant variety (Roma) show its more sensitiveness to water stress, as the leaves were treated with aqueous solutions of SA in comparison to control. It is in line with observation for Roma [15].

3.2. Antioxidant enzymes assay

POD and CAT activity was increased one day after treatment of SA, decrease after 2 days and again increased after 3 days of SA treatment in Riogrande and Roma (Fig 2A; 2B). CAT and POD help to scavenge the stress damage were confirmed in these two varieties primarily. Antioxidant activity was more prominent in resistant variety suspects its role to reduce the ROS damage. Higher induction of antioxidant enzymes (POD, PPO) along with generation of H_2O_2 were reported in response to SA (1, 1.5, 2 mM) in *Cicer arietinum* L. [28]. Antioxidative enzymes POD, thioredoxin, superoxide dismutase, glutathione S-transferase were reported in tomato after SA treatment to protect plant from damage[29; 30]. H_2O_2 in plants is generated during photorespiration, mitochondrial electron transport and β -oxidation of fatty acids, and its intracellular level are stringently controlled for maintaining homeostasis in the cell [31]. Immune response is suppressed by inhibition of host cell's POD activity, thereby inhibiting ROS production [32].

3.3. Iron ABC transporter

Iron ABC transporter was found to be expressed within one day of SA treatment in both Roma and Riogrande (Fig 3A). SA enhanced Iron ABC transporter expression in Roma and in Riogrande after 1, 2 and 3 days of SA treatment (Fig 3A). Iron ABC transporter, was also upregulated in the proteomics analysis of tomato cvs and soybean in response to application of either JA or SA [15; 20]. Iron ABC transporter had a role in stress resistance in plants, and induced in response to bacteria, fungus, heavy metal stress, ET, SA, JA, ABA, resistance to HR, iron homeostasis, cell viability and iron acquisition in plants [30; 33; 34]. Iron ABC transporter, photosynthesis related gens, defense genes, antioxidant enzymes were induced in response to SA treatment can be used as genetic tools to investigate disease resistance [35]. ABC transporter genes reduction indirectly resulted in reduced photosynthesis, and production of stress in apple [36]. Iron ABC transporter is also induced in JA dependent pathway in response to infection by bacterial and fungal infections in tobacco [37]. Its expression in response to JA, SA, bacterial and fungal pathogens; made it favorite gene responsible for susceptibility or resistance of tomato cultivars Roma and Riogrande.

3.4. ATP Synthase

ATP synthase was found to be expressed within one day of SA treatment in Roma along with control, while in Riogrande it is expressed in control and 3 days after SA treatment (Fig 3B). It is in line with observation that insusceptible cultivars ATP synthase is up-regulated and in resistant cultivar it is expressed after 2 days [19]. Proteomic approach show reduced expression of ATP synthase with 1, 2 and 3 days of SA treatment in same cultivar Roma [15]. Wu et al. [10] via gel free techniques (LC MS/MS) reported phosphoproteins such as ATP synthase and iron ABC transporters are upregulated by SA in maize. ATP synthase is expressed in response to biotic/abiotic stresses for production of resistance in plants by suppressing HR response, along with increased metabolism of glucose [38]. Janda et al. [39] reported regulatory effects of SA on photosynthetic electron transport processes. Chloroplastic-ATP synthase was found to be important to trigger ET, SA/JA pathway in response to insect herbivory [40]. Transgenic tobacco with ATPase exhibited resistance in response to *P. syringae*, suggesting its role in defense via JA or SA [41]. Upregulation of ATP synthase after 1st day of treatment in susceptible (Pant Bahr) and medium resistant varieties of tomato and after 2 days in resistant variety of tomato implied its importance as defense gene in response to biotic stresses,

insect herbivory, and suppression of HR made it a good candidate transgenic tomato with enhanced resistance to bacterial wilt.

3.5. Chaperonin

Chaperonin was expressed in control and 2 days after SA treatment in Roma (Fig 3C) and after 3 days of SA treatment in Riogrande (Fig 3D). SA enhanced Chaperonin expression in Roma and in Riogrande after 2 and 3 days of SA treatment. Proteomic approach show enhanced expression of Chaperonin in susceptible variety (Pant Bahr) and unchanged expression in medium resistant variety (Roma) after SA treatment. Wu et al. [27] reported that after inoculation with *P. solanacearum*; ATP synthase was expressed in susceptible and chaperonin was expressed in resistant maize cultivar in SA pathway. Moshe et al. [42] reported up regulation of chaperonin in resistant tomato cultivar in response to tomato leaf curl virus infection in comparison to susceptible cultivar. Proteomics analysis of *Arabidopsis* associate chaperonin-60 with photosynthesis for increasing duration of stress [43]. Cueto-Ginzo et al. [44] also reported SA balance photosynthetic reduction during infection of potato virus in tomato by increasing mesophyll conductance. Molecular Chaperone is another candidate found to be important for resistance in resistant cultivars in response to stresses in comparison to accumulation of ROS species in susceptible cultivars. Chaperone was expressed in resistant, medium resistant and in susceptible cultivars; more prominent in resistant variety (Riogrande) (Fig 3D).

4. CONCLUSION

Proteomics technique was used for identification of iron ABC transporter, ATP synthase and chaperonin in resistant and susceptible varieties of tomato [15]. Proteomics analysis data show chaperonin, ATP synthase, iron ABC transporter which were upregulated/remain unchanged in one variety in comparison to other in response to application of SA are candidates proteins that can be responsible for susceptibility or resistance of tomato cultivars. RTPCR show iron ABC transporter expression remain same in resistant and medium resistant cultivars. While chaperonin expression was important; as expressed strongly after 3 days of SA application in resistant cultivars also

supported by previous reports. Similarly ATP synthase expression was after 2 days in Riogrande. CAT and POD enzymatic activity was increased in both cultivars after one and more after 3 days of SA treatment little more in Riogrande. Genes encoding proteins related to protein folding and photosynthetic genes are targets for biotic stress related pathway in tomato.

References

1. Peterson DG, Pearson WR and Stack SM. Characterization of the tomato (*Lycopersicon esculentum*) genome using *in vitro* and in situ DNA reassociation. *Genome*. 1998;41(3):346-56.
2. Kimura S, Sinha N. Tomato (*Solanum lycopersicum*): a model fruit-bearing crop. *Cold Spring Harb Protoc*. 2008;11:1-9.
3. Kissoudis C, Chowdhury R, van Heusden S, de Wiel CV, Finkers R, Visser RGF, *et al*. Combined biotic and abiotic stress resistance in tomato. *Euphytica*. 2015;202(2):317-32.
4. Pangesti N, Pineda A, Pieterse CMJ, Dicke M, VanLoon JJA. Two-way plant-mediated interactions between root associated microbes and insects: from ecology to mechanisms. *Front Plant Sci*. 2013;4:414. doi: 10.3389/fpls.2013.00414.
5. Schenk PM, Kazan K, Wilson I, Anderson JP, Richmond T, Somerville SC, *et al*. Coordinated plant defense responses in *Arabidopsis* revealed by microarray analysis. *Proc Natl Acad Sci USA*. 2000;97(21):11655-60.
6. Da Cunha L, McFall AJ, Mackey D. Innate immunity in plants: a continuum of layered defenses. *Microb Infect* 2006;8(5):1372-81.
7. Abu Qamar S, Luo H, Laluk K, Mickelbart MV, Mengiste T. Crosstalk between biotic and abiotic stress responses in tomato is mediated by the AIM1 transcription factor. *Plant J* 2009;58(2):347-60.
8. Clarke JD, Volko SM, Ledford H, Ausubel FM, Dong X. Roles of salicylic acid, jasmonic acid, and ethylene in cpr-induced resistance in *Arabidopsis*. *Plant Cell*. 2000;12(11): 2175-90.
9. Kunkel BN, Brooks DM. Cross talk between signaling pathways in pathogen defence. *Curr Opin Plant Biol*. 2002;5:325-31. doi:10.1016/S1369-5266(02)00275-3.
10. Wu L, Hu X, Wang S, Tian L, Pang Y, Han Z, *et al*. Quantitative analysis of changes in the phosphoproteome of maize induced by the plant hormone salicylic acid. *Sci Rep*. 2015;5:18155. doi: 10.1038/srep18155.

- 220 11. An C, Mou Z. Salicylic acid and its function in plant immunity. J Integr Plant Biol. 2011;53:412-28.
- 221 12. Zandalinas SI, Rivero RM, Martinez V, Gomez-Cadenas A, Arbona V. Tolerance of citrus plants to the
- 222 combination of high temperatures and drought is associated to the increase in transpiration modulated by
- 223 a reduction in abscisic acid levels. BMC Plant Biol. 2016;16:105.
- 224 13. Qi PF, Johnston A, Balcerzak M, Rocheleau H, HarrisLJ, Long XY, et al. Effect of salicylic acid
- 225 on *Fusarium graminearum*, the major causal agent of fusarium head blight in wheat. Fungal Biol.
- 226 2012;116(3):413-26.
- 227 14. Dong CJ, Wang XL, Shang QM. Salicylic acid regulates sugar metabolism that confers tolerance to
- 228 salinity stress in cucumber seedlings. Sci Hort. 2011;129(4):629-36.
- 229 15. Afroz A, Khan MR, Komatsu S. Determination of proteins induced in response to jasmonic acid and
- 230 salicylic acid in resistant and susceptible cultivars of tomato. Protein Pept Lett.2010;17(7):836-46.
- 231 16. Yasuda M, Ishikawa A, Jikumaru Y, Seki M, Umezawa T, Asami T, et al. (2008) Antagonistic
- 232 interaction between systemic acquired resistance and the abscisic acid-mediated abiotic stress response
- 233 in *Arabidopsis*. Plant Cell. 2008;20(6):1678-92.
- 234 17. Vlot AC, Dempsey DMA, Klessing D. Salicylic acid, amulti-faceted hormone to combat disease. Annu
- 235 Rev Phytopathol. 2009;47:177-206. doi: 10.1146/annurev.phyto.050908.135202.
- 236 18. Horvath E, Szalai G, Janda T. Induction of abiotic stress tolerance by salicylic acid signaling. J Plant
- 237 Growth Regul2007;26(3):290-300.
- 238 19. Kohzuma K, Dal Bosco C, Meurer J, Kramer DM. Light-and metabolism-related regulation of the
- 239 chloroplast ATP synthase has distinct mechanisms and functions. J Biol Chem. 2013;288(18):13156-
- 240 20.Eichhorn H, Klinghammer M, Becht P, Tenhaken R. Isolation of a novel ABC-transporter gene from
- 241 soybean induced by salicylic acid. J Exp Bot. 2006;57:2193-201. doi:10.1093/jxb/erj179.
- 242 21. Malik CP, Singh MB. Extraction and estimation of amino acids and keto acids. In: Malik CP, Singh
- 243 MB, editors. Plant Enzymology and Histo-Enzymology. New Delhi: Kalyani Publishers; 1980: p. 286.
- 244 22. Aebi H. Catalase *in vitro*. Methods in Enzymology 105; 1984:121-126.
- 245 23. Hayward AC. Biology and epidemiology of bacterial wilt caused by *Pseudomonas solanacearum*.
- 246 Annu Rev Phytopathol. 1991;29:65-87.

- 247 24. Glick E, Levy Y, Gafni Y. The viral etiology of tomato yellow leaf curl disease-a review. *Plant Prot Sci.*
248 2009;45(3):81-97.
- 249 25. Afroz A, Khan MR, Ahsan N, Komatsu S. Comparative proteomic analysis of bacterial wilt susceptible
250 and resistant tomato cultivars. *Peptides* 2009;30(9):1600-7.
- 251 26. Novakova M, Sasek V, Dobrev PI, Valentova O, Burketova L. Plant hormones in defense response of
252 *Brassica napus* to *Sclerotiniasclerotiorum*. Reassessing the role of salicylic acid in the interaction with a
253 necrotroph. *Plant PhysiolBiochem.* 2014;80:308e317.
- 254 27. Wu L, Wang S, Chen X, Wang X, Wu L, Zu X, et al. Proteomic and phytohormone analysis of the
255 response of maize (*Zea mays* L.) seedlings to sugarcane mosaic virus. *PLoS ONE* 2013;8(7): e70295.
256 doi:10.1371/journal.pone.0070295.
- 257 28. War AR, Paulraj MG, War MY, Ignacimuthu S. Role of salicylic acid in induction of plant defense
258 system in chickpea (*Cicer arietinum* L.). *Plant Signal Behav.* 2011;6(11):1787-92.
- 259 29. Wan XL, Zhou Q, Wang YY, Wang WE, Bao MZ, Zhang JW. Identification of heat-responsive genes
260 in carnation (*Dianthus caryophyllus* L.) by RNA-seq. *Front Plant Sci.* 2015;6:519. doi:
261 10.3389/fpls.2015.00519
- 262 30. Singh AP, Dixit G, Mishra S, Dwivedi S, Tiwari M, Mallick S, et al. Salicylic acid modulates arsenic
263 toxicity by reducing its root to shoot translocation in rice (*Oryza sativa* L.). *Front Plant Sci.* 2015;6:340.
264 doi: 10.3389/fpls.2015.00340
- 265 31. McClung CR. Regulation of catalases in *Arabidopsis*. *Free RadicBiol Med.* 1997;23:489-96.
266 doi:10.1016/S0891-5849(97)00109-3.
- 267 32. Liu P, Zhang H, Yu B, Xiong L, Xia Y. Proteomic identification of early salicylate- and flg22-responsive
268 redox-sensitive proteins in *Arabidopsis*. *Sci Rep.* 2015;5:8625. doi 10.1038/srep08625.
- 269 33. Song WY, Park J, Mendoza-Cozat DG, Suter-Grotemeyer M, Shim D, et al. Arsenic tolerance in
270 *Arabidopsis* is mediated by two ABCC-type phytochelatin transporters. *Proc Natl AcadSci USA.*
271 2010;07:21187-92. doi:10.1073/pnas.1013964107.
- 272 34. Borghi L, Kang J, Ko D, Lee Y, Martinoia E. The role of ABCG-type ABC transporters in
273 phytohormone transport. *BiochemSoc Trans.* 2015;43:924-30. doi:10.1042/BST20150106.

- 274 35. Zhang X, Dong J, Liu H, Wang J, Qi Y, Liang Z. Transcriptome sequencing in response to salicylic
275 acid in *Salvia miltiorrhiza*. PLoS ONE 2016;11(1):e0147849. doi:10.1371/journal.pone.0147849.
- 276 36. Wu T, Wang Y, Zheng Y, Fei Z, Dandekar AM, Xu K, et al. Suppressing sorbitol synthesis
277 substantially alters the global expression profile of stress response genes in apple (*Malus domestica*)
278 leaves. Plant Cell Physiol. 2015;56(9):1748-61. doi: 10.1093/pcp/pcv092.
- 279 37. Stukkens Y, Bultreys A, Grec S, Trombik T, Vanham D, Boutry M. NpPDR1, a pleiotropic drug
280 resistancetype ATP binding cassette transporter from *Nicotianaplumbaginifolia*, plays a major role in plant
281 pathogen defense. Plant Physiol. 2005;139:341-52.
- 282 38. Sugimoto M, Yamaguchi Y, Nakamura K, Tatsumi Y, Sano H. A hypersensitive response-induced
283 ATPase associated with various cellular activities (AAA) protein from tobacco plant. Plant Mol Biol.
284 2004;56(6):973-85.
- 285 39. Janda K, Hideg E, Szali G, Kovacs L, Janda T (2012) Salicylic acid may indirectly influence the
286 photosynthetic electron transport. J Plant Physiol.2012;169(10):971-8.
- 287 40. Schmelz EA, LeClere S, Carroll MJ, Alborn HT, Teal PE. Cowpea chloroplastic ATP synthase is the
288 source of multiple plant defense elicitors during insect herbivory. Plant Physiol. 2007;144(2):793-805.
- 289 41. Lee MH, Sano H. Suppression of salicylic acid signaling pathwaysby an ATPase associated with
290 various cellular activities(AAA) protein in tobacco plants. Plant Biotech. 2007;24:209-15.
- 291 42. Moshe A, Pfannstiel J, Brotman Y, Kolot M, Sobol I, Czosnek H, et al. Stress responses to tomato
292 yellow leaf curl virus (TYLCV) infection of resistant and susceptible tomato plants are different.
293 Metabolomics. 2012;S1:006. doi:10.4172/2153-0769.S1-006.
- 294 43. Salvucci ME. Association of Rubisco activase with chaperonin-60 β : a possible mechanism for
295 protecting photosynthesis during heat stress. J Exp Bot.2008;59(7):1923-33. doi:10.1093/jxb/erm343
- 296 44. Cueto-Ginzo AI, Serrano L, Bostock RM, Ferrio JP, Rodriguez R, Arcal L, et al. Salicylic acid mitigates
297 physiological and proteomic changes induced by the SPCP1 strain of Potato virus X in tomato plants.
298 PhysiolMol Plant Path.2016;93:1e11.
- 299 **Abbreviations:**Salicylic acid (SA), Systemic acquired resistance (SAR), Absciscic acid (ABA),
300 Gibberellic acid (GA), Ethylene (ET), Jasmonic acid (JA), Pathogenesis related (PR), Catalase (CAT),
301 Peroxidase (POD), ATP-binding cassette transporters (ABC transporters).

Table 1. Primers for genes encoding protein sequences, melting temperature TM and their functional category.

Protein Name	Protein database/ Tm	Forward Primer	Functional Category
ATP synthase epsilon chain (F)	gi 114610/ 55°C	5-3' TGACCTTAAATCTTAGTGTACTGACC	Glycolysis pathway
ATP synthase epsilon chain (R)		5-3' TTATGAAATCGGATTGATAGCC	
60 kDa chaperonin (F)	gi 603912/ 53°C	5-3' GAGATTGTGTTTCGACCAGGAG	Protein destination and storage
60 kDa chaperonin (R)		5-3' ACACGCCATCGTTGATAACC	
Iron ABC transporter (F)	gi 57504871/ 53°C	5-3' ATGAAAAAATTTTAATCATTATGAGTTTA	Electron transport and photosynthesis
Iron ABC transporter (R)		5-3' TTAAAAATTTTGATTTTCTAATCTTGTTTT	

Figures




Fig 1

Fig 1. Morphological variation of *Solanum lycopersicum* cvs Roma (a, b, c and d) and Riogrande (e, f, g & h) in control and 1, 2 and 3 days after SA treatment.




Fig 2A




Fig 2B

Fig 2A. Peroxidase activity in *S. lycopersicum* cvs Roma and Riogrande 1, 2 and 3 days after 1 mM SA treatment. 2B. Catalase activity in tomato cvs Roma and Riogrande 1, 2 and 3 days after 1 mM SA treatment.




Fig 3

Fig 3. RTPCR of 3-W-Old *S. lycopersicum* leaves after 1, 2 and 3 day treatment of SA using **3A**. **ABC** transporter primer (220 bp). L 2-5 containing PCR products of Roma control, 1, 2 & 3 days after treatment of SA. L 6-9 containing PCR products of Riogrande control, 1, 2 & 3 days after treatment of SA. L1 100 bp marker. **3B**. With ATP Synthase primer (520 bp). L 1: 1 Kb ladder, L2-L5: PCR product of control, 1, 2 & 3 days after treatment of SA of Roma respectively, L6-L9 contain PCR product of control, 1, 2 & 3 days after treatment of SA of Riogrande respectively, L11- 13 contain PCR product of Riogrande 1, 2 & 3 days after SA treatment respectively as repeat. **3C**.With Chaperonin primer (220 bp). L 1: 100 bp ladder, L2: PCR product of control, L5, L6, L7 1, 2 & 3 days after treatment of SA of Roma respectively. **3D**. With Chaperonin primer (220 bp). L 1: 100 bp ladder, L2: PCR product of control, L3, L4, L5 1, 2 & 3 days after treatment of SA of Riogrande respectively.




Fig S1

Fig S1. (S1A) RNA of 3-W-old *S. lycopersicum* seedlings leaves of Roma. RNA extracted from 3-W-Old tomato leaves. L 1 is showing the bands of RNA of control while L 2, 3, 4 & 5 are showing of treated (S1B) cDNA samples from RNAs extracted after 1, 2 & 3 day of SA treatment of Roma and Riogrande.