

Original Research Article

**Effects of Feed Additives on Productive and Reproductive Performance of
Khaki Cambell Duck in Kuluarchor Upazilla of Kishoreganj District,
Bangladesh**

ABSTRACT

Khaki Campbell duck being an important poultry species can contribute efficiently in increasing egg and meat production in the Kuliarchar Upazilla, Kishoreganj district. The study was carried out in Kuliarchar upazilla, Kishoreganj during the period of August 2014 to May 2015. A total of 120 female day-old ducklings (Khaki Campbell) were procured from the local farmer from Kuliarchor Upazilla. The ducklings were randomly distributed into 4 treatment groups i.e group A, group B, group C and group D (untreated control), were treated by Digester[®], Vitapoultry[®], Alqurzim[®] respectively which is marketed by ACI animal health Limited. All ingredients were supplied 2g/kg feed daily. Body weight was increased day by day with the helping of feed additives. Digester[®], Vitapoultry[®], Alqurzim[®] had direct effects on the growth of ducklings. The weight was higher in group B treated by Vitapoultry[®] in every stage of weighing compare with other. The body weight of control group was lower than the treated group. There was a significant ($p < 0.01$) difference between the treated group and control group. The egg production of Khaki Campbell ducks recorded at 24-28 weeks, 28-32 weeks, 32-36 weeks, 36-40 weeks of age at group A, group B, group C and group D were 19.70 ± 0.64 , 19.78 ± 0.61 , 18.76 ± 0.56 , 18.51 ± 1.25 ; 22.81 ± 0.67 , 22.67 ± 0.70 , 23.56 ± 0.87 , 21.56 ± 0.67 ; 18.44 ± 0.45 , 20.09 ± 0.38 , 17.91 ± 0.56 , 17.62 ± 0.26 ; 14.61 ± 0.69 , 15.83 ± 1.65 , 15.16 ± 0.46 , 14.42 ± 0.79 number respectively. There was a significant difference ($p < 0.01$) between the treated group and control group in 32-36 weeks of age. The weight of the egg of treated group was higher than the control group. The

onset of puberty of Khaki Campbell recorded at group A, group B, group C and group D were 23.9±0.12, 22.54±0.17, 23.14±0.57, 24.14±0.61 weeks respectively. The fertility of egg (%) of Khaki Campbell recorded at group A, B, C and D were 83.78±0.57, 85.44±2.46, 84.28±0.24 and 78.49±2.74 respectively. The Hatchability (%) of Khaki Campbell recorded at group A, B, C and D were 67.27±8.09, 68.16±7.68, 64.94±0.97, 58.85±3.83. The hatchability (%) highest in treated groups compared with the control group. Feed additives had tremendous result both productive and reproductive performance of Khaki Campbell duck compared with control group in Kuliarchar Upazilla, Kishoreganj district. Specifically Vitapoultry[®] had very good result when mixed with the basal feed and provided daily to the duck. So, my recommendation is to use vitaamin, mineral and amino acid containing feed additives with basal feed for enhancing the productive and reproductive performance not Khaki Campbell but all other breeds of duck in Bangladesh. Further studies need to develop an appropriate implementation of using feed additives with basal feed in ducks in Bangladesh.

Keyword: Duck, feed additives, performance, Kishoreganj, Bangladesh.

1. INTRODUCTION

The economy of Bangladesh largely depends on agriculture and it is the principal occupation of the rural people. The Government has identified agriculture and rural development as the topmost priority sector for rapid poverty reduction. The rural economy as a whole contributes more than 56% of total GDP where the contribution of rural non-farm sector is 36.71% [1]. Agriculture generates 48.4% of total employment, contributes a quarter of total export earnings and provides food security for the increasing population [2]. The supply and demand gap of animal protein can be met by increasing the production of poultry meat. Among poultry, raising

of ducks is more profitable in Bangladesh. Ducks can be easily brooded, considerably cheaper to raise, do not require any elaborate and expensive housing and equipment, needless care and they are more resistant to diseases and have no cannibalism and agnostic behavior [3,4]. Duck population in Bangladesh has been reported to be 45.12 million [2] mostly of indigenous type although genetic dilution in some regions has occurred due to distribution of high yielding breeds or strains. Ducks in Bangladesh are traditionally reared as family poultry following free range scavenging system. Farmers, who cannot afford to keep large animals because of the big investment required, can easily maintain a few chicken or ducks within their homestead premises [5]. Ducks rank second, next to chicken in the country in terms of total egg and meat production [4]. It has been stated that national share of egg production from commercial and family poultry is almost equal and that of meat production is 60:40 [6] in Bangladesh. Ducks being an important poultry species, can contribute efficiently in increasing egg and meat production in the haors or low laying areas of north-eastern districts. Ducks are natural foragers and thus help to control weed and pests that attack field crops, particularly rice which ultimately helps the farmer [7]. Ducks ranked 2nd place as the supplier of eggs and meat and about 98% of ducks in Bangladesh are traditionally reared under the existing scavenging system [8]; 90 to 95% of which are of Desi (native) type [4]. In Bangladesh, the available duck genotypes are Khaki Campbell, Indian Runner, Jinding, Pekin, Muscovy and Desi [4]. Among these, Pekin is an excellent meat producing duck. But they have poor scavenging ability and high mortality under extensive condition [4]. On the contrary, Desi birds are well adapted to poor management and rural conditions of Bangladesh [9,10,11]. The haor basin located in the north-eastern region of Bangladesh is a wetlands ecosystem considered to be of international ecological importance due to the extensive waterfowl population that uses the basin as its habitat. There are altogether 423

71 small and large haors in Bangladesh [1] comprising an area of about 8000 km² dispersed in the
72 districts of Sunamgonj, Sylhet, Moulvibazar, Hobigonj, Netrokona and Kishoreganj [13].
73 Despite the economic importance of the Haors, people in the region are poorer than any other
74 parts of the country. More than 28% of the total population here lives below the lower poverty
75 line [13]. Development potentials are huge, but need an integrated approach for maximizing the
76 utilization of resources, both human and natural resources [13]. Ducks in the haors area of
77 Bangladesh provides self-employment for landless and small farmers. There is a great
78 potentiality of improving the productivity of ducks in coastal and *haor* (large marshy land) areas
79 through supplementary feeding. Recently, Pervin et al. [14] reported that the growth performance
80 of *desi* ducklings could be improved by supplementation of improved diets under scavenging
81 system of rearing. Duck keeping is an integral part of the rural farming system that provides
82 family income for the small, marginal and landless poor. It was found 78 per cent of egg and 86
83 per cent of poultry meats are produced by the smallholders under scavenging condition. Rearing
84 of ducks gives maximum return with minimum cost. Ducks are efficient converter of agricultural
85 by- products; kitchen wastes, seeds, grains, garden left over, insects, green grasses and all other
86 human refusal that would otherwise waste. Ducks occupy second place in comparison with
87 chicken in producing meat and egg in the country. Ducks are traditionally raised under
88 scavenging [8] by the smallholders in haors and low-lying areas with little or no feed
89 supplementation. Duck production in the Kishoreganj districts of Bangladesh provides self-
90 employment for landless and small farmers. There is a great potentiality of improving the
91 productivity of ducks in low-lying and haors areas through supplementary feeding. Khaki
92 Campbell duck being an important poultry species can contribute efficiently in increasing egg
93 and meat production than chicken in the haors or low lying areas in Kishoreganj districts. No

systematic study has yet been done to assess the potentiality of existing duck rearing system in the haor area. In consideration of these factors, this study was designed to fulfill with the following objectives:

- To study the effects of feed additives on productive and reproductive performance of Khaki Campbell in Kuliarchar Upazilla, Kishoreganj district of Bangladesh.

2. MATERIALS AND METHODS

The study was carried out in Kuliarchar Upazilla, Kishoreganj during the period of August 2014 to May 2015.

2.1 Study Area

The study was carried out in Ramdi, Kuliarchar upazilla, Kishoreganj district. Kuliarchar is located at 24.1542°N 90.9000°E . It has 26143 units of house hold and total area 104.01 km². It is in north east side of Bangladesh [15]. As of the 1991 Bangladesh census, Kuliarchar has a population of 133327. Males constitute 50.61% of the population, and females 49.39%. This Upazila's eighteen up population is 67006. Kuliarchar has an average literacy rate of 21.6% (7+ years), and the national average of 32.4% literate. Kuliarchar has 6 Unions and 1 Proshaba, 46 Mauzas/Mahallas and 131 villages.




Figure A. Map of my study area

2.2 Study time

The study was carried out in Kuliarchar Upazilla, Kishoreganj during the period of August 2014 to May 2015.

2.3 Sampling pattern

The data were collected from farmers directly face to face interview and measurements were taken by own self with the help of a weigh balance. In order to evaluate effects of feed additives on productive and reproductive performance of Khaki Campbell duck the following parameters were well thought-out:

1. Body weight
2. Egg weight
3. Egg production
4. Onset of puberty
5. Fertility (%)
6. Hatchability (%)

131

132 2.4 Materials and facilities used in the study

133

134 A total of 120 female day-old ducklings (Khaki Campbell) were procured from the local farmer
 135 of Kuliarchor Upazilla. The ducklings were randomly distributed into 4 treatment groups that is
 136 group A, group B, group C and group D (untreated control) and each group contained 30 ducks.
 137 Group A, group B, and group C were treated by Digester[®], Vitapoultry[®], Alqurzim[®] respectively
 138 which is marketed by ACI animal health Limited. Digester[®] is a probiotics which enhance the
 139 growth of the animal. Vitapoultry[®] is one of the best vitamin and mineral premix which enhance
 140 the growth rate, egg production as well as meat production of the poultry. Alqurzim[®] helps in
 141 improving the digestibility of the poultry. All ingredients were supplied 2g/kg feed daily. The
 142 individual body weight of ducklings was recorded.

143

144 **Table 1. List of the feed additives used in experiment**

145

Name of the feed additives	Composition	Dose
Digester [®]	Probiotics	2g/kg feed
Vitapoultry [®]	Vitamin, minerals and amino acids	2g/kg feed
Alqurzim [®]	Enzyme	2g/kg feed

146

147 2.5 Experimental programme

148

149 The experiment was carried out for a period of 40 weeks to study the effects of feed additives on
 150 body weight, egg production, egg weight, puberty, fertility, hatchability of Khaki Campbell
 151 ducks. One hundred and twenty female day-old ducklings of Khaki Campbell were divided into

four treatment groups. There were 30 ducklings in each treatment group. During experimental period all Khaki Campbell ducks were provided with a balanced diet and feed additives with abundant fresh water.

2.6 General management

Before the onset of experiment the house was cleaned, disinfected and dried thoroughly. Virocid[®] was used as a disinfectant to kill the organisms at the of the brooder house. Feed and water was provided adlibitum during the brooding period. To facilitate *ad lib* intake, fresh feed and water was provided daily in the morning and afternoon. The left over feed was collected, weighed and dried to know the daily feed intake. Routine management practices were adopted for all treatment groups.

2.7 Statistical Analysis

Mean value (MV) and standard errors (SE) for different traits were estimated with the help of statistical analysis system computer software.



Figure 1. Day old Khaki Campbell ducklings after hatching




Figure 2. Feeding of Khaki Campbell ducks and ducklings with feed additives riched feed.



Figure 3. Feed Additives used in my study



Figure 4. Adult laying duck during feeding in dry season

3. RESULTS

The study was carried out in Kuliarchar upazilla, Kishoreganj during the period of August 2014 to May 2015. A total of 120 female day-old ducklings (Khaki Campbell) were procured from the local farmers of Ramdi, Kuliarchor Upazilla. The ducklings were randomly distributed into 4

treatment groups that is group A, group B, group C and group D (untreated control). Group A, group B, and group C were treated by Digester[®], Vitapoultry[®], Alqurzim[®] respectively which is marketed by ACI animal health Limited. All feed additives mixed with basal feed @ 2g/kg daily.

Table 2. Effects of feed additives on Productive performance of Khaki Campbell Duck

Age (Weeks)	Body weight (g)				Significance
	Group A	Group B	Group C	Group D	
	Digester [®] (2g/kg feed)	Vitapoultry [®] (2g/kg feed)	Alqurzim [®] (2g/kg feed)	Control	
Day old	39.79±0.20	41.39±0.69	40.59±1.57	39.10±0.87	NS
4	230.23±0.21	256.45±0.98	213.43±0.11	196.51±0.26	NS
8	447.40±0.81	481.36±0.51	438.25±0.61	385.30±0.76	S
12	882.86±0.45	921.87±0.60	857.54±0.87	828.66±0.83	S
16	1033.00±0.34	1061.88±0.57	993.98±0.47	917.87±0.67	S
20	1291.23±5.13	1359.11±7.03	1237.80±0.87	1156.27±1.38	S
24	1735.23±5.77	1784.23±2.56	1716.32±2.15	1545.78±1.12	S
40	1584.94±7.47	1616.7±8.72	1547.9±7.96	1406.9±8.82	S
Productive Performance: (Egg Production)					
24-28	19.70±0.64	19.78±0.61	18.76±0.56	18.51±1.25	NS
28-32	22.81±0.67	22.67±0.70	23.56±0.87	21.56±0.67	NS
32-36	18.44±0.45	20.09±0.38	17.91±0.56	17.62±0.26	S
36-40	14.61±0.69	15.83±1.65	15.16±0.46	14.42±0.79	NS
Productive Performance: (Egg Weight)					
24-28	59.65±0.20	60.06±0.71	58.52±0.77	57.89±0.06	NS
28-32	58.07±0.33	58.34±0.55	58.13±0.86	57.96±0.27	NS
32-36	60.76±0.86	62.67±0.09	60.02±1.00	59.86±0.48	S
36-40	66.70±0.24	68.16±0.62	65.46±0.63	64.36±0.91	S

S means Significant, NS means Non Significant

Table 4. Effects of feed additives on Reproductive performance of Khaki Campbell duck

Traits					Significance
	Group A	Group B	Group C	Group D	
	Digester [®] (2g/kg feed)	Vitapoultry [®] (2g/kg feed)	Alqurzim [®] (2g/kg feed)	Control	
On set Puberty (Week)	23.9±0.12	22.54±0.17	23.14±0.57	24.14±0.61	NS
Fertility (%)	83.78±0.57	85.44±2.46	84.28±0.24	78.49±2.74	S

Hatchability (%)	67.27±8.09	68.16±7.68	64.94±0.97	58.85±3.83	S
------------------	------------	------------	------------	------------	---

S means Significant, NS means Non Significant

4. DISCUSSION

The body weight of Khaki Campbell ducks recorded at day one, 4 weeks, 8 weeks, 12 weeks, 16 weeks, 20 weeks, 24 weeks and 40 weeks of age during the experiment (Table 2). Body weight of all groups reduced at 40 weeks of age. This down ponderal weigh in lay period by comparison with breeding period is essentially caused by an intensity of lay. The weight was higher in group B treated by Vitapoultry® in every stage of weighing compare with other. The body of control group was lower than the treated group. There was a significant ($p<0.01$) difference between the treated group and control group. The present findings were almost similar with the Chakravarti and Mohan [16]. They recorded the body weight of Khaki Campbell duck at 8 weeks, 12 weeks, 20 weeks and 72 weeks of age were 448.97 ± 0.81 , 921.86 ± 0.60 , 1359.11 ± 7.03 and 1590.86 ± 1.74 g respectively. They also reported that during lay period ducks have given a weight of 1760 ± 151.6 g and 1715 ± 159.5 g respectively for witness and experimental diets. They found the final body weight and body weight gain were highest in subgroups feed *ad-libitum* with weak significant difference with the group of control. The egg production of Khaki Campbell ducks recorded (Table 2). There was a significant difference ($p<0.01$) between the treated group and control group in 32-36 weeks of age. The feed additives enhance the increasing of egg production in duck. The present findings were almost similar with the findings of Chakravarti and Mohan [16]. The reported that Khaki Campbell had high egg productivity than other breed when supplied feed additives with basal breed. Awad et al. [18] reported that egg number per duck were insignificantly higher for ducks fed diet supplemented with different betaine levels as compared to those fed the control diet. The egg weight of Khaki Campbell ducks recorded (Table 2). Significance difference ($p<0.01$) found in 32-36 weeks and 36-40 weeks of age. Feed

additives directly effects on egg weight of ducks. The treated group Vitapoultry[®] gave highest weight of eggs compared with other. The results was in agreement with those reported by Hetzel [19] who reported that Khaki Campbell ducks laid high weight egg required compared with other breeds. Awad et al. [18] reported that the egg weight of duck increased when supplemented of feed additives with basal feeds. They found egg weight at 24 weeks and 40 weeks of age 59.89±0.06, 58.52±0.77, 59.65±0.20, 60.06±0.71 g and 70.46±0.63, 70.36±0.91, 71.70±0.24, 71.16±0.62 g respectively. The onset of puberty of Khaki Campbell recorded at group A, group B, group C and group D were 23.9±0.12, 22.54±0.17, 23.14±0.57, 24.14±0.61 weeks respectively. The fertility of egg (%) of Khaki Campbell recorded at group A, group B, group C and group D were 83.78±0.57, 85.44±2.46, 84.28±0.24 and 78.49±2.74 respectively (Table 4). The fertility (%) of egg were significantly different ($p<0.01$) between the groups. The findings were almost similar with Awad et al. [18]. He found fertility rate (%) 87.78±0. Fertility percentage was insignificantly improved by supplementing 0.5 and 1.0 g B/kg diet. The hatchability (%) highest in treated groups compared with the control group. The present study similar with the findings of Awad et al. [18]. He recorded the hatchability (%) was 63.61±7.68. He reported that hatchability percentage fertile eggs were significantly improved by supplementing 0.5 g B/kg diet as compared to the control.

5. CONCLUSION AND RECOMMENDATION

A total of 120 female day-old ducklings (Khaki Campbell) were procured from the local farmers of Ramdi, Kuliarchor Upazilla. The ducklings were randomly distributed into 4 treatment groups that is group A, group B, group C and group D (untreated control). Group A, group B, and group C were treated by Digester[®], Vitapoultry[®], Alqurzim[®] respectively which is marketed by ACI animal health Limited. All feed additives mixed with basal feed @ 2g/kg daily. Feed additives

had tremendous result both productive and reproductive performance of Khaki Campbell duck compared with control group in Kuliarchar Upazilla, Kishoreganj district. Specifically Vitapoultry[®] had very good result when mixed with the basal feed and provided daily to the duck. So, it is recommended that use of vitamin, mineral and amino acid containing feed additives with basal feed might be helpful for enhancing the productive and reproductive performance of Khaki Campbell ducks in Bangladesh. Further studies need to develop an appropriate implementation of using feed additives with basal feed in ducks in Bangladesh.

6. REFERENCES

1. BER. Bangladesh Economic Review, 2012. Ministry of Finance, Government of the Peoples' Republic of Bangladesh, Dhaka.
2. Islam S. An Economic Study on Different Farming Systems in Dingaputa Haor Area of Netrokona District, 2012. M.S. Thesis, Department of Agricultural Economics, Bangladesh Agricultural University, Mymensingh.
3. Batty J. Domesticated ducks and geese. *Spur Publications*, 1979. Saiga Publishing Co. Ltd. England, 116 and 143.
4. Ahmed S. Duck production in Bangladesh. In "Duck Production Science and World Practice". Farrell, D. J. and P. Stapleton, (Ed.). *University of New England, Armidale, Australia*, 1986; 342-350.

- 264 5. Das SC, Chowdhury SD, Khatun MA, Nishibori M, Isobe N and Yoshimura Y. Poultry
265 production profile and expected future projection in Bangladesh. *World's Poultry*
266 *Science Journal*, 2008; 64:99-116.
- 267 6. Bhuiyan AKFH. Implementation of National Livestock Development Policy (2007) and
268 National Poultry Development Policy (2008): Impact on smallholder livestock rearers.
269 Keynote paper presented at the South Asia Pro Poor Livestock Policy Programme
270 (SAPPLP)-BRAC workshop held at BRAC Centre, Dhaka, 2011.
- 271 7. Manda M, Uchida H, Nakagama A, Matsumoto S, Shimoshikiryo K and Watanable S. Effects
272 of Aigamo ducks (crossbred of wild and domestic ducks) herding on weeding and pest
273 control in the paddy fields. *Japanese Poultry Science*, 1993; 30: 365-370.
- 274 8. Salahuddin M, Barua A, Rashid N and Howlider MAR. A study on the relationship of egg
275 weight with fertility and hatchability of Desi ducks. *Poultry Guide*, 1991; 28(3): 34-37.
- 276 9. Khan AG. Improvement of Desi birds-part 2. *Poultry Adviser*, 1983; 16: 67-78.
- 277 10. Hussain MS, Abdin MA, Quddus MA, Hossain AMM, Banu T, Ara S and Ahmed D.
278 Women's contribution to the homestead agricultural production system in Bangladesh
279 Academy for Rural Development, 1988, Kotbari, Comilla.
- 280 11. DLS. Livestock Population, 1990. Directorate of Livestock Services. Farmgate, Dhaka.
- 281 12. Alam MS, Quayu MA and Isla MA. Crop Production in the Haor Areas of Bangladesh:
282 Insights from Farm Level Survey. A Scientific Journal of Krishi Foundation, 2011, 8(2),

- 283 Agricultural Economics Division, *Bangladesh Rice Research Institute*, Gazipur- 1701,
284 Bangladesh: 88-97.
- 285 13. BHWDB. Bangladesh Haor and Wetland Development Board, 2011. Ministry of Water
286 Resources, Government of the People's Republic of Bangladesh, Dhaka.
- 287 14. Pervin W, Chowdhury SD, Ali MA, Khan JU and Raha SK. Growth performance of
288 indigenous (*desi*) ducklings receiving diets of varying nutrient concentrations. In:
289 Proceedings of 8th Poultry Show and Seminar, *World's Poultry Science Association*,
290 Bangladesh Branch, 2013; 45-50.
- 291 15. BBS. Population Census Wing 2006.
- 292 16. Chakravathi PV and Mohan B. Comparison of performance of Khaki Campbell Ducks under
293 organized farm and field conditions, *Shanlan International Journal of Veterinary*
294 *Science*, 2014; 2(2): 1-3.
- 295 17. Fulbert AC, Ngokaka and Mompoundza P. Productivity of Khaki Campbell Ducks (*Cairina*
296 *moschata*) under the Local Agricultural Produce and by Products in the Breeding
297 Conditions of Brazzaville-Congo, *Pakistan Journal of Nutrition*, 2008; 8 (12): 1910-
298 1914.
- 299 18. Awad AL, Fahim HN, Ibrahim AF and Beshara MM. Effect of dietary betaine
300 supplementation on productive and reproductive performance of domyati ducks under
301 summer conditions. Animal Production Research Institute, Agricultural Research Center,
302 Ministry of Agriultural, *Dokki, Giza. Egypt. Poultry Science*, 2014; 34 (2): 453-474.

- 303 19. Hetzel DJS. The growth and carcass characteristics of crosses between Alabio and Tegal
304 ducks and Muscovy and Pekin drakes. *British Poultry Science*, 1983; 24: 555-563.

305