

Integrated management of *Striga hermonthica* on Sorghum

Abstract

The phytoparasite, *Striga hermonthica* (Del.) Benth., is a major biotic constraint to cereals production in sub-Saharan Africa. The experiment was designed to investigate the effects of nitrogen and the herbicides triclopyr, chlorsulfuron and their combination on *Striga* incidence and sorghum (cv. Wad-Ahmed) growth and yield. Treatments were arranged in Complete Randomized Block Design (CRBD). All treatments at 45 days after sowing (DAS), reduced *Striga* emergence albeit not significantly as compared to the control. At 60 DAS, untreated control (3 plants/ m²) sustained the highest *Striga* emergence significantly as compared to other treatments. Sorghum height, irrespective of date of observation did not show significant differences between treatments. At 60 DAS, triclopyr at 0.49 and 0.73 kg a.e ha⁻¹ supplemented with nitrogen at 43.8 kg ha⁻¹, increased sorghum height by 11.9 and 9.4%, respectively, as compared to untreated control. Chlorsulfuron at 2.38 g a.i ha⁻¹ when supplemented with nitrogen and irrespective of rate, caused considerable decreased in peduncle length by 46.2-35.7%, respectively when compared to the control. However, triclopyr at 0.73 kg a.e. ha⁻¹ alone or in combination with nitrogen at 87.6 kg ha⁻¹ had no effects on peduncle length when compared to the untreated control. All treatments significantly increased head length over the untreated control. All treatments increased head length over the untreated control albeit not significant. Chlorsulfuron at the highest dose when combined with nitrogen at the highest rate resulted in the highest increase in head length 21.7%. Triclopyr at 0.49 and 0.73 kg a.e. ha⁻¹, in supplementation with nitrogen at 87.6 kg ha⁻¹ displayed the highest increase in straw weight 23.2-28.4%, respectively. Triclopyr at 0.49 kg a.e. ha⁻¹, tank mixed with chlorsulfuron at 1.19 g a.i ha⁻¹ and in supplemented with nitrogen at 87.6 kg ha⁻¹ caused straw weight increase by 5.4-8.6% over the untreated control.

Keywords: Nitrogen, chlorsulfuron, triclopyr, *Striga*, sorghum, emergence

Introduction

The phytoparasite, *Striga hermonthica* (Del.) Benth., infests millions of hectares of cultivated fields of cereal crops in sub-Saharan Africa. Yield losses are often 70 but can be as high as 100% and farmers often abandon infested fields and move to new areas (Kim, 1995). It is an obligate parasitic plant that attaches to host roots to deplete them of nutrients. In Sub-Saharan Africa, the most destructive *Striga* species, *Striga hermonthica*, parasitizes major food crops affecting two-thirds of the arable land and over 100 million people. One potential weakness in the *Striga* infection process is the way it senses the presence of a host crop (Smith, 2016).

Striga hermonthica is a widespread parasitic weed in Africa, causing severe damages to the major cereal crops: millet, sorghum and maize. Considering the subsistence nature of farming in Africa, the development of resistant or tolerant varieties is the most promising cost-effective approach to overcoming the weed problem (Efron, 1993).

Witchweeds (*Striga* spp.) are noxious parasitic weeds that cause considerable crop damage in the semiarid tropics (Rich, *et al* 2003).

The witchweed *Striga hermonthica* is proving to be a nightmare for smallholder farmers in the Sahel and Savannah regions of Africa. The close and intricate biological association between this phytoparasite and the cereal host makes its management difficult (Lendzemo, 2004). Prodigious seed production, prolonged viability and ease of dissemination of the seeds together with the subterranean nature of the early stages of parasitism make *Striga* a difficult to control invasive weed (Babiker 2007).

Herbicides are considered to have the most potential, however, they have to be appropriate, cost-effective and affordable (Hess and Lenne 1999). Damages by chemical persistence and availability are other major constraints that limit the successful usage of herbicides for parasitic weed control. In addition, in developing countries, the income of subsistence farmers is usually too low to afford to purchase inputs including herbicides (Aly 2007).

Parasitic weeds control, has proved to be extremely difficult. Unusual seed production, prolonged viability and ease of distribution of the seeds together with the subterranean nature of the early stages of parasitism make *Striga* a difficult weed to control (Babiker 2007). However, several control methods have been investigated positively, but often inconsistent results across seasons and sites. These methods include biological, cultural and chemical methods and they are well reviewed by Parker and Riches, (1993) and summed up by (Kroschel 2001; Elzein and

63 Kroschel 2003; Babiker 2007; Joel, et al. 2007 and Aly 2007).

64 **Materials and Methods**

65 A field experiment was undertaken at the College of Agricultural Studies, (CAS), at Shambat
66 during the season 2011/2012. The experiment was designed to investigate the effects of nitrogen
67 and the herbicides triclopyr, chlorsulfuron and their combination on *Striga* incidence and
68 sorghum (cv. Wad-Ahmed) growth and yield. Treatments were arranged in Complete
69 Randomized Block Design (CRBD).

70 The experimental area was disc ploughed, harrowed, levelled, ridged and divided into sub-plots.
71 Sorghum (Wad Ahmed) seeds, supplied by the Arab Seed Production Company, were treated
72 with the fungicide Thiram. Seeds were sown in holes at a within row spacing of 20 cm. In
73 experiments involving fertilization, nitrogen in form of urea was applied as broadcast by hand.
74 All sub-plots were artificially infested with *Striga hermonthica* (Del.) Benth seeds. *Striga* seeds
75 used in this study were collected from under sorghum in Gazira, Sudan. Inoculums were
76 prepared by through mixing of 1 g of clean *Striga* seeds with 1 kg of soil. Inoculums were
77 applied to the soil at time of sowing. The crop was sown in July. Sorghum seedlings were
78 thinned to two per hole 15 days after emergence. The herbicide triclopyr as Garlon was applied
79 30 DAS by knapsack sprayer as aqueous spray. Weeds were removed by hand to avoid
80 damaging emerged *Striga* plants at biweekly intervals for the first six weeks. Irrigation was
81 applied when necessary throughout the duration of the trial.

82 Treatment effects, were assessed by determining i) number of emerged *Striga* plants per meter
83 square 45, 60 and 90 days after sowing (DAS), ii) peduncle length in cm, iii) diameter at flag
84 leaf in cm, iv) threshing percentage, v) *Striga* dry weight in g.

85 **Effects of nitrogen, triclopyr, chlorsulfuron and their combinations on *Striga* incidence** 86 **and sorghum growth and yield**

87 Nitrogen at 43.8 and 87.6 kg ha⁻¹ was applied by broadcasting. Triclopyr at 0.49 and 0.73 kg
88 a.e. ha⁻¹ implemented alone. Chlorsulfuron 2.38 g a.i ha⁻¹ implemented alone. The combinations
89 triclopyr at 0.49 a.e ha⁻¹ and urea at 43.8 and 87.6 kg ha⁻¹. The combinations triclopyr at 0.73 kg
90 a.e. ha⁻¹ and urea at 43.8 and 87.6 kg ha⁻¹. The combinations triclopyr at 0.49 a.e ha⁻¹,
91 Chlorsulfuron at 1.19 and urea at 43.8 and 87.6 kg ha⁻¹ were applied 30 days after sowing.

92 **Statistical analysis**

93 Data collected from the experiment was subjected to statistical analysis using GenStat

(PC/Windows 7), VSN International Ltd., UK statistical package (Rothamsted Experimental Station), Means were separated for significance using Least Significant Difference (LSD).

Results

Striga

Striga emergence

At 45 days after sowing (DAS), *Striga* emergence in the untreated control displayed 2 plants/m². All treatments reduced *Striga* emergence albeit not significantly as compared to the control. At 60 DAS, untreated control (3 plants/m²) sustained the highest *Striga* emergence significantly as compared to other treatments (Table 1).

Table 1. Effects of chlorsulfuron, nitrogen and their combinations on *Striga* emergence

Treatment	Herbicide rate ha ⁻¹	<i>Striga</i> emergence (plants/m ²) (DAS)	
		45	60
Untreated control	-	2(2)	2(3) ^a
Chlor. ⁱ	2.38	1(0)	1(0) ^b
Chlor.+ 1N	2.38	1(1)	1(0) ^b
Chlor.+ 2N	2.38	1(0)	1(0) ^b
Tr. ⁱⁱ	0.49	1(0)	1(0) ^b
Tr. + 1N	0.49	1(0)	1(0) ^b
Tr. + 2N	0.49	1(0)	1(0) ^b
Tr. + Chlor.	0.49+1.19	1(1)	1(1) ^b
Tr. + Chlor.+1N	0.49+1.19	1(0)	1(0) ^b
Tr. + Chlor.+2N	0.49+1.19	1(0)	1(0) ^b
Tr.	0.73	1(0)	1(0) ^b
Tr. + 1N	0.73	1(1)	1(0) ^b
Tr. + 2N	0.73	1(0)	1(0) ^b
P ≤ 0.05	-	NS	0.001***
CV%	-	26.1	20.7

Means within a column having the same superscript letter(s) are not significantly different according to LSD test. Bracketed data are real. NS = Non significant. ***= P ≤ 0.001. ⁱ = Chlorsulfuron/ g a.i. ha⁻¹. ⁱⁱ = Triclopy/kg a.e ha⁻¹, 1N= Nitrogen at 43.8 kg ha⁻¹ 2N= Nitrogen at 87.6 kg ha⁻¹.

Plant height

Sorghum height, irrespective of rate of observation did not show significant differences between treatments (Table 2). At 45 DAS, combination between triclopyr at 0.49 kg a.e.ha⁻¹ and nitrogen at 43.8 kg ha⁻¹ displayed the highest plant height, while chlorsulfuron alone at 2.38 g a.i ha⁻¹ displayed the lowest (Table 2). At 60 DAS, triclopyr at 0.49 and 0.73 kg a.e ha⁻¹ supplemented with nitrogen at 43.8 kg ha⁻¹, increased sorghum height by 11.9 and 4.9%, respectively, as compared to untreated control. Chlorsulfuron at 2.38 g a.i ha⁻¹ in combination with nitrogen at 87.6 kg ha⁻¹ and triclopyr alone at 0.49 kg a.e. ha⁻¹ reduced sorghum height by 7.7-8.5%, but not significantly, as compared to the control (Table 2).

Table 2. Effects of chlorsulfuron, nitrogen and their combinations on sorghum height

Treatment	Herbicide rate ha ⁻¹	Sorghum height (cm) (DAS)	
		45	60
Untreated control	-	77.6	107.5
Chlor. ⁱ	2.38	56.3	102.7
Chlor.+1N	2.38	77.1	107.1
Chlor.+ 2N	2.38	67.7	98.4
Tr. ⁱⁱ	0.49	67.4	99.2
Tr. + 1N	0.49	83.3	120.3
Tr. + 2N	0.49	79.0	110.5
Tr. + Chlor.	0.49+1.1	74.4	111.6
Tr. + Chlor.+1N	0.49+1.1	79.5	113.6
Tr. + Chlor.+2N	0.49+1.1	68.3	109.0
Tr.	0.73	67.2	107.2
Tr. + 1N	0.73	74.9	117.6
Tr. + 2N	0.73	67.1	108.0
P ≤ 0.05		NS	NS
CV%	-	17.1	10.5

NS= non- significant. ⁱ = Chlorsulfuron/ g a.i. ha⁻¹. ⁱⁱ = Triclopy/kg,
1N= Nitrogen at 43.8 kg ha⁻¹ 2N= Nitrogen at 87.6 kg ha⁻¹.

Peduncle length

Chlorsulfuron at 2.38 g a.i ha⁻¹ alone and when supplemented with nitrogen irrespective of rate, caused considerable decrease in peduncle length by 41.0-51.3% when compared to the control (Table 3). Triclopyr at 0.49 kg a.e. ha⁻¹ alone and when added subsequent to nitrogen at 43.8 kg ha⁻¹ decreased peduncle length by 15.4%. Triclopyr at 0.73 kg a.e. ha⁻¹ when combined with nitrogen irrespective of rate increased peduncle length by 5.1-25.6%. However, triclopyr rate at the same rate when tank mixed with chlorsulfuron at 1.19 g a.i ha⁻¹ and nitrogen at 43.8 kg ha⁻¹ increased peduncle length by 17.9% (Table 3).

Diameter at flag

In general, all treatments had no significant effects on diameter at flag leaf when compared to the untreated control (Table 3). Chlorsulfuron at 2.38 g a.i ha⁻¹ alone and when supplemented with nitrogen irrespective of rate, caused an increase in diameter at flag leaf by 5.0-6.6%. Triclopyr at 0.49 kg a.e. ha⁻¹ alone and when combined with nitrogen at 87.6 kg ha⁻¹ displayed an increase in diameter at flag leaf by 5.2-7.0%. Triclopyr at the lower dose when combined with nitrogen at the lower rate and Triclopyr at the higher dose in combination with nitrogen at the lower rate resulted in an increase of 2.5-.8%, respectively (Table 3).

Table 3. Effects of chlorsulfuron, nitrogen and their combinations on sorghum exertion

Treatment	Herbicide rate ha ⁻¹	Head Attributes	
		Peduncle length /cm	Diameter at flag Leaf/cm
Untreated control	-	1.9(3.9)	5.15
Chlor. ⁱ	2.38	1.5(2.3)	4.70
Chlor.+ 1N	2.38	1.3(1.9)	5.41
Chlor.+ 2N	2.38	1.4(2.0)	5.49
Tr. ⁱⁱ	0.49	1.8(3.3)	5.51
Tr. + 1N	0.49	1.8(3.3)	4.41
Tr. + 2N	0.49	1.6(2.7)	5.42

Tr. + Chlor.	0.49+1.19	1.7(3.1)	5.14
Tr. + Chlor.+1N	0.49+1.19	2.1(4.6)	5.28
Tr. + Chlor.+2N	0.49+1.19	1.5(3.0)	5.33
Tr.	0.73	1.9(3.8)	4.92
Tr. + 1N	0.73	2.2(4.9)	5.19
Tr. + 2N	0.73	1.9(4.1)	4.33
P ≤ 0.05	-	NS	NS
CV%	-	31.2	17.7

Bracketed data are real. NS = Non significant. ***= $P \leq 0.001$. i = Chlorsulfuron/ g a.i. ha^{-1} . $i i$ = Triclopyr/kg a.e ha^{-1} , 1N= Nitrogen at 43.8 kg ha^{-1} 2N= Nitrogen at 87.6 kg ha^{-1} .

Head length

All treatments significantly increased head length over the untreated control. Chlorsulfuron at the highest dose when combined with nitrogen at the highest rate resulted in the highest increase in head length 27.6% (Table 4). Chlorsulfuron at 2.38 g a.i ha^{-1} when supplemented with nitrogen at the lower rate, and triclopyr at 0.73 kg a.e. ha^{-1} when supplemented with nitrogen at the lower rate, displayed head length increase by 18.7%. Triclopyr at 0.49 kg a.e. ha^{-1} , alone and when supplemented with nitrogen irrespective of nitrogen rate, caused an increase in sorghum head length by 8.1-15.4%. However, triclopyr at 0.73 kg a.e. ha^{-1} alone or in combination with nitrogen at 87.6 kg ha^{-1} displayed the least sorghum head length of 6.5% (Table 4).

Head circumference

All treatments increased head circumference over the untreated control albeit not significant (Table 4). Chlorsulfuron at 2.3 8 g a.i ha^{-1} in supplementation with nitrogen at 87.6 kg ha^{-1} and triclopyr at 0.49 kg a.e. ha^{-1} , tank mixed with chlorsulfuron at 1.19 g a.i ha^{-1} in supplementation with nitrogen at 43.8 kg ha^{-1} displayed the heist increase in head circumference 31.3%. Chlorsulfuron at 2.3 8 g a.i ha^{-1} in supplementation with nitrogen at 43.8 kg ha^{-1} gave an increase in head circumference by 28.9%. Triclopyr at the lower rate alone and triclopyr at the highest dose when applied subsequent to nitrogen at 87.6 kg ha^{-1} showed an increase in head circumference by 18.1% (Table 4).

Triclopyr at 0.49 kg a.e. ha^{-1} alone, or when added subsequent to nitrogen irrespective of rate, displayed an increase in sorghum head circumference by 8.4-21.7%. However, triclopyr at 0.73

174 kg a.e. ha⁻¹ alone or in combination with nitrogen at 43.8 kg ha⁻¹ caused the lowest head
175 circumference effect 8.4-26.5%, respectively (Table 4).

176 **Table4. Effects of chlorsulfuron, triclopyr, nitrogen and their combinations on sorghum**
177 **head length and circumference**

Treatment	Herbicide rate ha ⁻¹	Head Attributes	
		Length/cm	Circumference/cm
Untreated control	-	12.3	8.3
Chlor. ⁱ	2.38	13.2 ^{bcd}	9.4
Chlor.+ 1N	2.38	14.6 ^b	10.7
Chlor.+ 2N	2.38	15.7 ^a	10.9
Tr. ⁱⁱ	0.49	14.2 ^{bc}	9.8
Tr. + 1N	0.49	13.3 ^{bcd}	10.1
Tr. + 2N	0.49	13.4 ^{bcd}	9.0
Tr. + Chlor.	0.49+1.19	13.8 ^{bc}	9.8
Tr. + Chlor.+1N	0.49+1.19	13.6 ^{bcd}	10.9
Tr. + Chlor.+2N	0.49+1.19	14.5 ^{ab}	9.5
Tr.	0.73	13.1 ^{cd}	9.0
Tr. + 1N	0.73	14.6 ^{ab}	10.5
Tr. + 2N	0.73	13.1 ^{cd}	9.8
P ≤ 0.05	-	0.003**	NS
CV%	-	7.1	13.6

178 Means within a column having the same superscript letter(s) are not significantly different according to
179 LSD test. NS = Non significant. **= P ≤ 0.01. ⁱ = Chlorsulfuron/ g a.i. ha⁻¹. ⁱⁱ = Triclopyr/kg a.e ha⁻¹,
180 1N= Nitrogen at 43.8 kg ha⁻¹ 2N= Nitrogen at 87.6 kg ha⁻¹.

182 **Straw weight**

183 Triclopyr at 0.49 and 0.73 kg a.e. ha⁻¹, in supplementation with nitrogen at 87.6 kg ha⁻¹
184 ¹Displayed the highest increase in straw weight 23.2-28.4%, respectively when compared to the
185 control (Table 5). Triclopyr at 0.49 kg a.e. ha⁻¹ and 0.73 kg a.e. ha⁻¹, in supplementation with
186 nitrogen at 43.8 kg ha⁻¹ displayed 15.2-39.2% decrease in straw weight. Chlorsulfuron at 2.38 g
187 a.i ha⁻¹ alone or in supplementation with nitrogen irrespective of rate displayed an increase in
188 sorghum straw weight by 3.6, 7.0 - 8.6%, respectively. Triclopyr at 0.49 kg a.e. ha⁻¹, tank mixed

189 with chlorsulfuron at 1.19 g a.i ha⁻¹ and in supplemented with nitrogen at 87.6 kg ha⁻¹ caused
190 straw weight increase by 5.4-8.6% over the untreated control (Table 5).

191
192
193
194

Table 5. Effects of chlorsulfuron, nitrogen and their combinations on sorghum straw dry weight

Treatment	Herbicide rate ha ⁻¹	Straw weight g/m ²
Untreated control	-	53
Chlor. ⁱ	2.38	55
Chlor.+ 1N	2.38	58
Chlor.+ 2N	2.38	57
Tr. ⁱⁱ	0.49	38
Tr. + 1N	0.49	56
Tr. + 2N	0.49	69
Tr. + Chlor.	0.49+1.19	58
Tr. + Chlor.+1N	0.49+1.19	48
Tr. + Chlor.+2N	0.49+1.19	56
Tr.	0.73	52
Tr. + 1N	0.73	46
Tr. + 2N	0.73	74
P ≤ 0.05	-	NS
CV%	-	28.9

195 NS = Non significant. ⁱ = Chlorsulfuron/ g a.i. ha⁻¹. ⁱⁱ = Triclopyr/kg a.e ha⁻¹, 1N= Nitrogen at 43.8 kg
196 ha⁻¹ 2N= Nitrogen at 87.6 kg ha⁻¹.
197

198 Discussion

199

200 The herbicides triclopyr and chlorsulfuron irrespective of doses, effected sizeable to significant
201 decrease in *Striga hermonthica* infestation. In general the herbicides triclopyr and chlorsulfuron
202 when applied subsequent to nitrogen as urea, at 43.8 (1N) and 87.6 (2N) kg ha⁻¹, or combined
203 with their tank mixtures were more exploitive to the parasite than when each was applied alone.
204 At 45 DAS *Striga* emergence was inconsistent and maximum emergence was 1 plant m⁻². At 60
205 DAS the parasite displayed maximum emergence (3 plants m⁻²) in the untreated control sub-

plots, followed by sub-plot treated with triclopyr at 0.49 kg a.e. ha⁻¹ tank mixed with chlorsulfuron at 1.19 g a.i. ha⁻¹.

Chlorsulfuron is acetolactate synthase (ALS) inhibitor (Forlani, et al. 1991; Dastgheib and Field 1998). The herbicide inhibits synthesis of the branched amino acids L-leucine, L- isoleucine and L-valine, and thus may interfere with protein synthesis and cell division (Ray 1984). These findings indicate clearly that chlorsulfuron alone delayed *Striga* emergence early in the season these findings are in agreement with those previously reported by Babiker and Reda 1991; Babiker, et al. 1993; Babiker 2002; Ahonsi, et al. 2004 and Adam 2007.

Chlorsulfuron at 2.38 g a.i ha⁻¹ alone and when supplemented with nitrogen irrespective of rate, caused considerable decrease in peduncle length by 41.0-51.3% when compared to the control (Table 3). Triclopyr at 0.49 kg a.e. ha⁻¹ alone and when added subsequent to nitrogen at 43.8 kg ha⁻¹ decreased peduncle length by 15.4%. In general, all treatments had no significant effects on diameter at flag leaf when compared to the untreated control. All treatments significantly increased head length over the untreated control. Chlorsulfuron at the highest dose when combined with nitrogen at the highest rate resulted in the highest increase in head length 27.6% (Table 4). All treatments increased head circumference over the untreated control albeit not significant.

The general trend in growth of the chlorsulfuron treated sorghum compared to the untreated Sorghum height, irrespective of rate of observation did not show significant differences between treatments (Table 2). At 60 DAS triclopyr, alone, at the two rates (0.49 and 0.73 kg a.e.ha⁻¹) and when applied subsequent to nitrogen irrespective of rate suppressed *Striga* emergence completely.

These results, which are in line with those obtained with chlorsulfron (Tables 4.5 and 4.14), and those previously reported by Last (1960a) and Dafaallah, et al. (2008) for 2,4-D and Last (1960b) for nitrogen, indicate clearly the consistency in performance of the combinations of the herbicides and nitrogen. Furthermore, the results indicate that, in addition to improved performance, nitrogen allows the use of low herbicide rates. Low herbicide rates in addition to economic benefits, safeguard against phytotoxicity and prolonged persistence which may be expected from high rates and/or variations in edaphic and environmental conditions.

CONCLUSIONS

- *Striga* is a difficult weed to control.
- *Striga* management requires integrated practices comprising different components.
- Sorghum (cv. Wad Ahmed), *Striga* tolerant, the herbicides triclopyr or chlorsulfuron provide efficient packages for *Striga* management.
- Sorghum (cv. Wad Ahmed), *Striga* tolerant, the herbicides triclopyr or chlorsulfuron when applied subsequent to nitrogen or when tank mixed with each other provide efficient packages for *Striga* management.

References:

1. Adam, A.G. 2007. Effects of Urea and the herbicides oxyfluorfen and chlorsulfuron/2,4-D tank mixtures on control of *Striga hermonthica* and sorghum growth and yield. M.Sc. Thesis, University of Gezira. pp 60.
2. Ahonsi, M.O. Bernera, D.K. Emechebeb, A.M. and Lagoke, S.T. 2002. Effects of soil pasteurisation and soil N status on severity of *Striga hermonthica* (Del.) Benth. In: maize International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria Institute for Agricultural Research, Samaru, PMB 1044, Ahmadu Bello University, Zaria, Nigeria Soil Biology and Biochemistry 34:1675 -1681.
3. Aly, R. 2007. Conventional and biotechnological approaches for control of parasitic weeds. The Society for *in-vitro* biology. 43:304-317.
4. Assefa, Y. Staggenborg, S.A and Prasad, P.V. 2010. Grain sorghum water requirement and responses to drought stress: A Review. Plant management network. <http://www.plantmanagementnetwork.org> (Accessed January 6, 2014 at 9:16 am).
5. Awika, J.M. and Rooney, L.W. 2004. Sorghum phytochemicals and their potential impact on human health. Phytochemistry. 65:1199–1221.
6. Babiker, A.G.T. 2007. Striga: The spreading scourge in Africa. Regulation of plant Growth and Development 42:74-87.

- 265 7. Babiker, A.G.T. 2002. *Striga* Control in Sudan: An Integrated Approach. In: Leslie, J. F.
266 (Ed.). Sorghum and millets diseases. Ames, Iowa State Press: Iowa. pp 159-163.
- 267 8. Babiker, A.G.T., Ahmed, N.E., Mohamed, A.H., Eltayeb, S.M., El Mana, M.E.1993.
268 *Striga hermonthica* on sorghum. IN: Chemical and cultural control. Brighton crop
269 protection conference-weeds. 907-911.
- 270 9. Babiker, A.G.T. and Reda, F. 1991. Recent advance in *Striga* research in Sudan and
271 Ethiopia. In: (Ransome, J.K., Musselman, L. J., Worsham, A.D. and Parker, C. eds.).
272 Proceedings of 5th International Symposium on Parasitic Weed. Nairobi, Kenya. pp.180-
273 189.
- 274 10. Dastgheib, F. and Field, R. J. 1998. Acetolactate synthase activity and chlorsulfuron
275 sensitivity of wheat cultivars. Weed Research. 38:63-68.
- 276 11. Elzein, A. and Kroschel, J. 2003. Progress on management of parasitic weeds. In:
277 Labrada (Ed.). Weed Management in Developing Countries. FAO Plant Production and
278 Protection Addendum I, Rome pp 109-143.
- 279 12. E fron,Y. (1993) Screening Maize for Tolerance to *Striga hermonthica*. Plant Breeding,
280 110: 192–200. <http://onlinelibrary.wiley.com/doi/10.1111/j.1439>.
- 281 13. FAO, 2008. Latest forecast for world production of coarse grains. In Food outlook:
282 Global market analysis. <http://www.fao.org/> (Accessed: January 6, 2014 at 6.38 am).
- 283 14. Forlani, G. Nielsen, E. Landi, P. and Tuberosa, R. 1991. Chlorsulfuron tolerance and
284 acetolactate synthase activity in Corn (*Zea mays* L.). Inbred Lines. Weed Science
285 Society of America. 39:553-557.
- 286 15. Hess, D.E. and Lenne, J.M. 1999. Importance of *Striga* as a constraint to sorghum and millet
287 production. Report on the ICRISAT sector review for *Striga* control in sorghum and millet.
288 Bamako, Mali. pp 4-8.
- 289 16. Joel, D.M, Hershenhorn, J. Eizenburg, H. Aly, R. Ejeta, G. Rich, P.J. Ransom, J.K.
290 Sauerborn, J. and Rubiales, D. 2007. Biology and management of weedy root parasites.
291 Horticultural Reviews. Wiley, London, pp 267-349.

17. Kim, S. K. Adetimirin, V. O. and Akintunde, A. Y. (1995) Nitrogen Effects on *Striga hermonthica* Infestation, Grain Yield, and Agronomic Traits of Tolerant and Susceptible Maize Hybrids. Crop Science, 37: 711-716.
<https://dl.sciencesocieties.org/publications/cs/abstracts>
18. Kroschel, J. 2001. Atechnical manual for parasitic weed research and extension. Kluwer Academic Publishers, Dordrecht, The Netherlands. pp 256.
19. Lenzemo, V.W., 2004. The tripartite interaction between sorghum, *Striga hermonthica*, and arbuscular mycorrhizal fungi. PhD thesis, Wageningen University, Wageningen, The Netherlands, 112 pp. with English, French and Dutch summaries.
20. Parker, C. and Riches, C.R. 1993. Parasitic Weeds of the World: Biology and Control. CAB International, Wallingford, Oxon, UK. pp 332.
21. Ray, T.B. 1984. Site of Action of chlorsulfuron. Inhibition of valine and isoleucine biosynthesis in plants. Plant physiology. 75: 827-831.
22. Rich, P. J. Grenier, C. and Ejeta, G (2003) *Striga* Resistance in the Wild Relatives of Sorghum, 44: 2221-2229.
23. Smith, D. H. Toh, S. Tsuchiya, Y and McCourt, Peter (2016) Small-molecule antagonists of germination of the parasitic plant *Striga hermonthica*. Nature Chemical Biology, 12: 724–729. <http://www.nature.com/nchembio/journal/v12/n9/full/>
24. Last, F.T. 1960a. Incidence of *Striga hermonthica* (Del.) Benth. on two varieties of irrigated sorghum differently manured, spaced and thinned. Tropical Agriculture, Trinidad and Tobago. 37: 309-319.
25. Last, F.T. 1960b. Effect of cultural treatments on the incidence of *Striga hermonthica* (Del.) Benth. and yields of sorghum in the Sudan: field experiments. Annals of Applied Biology. 48: 207-229.
26. Vasey, R. A., Scholes, J. D., and Press, M. C. 2005. Wheat (*Triticum aestivum*) is susceptible to the parasitic angiosperm *Striga hermonthica*, a major cereal pathogen in Africa. Phytopathology 95:1294-1300.