

Technical sheet of Process of traditional cassava inoculated for attieke productions in south of Côte d'Ivoire.

Abstract

Production of *attieke* in Cote d'Ivoire requires the use of inputs such as the cassava traditional inocula. Our study is a technical sheet of a process of traditional cassava inocula used for *attieke* production in south of cote d'Ivoire. The process of traditional cassava inocula for *attieke* production was: Cassava roots, peeling, pulp washing and cutting, cooking, cooling, packaging, fermentation. It contained aerobic mesophiles, total coliforms, fecal, *Bacillus*, *Staphylococcus* and moulds.

Keywords : *attieke*, bacteria, cassava traditional inocula, process

1.INTRODUCTION

Attieke is the major fermented plant food in Côte d'Ivoire. It is a steamed granular cassava (*Manihotesculenta*Crantz) meal ready to eat, couscou-like product, with slightly sour taste and whitish colour (Djeni et al., 2011). It is consumed two to three times a day with meat, fish or vegetables. The popularity of attieketo urban dwellers in recent years has been associated with its cheapness, lower bulk (as compared to other cassava product) and its characteristic of ready to eat food. Attieke is one of the few products whose fermentation is not spontaneous but involves the use of an inoculum. This inoculum is obtained after 2-3 days of spontaneous fermentation of cassava roots, thus colonized by a wide variety of microorganism which constitutes the main source of microbial activities during the cassava dough fermentation (Djeni et al., 2008).Thesemicroorganismsconstitute the bulk of the inoculum whichwouldparticipate in the fermentation of the cassavapulp for the production of the attieke. (Djeni et al., 2015).Our studyis a technicalsheet of aprocess of traditionalcassava inocula used for attieke production in south of côte d'Ivoire.

2 MATERIAL AND METHODS

34 **2.1 Materialvegetable**



35
36 **Photo 1: Cassava roots (photo A.K Kouame)**

37
38 The used plant **materialconsisted** of 12 months-**oldfreshlyharvestedcassavaroots** ofthe bitter
39 variety, IAC cultivar (ImprovedAfricanCassava).

40 **2.2 Traditionalcassava inocula preparation**

41 **Cassavaroots** are peeled, **cutintopieces**, **washedthreetimeswithfresh** water and
42 **thenground**,blanched for 10 to 20 minutes, drained, cooled in the open air for 15 minutes,
43 **thenwrapped** in jute bags. The **wholeis** put to ferment for 2 to 3 days in a hot place of the
44 kitchen

Cassavaroots



Peeling



Pulpwashing and cutting



Cooking (10-20 minutes ; 100°C)



Cooling (25-35°C)



Packaging



Fermentation (25-37°C ; 2-3 days)

Figure 1 : Process of cassavatradoinal inocula production used for cassavadough fermentation during attiekeproduction in southern parts of Côte d'Ivoire



Photo 2: Traditional cassava inocula (photo A.K Kouame)

2.3 Determination of pH and Total Titrable Acidity (TTA)

Thirty grams of cassava traditional inocula samples were blended with 70 ml of sterile distilled water and filtered through a Whatman filter paper. The pH of 30 ml of the filtered solution was determined using a pH-meter (pH-meter P107, Consort, Bioblock Scientific, Illkirch, France). TTA was determined using the standard method described by Amoa-Awua et al. 1996. Ten millilitres of filtered solution were titrated with NaOH 0.1 N, using 1% phenolphthalein as indicator. The volume of aliquot used was recorded to determine the amount of acid in the sample. The titrable acidity was calculated as percentage of lactic acid. The determinations were carried out in triplicates and the mean value recorded.

2.4 Enumeration and Identification of Spoilage Microorganisms

Preparation of stock solutions, inoculation of agar plates, cultivation and quantification of micro-organisms were carried out according to (Djeni et al., 2011). For all determinations, 10 g of the traditional cassava inocula samples were homogenized in a stomacher with 90 ml of sterile buffered peptone water (AES Laboratoire, Combourg, France). Tenfold serial dilutions of stomacher fluid were prepared and spread plated for determination of micro-organism counts. Enumeration of coliforms was carried out using plates of Violet Red Bile Lactose agar (VRBL, Merck 10660, Merck, Darmstadt, Germany). The cultures were incubated for 48 h at 30°C for total coliforms and 44°C for faecal coliforms. Yeasts and moulds were enumerated on plates of Sabouraud–chloramphenicol agar (Fluka, Bochemica 89579, Sigma-Aldrich Chemie GmbH, Bangalore, India), incubated at 30°C for 4 days. Aerobic mesophiles were enumerated on plates of plate count agar (PCA Oxoid Ltd, Basingstoke, UK) and incubated at 30°C for 2 days.

2.5 Isolation and Identification of Food-borne Pathogens

2.5.1 *Staphylococcus aureus*

Staphylococcus aureus was isolated and enumerated according to the method described by Capita et al., 2001. A volume of 01 ml of each dilution was surface plated on Baird-Parker agar (BPA) containing egg yolk tellurite emulsion (Oxoid) and incubated at 37°C for 24 and 48 h. The total number of colonies, colonies with different morphology to those of *Staphylococcus aureus* was counted. Five colonies from each sample were randomly selected, purified and tested for cell morphology, arrangement of the cells, Gram reaction, catalase activity, oxidase test, ability to produce acid anaerobically in a glucose-containing growth medium, coagulase activity, thermo-stable nuclease activity, acid production from mannitol and acetoin production. Only, the gram positive cocci were identified using the identification schemes proposed by Schleifer et Kloos, 1975. After the identification, the percentages of *Staphylococcus aureus* and the other strains were calculated. These percentages were later used to correct the results of the counts obtained from each BPA plate.

2.5.2 *Bacillus*

The quantitative estimation of spores of *B. cereus* was performed by a standard plate counting method. Isolations were achieved from heat-treated dilutions by plating on mannitol egg yolk polymyxin B agar (Kouame et al., 2012). Presumptive colonies of *B. cereus* were randomly selected based on characteristic colony feature, purified on the same medium and identified by morphological, cultural and biochemical characteristics according to the documented procedures (Cappuccino and Sherman, 2004).

2.5.3 *Salmonella*

The research of *Salmonella* in cassava traditional inocula, palm oil and water samples were achieved according to the procedures described in the global *Salmonella* surveillance and laboratory support project of the World Health Organization (Hendriksen, 2003). From each sample, 25 g was aseptically weighed and macerated in 225 ml of buffered peptone water (Oxoid) and incubated at 37°C for 24 h. A selective enrichment in Tetrathionate broth (Mueller-Kauffmann) and Rappaport Vassiliadis soy peptone broth using 1 ml of previously incubated buffered peptone water was achieved at 37°C for 24 h, followed by a

subcultivation on Salmonella Shigella agar incubation at 35°C for 24– 48 hours (Feng et al., 2007). Colourless, transparent and with a black centre colonies were further identified using biochemical tests.

2.6 Statistical Analysis

Descriptive statistics for microbiological data were calculated with Excel (Microsoft, Redmond, WA, USA). All statistical analyses were implemented in STATISTICA for Windows ver. 10 (Statsoft Iberica, Lisbon, Portugal). Parametric tests (one-way variance analysis with Duncan's test) at 5% significance level were performed to determine whether there were significant differences between markets regarding microbiological data collected.

3. Results and discussion

In Côte d'Ivoire, *attiéké* plays an important role in the population diet. It is part of the diet of many peoples. It is a typically Ivorian food, whose annual local consumption is estimated at over 450 000 tons (Djeni et al., 2008). The production of *attiéké* necessarily requires the use of ingredient such as the cassava traditional inocula. The traditional cassava inocula used for cassava dough fermentation during *attiéké* production in southern parts of Côte d'Ivoire is usually prepared by the method described above (figure 1). Cassava traditional inocula contain several microorganisms which play an important role in the fermentation of cassava dough for *attiéké* production. The total titrable acidity and pH content of cassava traditional inocula were respectively $1.67\% \pm 0.2$ and 4.94 ± 0.8 . This pH was due to the fact that production of the traditional cassava ferment necessarily requires spontaneous fermentation. Traditional cassava inocula contained aerobic mesophiles, total coliforms, fecal, *Bacillus*, *Staphylococcus* and moulds. Mean loadings of aerobic mesophiles, total coliforms, faecal coliforms, *Bacilli*, *Staphylococcus* and mould were respectively $(2.2 \pm 0.7)10^9$ CFU / g; $(3.7 \pm 1.7)10^5$ CFU / g; $(1.7 \pm 0.7)10^3$ CFU / g; $(2.7 \pm 1.2)10^7$ CFU / g; $(2.2 \pm 1.5)10^5$ CFU / g and $(6.3 \pm 1.9)10^6$ CFU / g (Table 1). Similar observations were reported by Ofuya and Akpoty, 1998 in their study on gari stability. Moreover, the multiplication of microorganisms in the cassava traditional inocula sample was facilitated by the product nature (wet product), its composition and the temperature of storage, which corresponded to some micro-organisms optimal growth temperature. In fact, according to (Giese, 1998; Amoa-Awu and Jakobsen, 1996), an environment containing high sugar and moisture contents

constituted a favourable medium for yeasts and moulds, Enterococci and coliforms development.

Table 1. pH, total titratable acidity and microbial population in cassava traditional inocula used in attieke process

Parameters	Values
pH	4.94± 0.8
Titratable acidity (%)	1.67± 0.2
Aerobic mesophiles (CFU.g ⁻¹)	(2.2±0.7)10 ⁹
Moulds (CFU.g ⁻¹)	(6.3±1.9)10 ⁶
Staphylococci (CFU.g ⁻¹)	(2.2±1.5)10 ⁵
Bacilli (CFU.g ⁻¹)	(2.7±1.2)10 ⁷
Total coliforms (CFU.g ⁻¹)	(3.7±1.7)10 ⁵
Faecal coliforms (CFU.g ⁻¹)	(1.7±0.7)10 ³
Salmonella (CFU.g ⁻¹)	ab

ab, absence in 25 g for Salmonella, values are expressed as mean ± standard deviation.

The presence of such micro-organisms in the cassava traditional inocula could be due to a further contamination after steaming, by the production environment, the material of production, the product handling and during the packaging (Kouame, 2013). These microorganisms constitute the bulk of the inoculum which would participate in the fermentation of the cassava pulp for the production of attieke. (Djeni et al., 2015).

4. Conclusion

Attieke is one of the few products whose fermentation is not spontaneous but involves the use of an inoculum. Traditional cassava inocula used for attieke productions in south of Côte d'Ivoire had an acidic pH. It contained a diversity of microorganisms which would participate in the fermentation of cassava pulp intended for the production of the attieke in Côte d'Ivoire.

REFERENCES

Amoa-Awua WK, Appoh FW, Jakobsen, M (1996). Lactic acid fermentation of cassava dough into 'agbelima'. Int J Food Microbiol; 31 : 87–98.

207 Amoa-Awua, W.K. and Jakobsen, M (1996). The role of microorganisms in the
 208 fermentation of AgbelimaCassavaDough. In TraditionalFermentedProcessing in Africa: The
 209 Third Biennal Seminar on AfricanFermented Food. EdsHalm, M. and Jakobsen, M. pp. 1–7.
 210 Ghana: FRI, DANIDA, KVL.

211 Capita R., Alonso-Calleja M, Garcia-Fernandez MC (2001).Assessment of Baird-Parker
 212 agar as screening test for determination of *Staphylococcus aureus* in poultrymeat. J
 213 Microbiol.;39 : 321–325.

214 Cappucino, J.G. and Sherman, N (2004).Microbiology: A LaboratoryManual, 6th edn.
 215 Singapore: Person Education.

216
 217 Djeni, N.T., N'Guessan, K.F., Toka, D.M., Kouame, K.A. and Dje, K.M. (2011) Quality
 218 of attieke (a fermented cassava product) from the three main processing zones in Côte
 219 d'Ivoire. Food Res Int 44, 410–416.

220 Djeni, N. T., N'guessan, K. F., Dadié, A. T., & Djè, K. M. (2008). Impact of different levels
 221 of a traditional starter on the fermentation of cassava dough for Attiéképroduction. Food, 2(2),
 222 145–151.

223 Djeni N.T, Bouatenin K.M.J.-P., Assohou N.M.C., Toka D.M., Menan E.H., Dousset
 224 X., Dje K.M.(2015).Biochemical and microbial characterization of cassava inocula from the
 225 three main attieke production zones in Côte d'Ivoire. Food control 50 : 133-140

226 Feng, P., Weagant, S. and Grant, M (2007).Enumeration of Escherichia Coli and the
 227 ColiformBacteria. BacteriologicalAnalyticalManual, 8th edn. USA: FDA/Center for Food
 228 Safety and Applied Nutrition.number of pages

229 Giese J (1993). Spices and seasoningblends a taste for allseasoning. J Food Technol.;49 :88-
 230 98.

231 Hendriksen, R.S (2003). LaboratoryProtocolsLevel 1: Training Course Isolation of
 232 Salmonella. A Global Salmonella Surveillance and Laboratory Support Project of the World
 233 HealthOrganization, 4th edn. Geneva: WHO,. ?????????

234 Kouame AK, Djéni TN, N'Guessan FK, Djè KM (2012).Postprocessingmicroflora of
 235 commercial attiéké (afermentedcassavaproduct) produced in the south of Côte d'Ivoire.
 236 Letters in AppliedMicrobiology. ;56 : 44-50.

237 Kouame AK (2013). Identification des dangers et des des points critiques de contrôle pour la
 238 mise en place d'un système HACCP pour la production de l'attieke en Côte d'Ivoire. Thèse

239 de doctorat unique Université NanguiAbrogoua, UFR des Sciences et technologies des
240 Aliments. Number of pages
241 **Ofuya CO and Akpoty P (1998)** Post-processingmicroflora and the shelfstability of gari
242 marketed in Port Harcourt. J ApplBacteriol. ; 64 :389–394.
243 **Schleifer KH and Kloos WE (1975).** Isolation and characterization of
244 staphylococci from humanskin. Int J SystBacteriol.;25 :50–61.
245