

Original Research Article

Heavy Metal Content of Plants from Njere River Bank in Umuosu Okaiuga Nkwoegwu, Umuahia North L.G.A of Abia State

ABSTRACT

Aims: To assess the heavy metal content of plants from Njere river bank in Umuosu Okaiuga Nkwoegwu, Umuahia North L.G.A of Abia state.

Study design: Plant samples were collected from upstream, midstream and downstream locations of the river bank. **And used for the study.**

Place and Duration of Study: Department of Chemistry, Michael Okpara University of Agriculture, Umudike between June 2009 and December 2009.

Methodology: Composite samples were made from the samples from the various locations. They were air-dried, milled, sieved and used for Atomic Absorption Spectroscopy.

Results: Total concentrations of Ni, Cd, Fe, Zn and Pb, soluble metal concentrations of Ni, Cd and Zn and exchangeable metal concentrations of Ni, Cd, Fe, Zn, and Pb were **all** below the W.H.O standard for all the plant samples.

Conclusion: The plants from Njere river bank is relatively safe for animal and human consumption.

Keywords: Heavy metals, plants, food chain, toxicity, bioavailability, accumulation

1. INTRODUCTION

Plants are a basic source of food needed for various biological processes in living organisms [1] [2] [3]. Plants growing in heavy metal environments undergo changes in their physiological and biochemical processes that affect their growth [4] and as such will ultimately lead to food insecurity. Heavy metals have been found in different kinds of plants and food crops including: fruits such as apples, pineapples, oranges, banana) [5]; vegetables such as fluted pumpkin [6] and even teas [7].

Heavy metals are elements that possess high relative atomic weight with atomic number that is greater than 20 and possess attributes of metals [8] [3]. They are highly recalcitrant [9], toxic and occurs naturally in the earth. Plants are able to take up heavy metals that are easily solubilized in soil solution and root exudates[10]. Heavy metals cause oxidative stress that alters cell function and enzyme activities [11]. Kibra, [12] reported that rice plants cultivated on Hg polluted soils showed decline in height. Ahmad et al. [13] reported that under Cd toxicity, the root and roots of wheat plants declined in their growth. Another study by Ghani, [14] showed a decline in the growth and protein content of maize plant grown under a combined heavy metal toxicity of Cd, Cr, Co, Mn, Hg and Pb. In a review article, Peralta-Videa et al. [15] documented the processes of plant uptake and bioaccumulation of heavy metals, including their negative effects on man.

Plants source is one of the known ways by which heavy metals are introduced into the food chain. This pose a serious challenge as humans and other lower animals may be potentially at risk as they consume plants contaminated by heavy metals. This research was therefore conducted to assess the heavy metal content of plants growing around the Njere river. The result of this experiment will help offer useful advice to the rural dwellers on possible mechanisms to employ to ensure production of healthy foods and crop yield in the area.

2. MATERIAL AND METHODS

2.1 SAMPLE COLLECTION AND PREPARATION

The plant sample were collected by forming composites of plants collected from the bank of the Njere river based on the upstream, midstream and downstream locations. The samples were labeled accordingly in clean polythene bags. The samples were dried at room temperature, milled and sieved using 0.01 μ m sieve. The ground plant sample was stored in clean polythene bags and well labeled. The samples were collected in April, 2009.

2.2 DETERMINATION OF TOTAL METAL CONCENTRATION

From the ground plant sample, 1g was weighed out into a conical flask. It was digested by adding 15mls of HNO₃, 10mls of HCL and heated at 100°C in a fume cupboard for 30 minutes. Thereafter, 25mls of deionized water was added and the heating continued for 90 minutes. It was allowed to cool, filtered with whatmann no 1 filter paper into a 50ml volumetric flask and made up to mark with deionized water. Ten mls of the filtered solution was measured into clean sample bottles for determination of Pb, Zn, Fe, Cd and Ni using Atomic Absorption Spectrophotometer (AAS 240 FS Agilent Technologies).

2.3 DETERMINATION OF WATER SOLUBLE METAL CONCENTRATION

One gram of plant sample was weighed into a flask to which was added 20mls of deionized water and allowed to stand for 30 minutes. It was filtered into 50mls volumetric flask using whatmann no 1 filter paper and made up to mark with deionized water. A quantity of 10mls of the filtrate was measured out into clean sample bottle for analysis with AAS.

2.4 DETERMINATION OF EXCHANGEABLE METAL FRACTION

One gram of soil sample and twenty mls of 1M MgCl₂ were poured into a clean conical flask. The pH of the solution was adjusted using potassium phosphate buffer, boiled for 60 minutes at ambient temperature. It was allowed to cool, filtered into a volumetric flask and made up to mark with deionized water. Ten mls of the filtrate was measured out into clean sample bottles for exchangeable metal fraction determination using AAS.

STATISTICAL ANALYSIS

Results are presented as means of duplicate determinations.

3. RESULTS AND DISCUSSION

The result of the Total metal ions in plant samples from around Njere river is depicted on Table 1. The Ni, Cd, Fe, Zn and Pb concentration in the plants from the upstream, midstream and downstream were all below the W.H.O standard for each metal.



Table 1. Total metal ions in plant samples from around Njere river

89
90

Total of metal ion	Location	mean conc of metal ion (ppm)	W.H.O standard
Ni	Upstream	0.242	1-10
Ni	Midstream	0.108	1-10
Ni	Downstream	0.145	1-10
Cd	Upstream	0.137	<0.5
Cd	Midstream	0.025	<0.5
Cd	Downstream	0.1	<0.5
Fe	Upstream	1.881	250
Fe	Midstream	1.338	250
Fe	Downstream	3.349	250
Zn	Upstream	1.265	5-100
Zn	Midstream	0.502	5-100
Zn	Downstream	0.718	5-100
Pb	Upstream	0.164	6-9
Pb	Midstream	0.021	6-9
Pb	Downstream	0.162	6-9

91
92

Values are means of duplicate determinations

93

94

95

96

97

98

99

100

101

Table 2. Water soluble metal ions in plants from around Njere river

Type of metal ion	Location	mean conc of metal ion (ppm)	W.H.O standard
Ni	Upstream	0.021	1-10
Ni	Midstream	0.187	1-10
Ni	Downstream	0.05	1-10
Cd	Upstream	0.008	<0.5

Cd	Midstream	0.045	<0.5
Cd	Downstream	0.002	<0.5
Fe	Upstream	0.056	250
Fe	Midstream	ND	250
Fe	Downstream	0.019	250
Zn	Upstream	0.262	5-100
Zn	Midstream	0.271	5-100
Zn	Downstream	0.258	5-100
Pb	Upstream	0.001	6-9
Pb	Midstream	ND	6-9
Pb	Downstream	0.003	6-9

Values are means of duplicate determinations

Table 3 shows the result of the Exchangeable metal ions in plants from around Njere river. In all the plant samples from the different locations, the concentrations of Ni, Cd, Fe, Zn, and Pb were all below the W.H.O standards of 1-10ppm, < 0.1ppm, 250ppm, 5-100ppm and 6-9ppm respectively.

Table 3. Exchangeable metal ions in plants from Njere river

Total of metal ion	Location	mean conc of metal ion (ppm)	W.H.O standard
Ni	Upstream	0.213	1-10
Ni	Midstream	0.227	1-10
Ni	Downstream	0.137	1-10
Cd	Upstream	0.118	<0.5
Cd	Midstream	0.1	<0.5
Cd	Downstream	0.075	<0.5
Fe	Upstream	0.135	250
Fe	Midstream	0.116	250
Fe	Downstream	0.109	250
Zn	Upstream	0.203	5-100
Zn	Midstream	0.18	5-100
	Downstream	0.1	5-100

Zn

Pb	Upstream	0.049	6-9
Pb	Midstream	0.043	6-9
Pb	Downstream	0.025	6-9

Values are means of duplicate determinations

The total heavy metal ion concentration is an indication of all the chemical forms present in a given sample. The result shows very low accumulation of total heavy metal ion by plants in the study area (Table 1). This could be possibly due to the fact that the study area is still a suburb with very minimal industrial activities. Most of the heavy metals assessed could be basically from natural sources such as weathering and volcanic eruptions. Dossa et al. [16] and Ives and Kendal [17] reported that a combination of agricultural, industrial and anthropogenic activities are key sources of increased heavy metals in the environment. However, Pueyo et al. [18] reported that total heavy metal ion concentration alone cannot be used to evaluate the mobility, bioavailability and any possible toxicity of metals.

The low concentration of soluble heavy metals accumulated by the plants could also correlate the concentration of total metal ions (Table 2). Increases in soil pH has been implicated in mobilization of heavy metals in plants [19], as well as soil properties [20].

The results obtained in this study have lent credence to the fact that the study area Umuosu Okaiga Nkwoegwu community is still under-developed with little or no industrial activities. Also, ingestion of the plants by animals and humans may be relatively safe.

CONCLUSION

Since man continues to depend on food for survival, there is need to ensure production of quality food products for health and sustainability of human and animal health. The urban cities still depend greatly on the rural communities for these food products. The Njere river bank can therefore serve as a good site for all round year crop production. The water can also support irrigation agriculture.

REFERENCES

1. Iweala EEJ, Olugbuyiro JAO, Durodola BM, Fubara-Manuel DR, Okoli AO. Metal contamination of foods and drinks consumed in Ota, Nigeria. *Res. J. Environ. Toxicol.* 2014; 8:92–97.
2. Izah SC, Aseiba ER, Orutugu LA. Microbial quality of polythene packaged sliced fruits sold in major markets of Yenagoa Metropolis, Nigeria. *Point J. Bot. Microbiol. Res.* 2015;1: 30–36.
3. Izah SC, Kigigha LT, Anene EK. Bacteriological Quality Assessment of *Malus domestica* Borkh and *Cucumis sativus* L. in Yenagoa Metropolis, Bayelsa state, Nigeria. *Br. J. Appl. Res.* 2016;1:5–7.
4. Oancea S, Foca N, Airinei A. Effects of heavy metals on plant growth and photosynthetic activity,” *Analele Științifice ale Universității “AL. I. CUZA1 IAS1, Tomul I, s. Biofizică, Fizică medicală și Fizică mediului.* 2005;1-4.
5. Unaegbu M, Engwa GA, Abaa QD, Aliozo SO, Ayuk EL, Osuji GA, Onwurah EI. Heavy metal, nutrient and antioxidant status of selected fruit samples sold in Enugu, Nigeria. *Int. J. Food Contam.* 2016;3-7.
6. Nwadinigwe CA, Udo GJ, Nwadinigwe AO. Investigations of Heavy Metals Concentrations in Leaves of *Telfairia occidentalis* Hook. F. (Fluted Pumpkin) in Nigeria. *Pol. J. Environ. Stud.* 2015;24:1733–1742.

- 156
157 7. Garba ZN, Ubam S, Babando AA, Galadima A. Quantitative Assessment of Heavy Metals from
158 Selected Tea Brands Marketed in Zaria, Nigeria. J. Phys. Sci. 2015; 26:43–51.
159 17.
160
- 161 8. Chounwou PB, Yedjou CG, Patlolla AK, Sutton DJ. Heavy Metals Toxicity and the Environment.
162 Natl. Inst. Health. 2012;101:133–164.
163
- 164 9. Ogunlana OO, Ogunlana OE, Akinsanya AE, Ologbenia OO. Heavy metal analysis of selected soft
165 drinks in Nigeria. J. Glob. Biosci. 2015;4:1335–1338.
166
- 167 10. Blaylock MJ, Huang JW. Phytoextraction of metals, in *Phytoremediation of Toxic Metals: Using Plants*
168 *to Clean up the Environment*, I. Raskin and B.D. Ensley, Eds., pp. 53–70, Wiley, New York, NY, USA,
169 2000.
170
- 171 11. Jadia CDCD, Fulekar MH. Phytoremediation of heavy metals: recent techniques. African Journal of
172 Biotechnology. 2009;8(6):921–928.
173
- 174
- 175 12. Kibra MG. Effects of mercury on some growth parameters of rice (*Oryza sativa* L.). Soil &
176 Environment. 2008;27(1): 23–28.
177
- 178 13. Ahmad I, Akhtar MJ, Zahir ZA, Jamil A. Effect of cadmium on seed germination and seedling growth
179 of four wheat (*Triticum aestivum* L.) cultivars. Pakistan Journal of Botany. 2012;44(5):1569–1574.
180
- 181 14. Ghani A. Toxic effects of heavy metals on plant growth and metal accumulation in maize (*Zea mays*
182 L.). Iranian Journal of Toxicology. 2010; 3(3):325–334.
183
- 184
- 185 15. Peralta-Videa JR, Lopez ML, Narayan M, Saupe G, Gardea-Torresdey J. The biochemistry of
186 environmental heavy metal uptake by plants: Implications for the food chain Int. J. Biochem. Cell Biol.
2009; 41(8–9):1665–1677.
- 187
- 188 16. Dossa LH, Sangaré M, Buerkert A, Schlecht E. Intra-urban and peri-urban differences in cattle
189 farming systems of Burkina Faso. Land Use Policy. 2015;48:401–411.
- 190
- 191 17. Ives CD, Kendal D. Values and attitudes of the urban public towards peri-urban agricultural land.
192 Land Use Policy. 2013;34:80–90.
- 193
- 194 18. Pueyo M, López-Sánchez JF, Rauret G. Assessment of CaCl_2 , NaNO_3 and NH_4NO_3 extraction
195 procedures for the study of Cd, Cu, Pb and Zn extractability in contaminated soils, Anal. Chim. Acta,
196 2004;504:217–226.
- 197
- 198 19. Naidu R, Bolan NS, Megharaj M, Juhasz AL, Gupta SK, Clothier BE, Schulin R. Chemical
199 bioavailability in terrestrial environments, in: Developments in Soil Science, 32, Chemical bioavailability in
200 terrestrial environments, edited by: Hartemink, A. E., McBratney, A. B., and Naidu, R., Elsevier, Oxford,
201 UK, 2008;1-6.
- 202
- 203 20. Li HF, Gray C, Mico C, Zhao FJ, McGrath SP. Phyto-toxicity and bioavailability of cobalt to plants in a
range of soils. Chemosphere, 2009; 75: 979–986.