

Histological Development of Grafting in Apple

In Cold and High Altitude Conditions

Oğuz Dolgun^{1*}Hamdi Zenginbal²

¹ Adnan Menderes University, Sultanhisar Vocational College, Aydın, Turkey

²Abant İzzet Baysal University, Bolu Vocational College, Bolu, Turkey

*Corresponding authors' e-mail: odolgun@gmail.com

ABSTRACT

The purpose of this study is to briefly describe graft fusion in relation tissue formation in some apple varieties grafted by chip and T buddings on MM106 in cold climatic and high altitude conditions. Effects of enviromental conditions on plant growing is well known. Plants grow strongly in warm and humid places. But, in cold climate and high altitude conditions grow weak. Therefore, graftings which made in different ecological condition may show different grafting formation time. In such places it is important that the graft fusion takes place rapidly. The short-term fusion will be effective in faster and stronger development of grafted plants. For his reason, it is important to know the most useful grafting method in high and cold places. In this research, it was observed that stocks and scions fits very well and no large gaps and although the callus formation was shorter in chip budding, it soon filled all the voids, and as a result, the cambial continuity and strong vascular connection occurred in short time. For these reasons, chip budding were found more suitable due to less necrotic tissues formation and wounds heal more quickly compared to classic "T" budding for production of apple sapling in cold climate and high altitude conditions.

Keywords: Grafting, Apple, Altitude, Graft Techniques, Histology

27 1. INTRODUCING

28 Effects of enviromental conditions on plant growing is well known.Plants grow
29 strongly in warm and humid places compared with cold climate regions [1,2,3,4]. In
30 this relation, also tissues formation in grafting is in relation with ecological conditions
31 and especially temperatures and humidity affect speed of graft formation and scion
32 growth after grafting [5,6].A good graft formation results high quality saplingsgrowing
33 in short time. But, grafts made in different ecological condition like cold and high
34 regions can show different graft formation time and saplings can show different
35 growth rate and quality. For this reason the establishment of nurseriesin places over
36 800 meters is not prefered. Nurseries are usually established where the sea level or
37 the maximum height of 300 meters above sea level. However, without the possibility
38 of establishing sea level nursery places, or when required to produce saplings in high
39 places, environmental conditions are effective in sapling production [7]. Therefore,
40 rapid realization of graft fusion is important in such places.Short-term graft fusion will
41 be effective in faster and stronger development of saplings. Therefore, it is important
42 to know the most appropriate grafting method for high and cold places.

43 On the other hand, histological developments in graft union formation can provide
44 some information about graft success. Major events in graft formation are cohesion of
45 scion and rootsck, proliferation of callus cells, connection of cambiums and formation
46 of vascular continuity.The structural integrity of the graft union not only holds the
47 grafted plant together but it is the reestablishment of anatomical and functional
48 continuity between xylem and phloem that allows for transportation of water and
49 minerals by the xylem, conduction of carbohydrates and other organics by the
50 phloem. The necrotic plates are the layers of crushed and desiccated cell walls at
51 the cut surface of both stock and scion.In addition, necrotic plates serve to seal
52 injured tissue from pathogens and to limit water loss. In most species, callus is not
53 from the vascular cambium itself, but rather from the secondary xylem and phloem
54 cells that were most recently formed from division of the vascular cambium. As the
55 new callus increases in volume, it ruptures the necrotic plate and begins to expand
56 into whatever spaces exist between stock and scion. New vascular cambium
57 differentiates inwards from the vascular cambiums cut ends of the stock and scion.
58 New vascular cambium cells begin to produce to the inside (xylem) and to the outside
59 (phloem). Regeneration and bridging of conducting elements (xylem tracheids and

vessels, and phloem sieve tubes) allows for transport across stock and scion. Besides, interlocking of new xylem fibers is largely responsible for mechanical strengthening [1,8,9,10,11,12,13,14,15,16].

The aim of this study is to briefly describe graft union in relation to tissue formation in some apple varieties grafted on MM106 in cold climatic and high altitude conditions.

2. MATERIAL AND METHOD

Grafting studies were carried out in the experimental orchard of Bolu Vocational College of Abant İzzet Baysal University. Bolu province is placed in the northern region of Turkey and it has cold climate conditions and high altitude (768m). Graft sections were examined under microscope in Sultanhisar Vocational College, Adnan Menderes University.

In this study, one year old semi-dwarf MM106 was used as rootstock. MM106 is a semi-dwarf apple rootstock that can be easily and cheaply produced when compared to dwarf M9. Because of this feature, it is preferred in recent years. Apple varieties; Golden Delicious, Granny Smith, Mondial Gala and Red Chief were grafted on root stock by using “T” and “Chip” budding techniques at 25 April 2012. 100 plants were budded for each combination.

Graft samples of each rootstock/scion combinations were taken at first 15th day and 30th days intervals until 4 months after grafting. Samples were fixed in ethanol (70%). Transverse sections in 0.2-0.3mm thickness were cut using a rotary microtome. All sections were examined and photographed with digital microscope Mic-D having colouring property. Five basic stages of graft formation; (1) development and positions of necrotic layers, (2) proliferation of callus cells, (3) formation of callus bridge at the graft interface (4), cambial continuity and (5) formation of vascular tissues were examined on samples. Meteorological records of region were given in Table 1.

(Table 1)

3. RESULTS

3.1 Histological observations in “T” budding samples

In 15 days samples, There was seen many necrotic layers in all combinations. Necrotic layers formed from dead cells occurred as a result of cuts was clearly visible in the cut sides of scion and especially bark of the stock which cover up scion in all combinations. It was seen low callus formation in all combinations, but necrotic layers were started to fragment by newly formed callus cells. It was seen that callus tissues were produced by cortex of scion and stock.

Sufficient callus formation were observed in 30 days sections of all combinations. Necrotic layers was fragmented mostly by the callus tissue. It was seen in the form of vesicular callus mass which spread into gaps between stock and scion. There was not observed tissue formation related cambial differentiation (Fig. 1).

An intensive callus activity and absorption of necrotic layers were seen in the graft interface of both scion and stock in 60 days sections. It was seen that the callus continued to proliferate and almost filled all gaps. An indistinct cambium differentiation were seen in callus tissue close places to uninjured cut end sides of graft members in 60 days sections (Fig. 2). Largely eliminated necrotic layers were seen as small points and thin black lines in callus tissue.

Cambial continuity was clearly seen in 90 days sections of combinations. Cambium bridge is curly form because of the callus mass and necrotic layers. Newly formed xylem and floem were seen as locally strands in callus tissue between scion and stock (Fig. 3). But vascular tissues was not sufficient for strong graft formation.

In the sections taken on 120th days samples, callus was almost fill all gaps and vascular bridge were established between scion and stock. It was not seen incompatibility related histologically development (Fig. 4).

(Figure 1-4)

3.2 Histological observations in “Chip” budding samples

In 15 days sections, It was seen clear but small amount necrotic layers placed close to cut interface of scion and stock. Callus production was low but sufficient. There are thin gaps between scion and stock especially xylem tissues of both stock and scion.

Newly formed callus was produced from cortex tissues close to cut sides of cambiums of stock and scion.

In 30 days sections, A strong callus formation were seen in all combinations. Necrotic layers formed soon after due to cuts was clearly seen in cut sides of stock and scion. Necrotic layers was broken by the newly formed callus tissues. There was not seen a clear evident related cambial differentiation (Fig. 5).

Clear cambial continuity was observed in 60 days samples of all combinations. An intensive cellular activity was seen in all combinations. Cellular activity was not only for callus proliferation but also absorption of necrotic layers. Necrotic layers were largely eliminated by callus. They were seen in cortex region close to bark as thin black lines and small black points in callus (Fig. 6). New cambium derived from callus cells started to produce vascular elements. Especially, new xylem cell strands were observed in callus along graft line close to bark (Fig. 7).

In the sections taken on 90th day and following days, it was seen that callus completely filled all gaps between stock and scion. Xylem and phloem tissues were successfully bridged scion and stock. Necrotic layers were absorbed by proliferated callus. There were seen a few necrotic layers as small points. It was not seen an abnormal histological or cellular development (Fig. 8).

(Figure 5-8)

4. DISCUSSIONS

Usually, a successful grafting includes formation of necrotic layers, callus proliferation, elimination of necrotic layers by proliferated callus, differentiation of some cells to cambium cells, establishing of cambial continuity and bridging of vascular tissues finally. Necrotic layer formation as a result of wounding is a defensive mechanism against to invasion of pathogens and it is an independent event. In subsequent days, cortex tissue starts to produce callus. In time, callus tissue continues to proliferation and spread into gaps placed between stock and scion. In the mean time, necrotic layers have been eliminated by callus. Callus absorbs most of the necrotic layers [17,18,19,20]. Also in this study, all grafts showed a similar necrotic layer formation and elimination. Especially in the "T" graft samples a high amount of necrotic layer formation was observed. It is a result of excessive cuts and

woundings especially in stock. It is well know that dead cells due to wounding are transformed to black necrotic strands or points. As result of this, it was seen necrotic strands and points in callus and cut surface even 90 days samples. Necrotic layers was not completely absorbed by callus which proliferated from cortex. This result is in agreement with previous researches which indicated wounding is much in “T” buddings compared chip budding. On the other hand,Chip budding samples showed that wounding is very small amount. As a result of small wounding, necrotic layers was small amount and generally placed in cut sides of cortex.It was clearly that callus cells proliferated from cortex paranchyma of stock and scion. In the course of time, callus which spreaded in gaps were fill all gaps between scion and stock and necrotic layers were absorbed. Thus, our results are in agreement with prevous researches [1,17,20].

In the present research, most clear difference was seen in establishing cambial continuity. Cambial continuity was clearly seen in 90 days samples of “T” budding, but in “chip” budding samples, cambial continuity were seen in 60 days samples. This situation may be a result of the excessive wounding and long filling time of the callus for gaps in “T” grafts. Because, it was seen that much necrotic layers and large gaps between scion and stock

But, in “chip” budding samples, it was seen small amount necrotic layers. Also, it was seen small gaps between xylem and cortex zones of scion and stock due to close cuts surfaces of graft members. Gaps between scion and stock were filled in short time by callus and a strong cambial continuity established in short time. In 90th and other periodsof samples of all combinations,smiliar observations were seen. Results is agreement with previous reseraches [10,22,23].According to Hartmann et all. [1] “T” budding is generally used in propagation of fruit trees and chip budding is not preffered because it is not simplier and faster. But, in contrast Kviklis [22] reported that chip budding gives higher graft success in apple.Researchers reported that chip budding was more effective than “T” budding [24]. Smilar results were also reported by many authors [25,26,27].

Kadan and Yarılıgac who studied apple and pear sapling production by using “T”budding in Van ecological condition which has cold climate and high altitude (1727 m) reported that it couldn’t be obtained any first or second class saplings [7].

5. CONCLUSION

Chip budding were found more suitable than classic “T” budding for production of apple sapling in cold climate and high altitude conditions due to less wounding, less necrotic layer formation and wounds heal more quickly. In addition, it was observed that stock and scions fit very well and there are no large gaps between them in chip budding. Callus formation is less and fills all gaps in short time. As a result of this, clear cambial continuity and strong vascular connection occurred in short time. Histological observations have shown that the chip budding is relatively more suitable due to quick and strong fusion of tissues in the shorter time in cold and high places.


REFERENCES

1. Hartmann H.T, Kester DE, Davies FT, Geneve LR. Plant propagation. principles and practices. seventh edition. regents / prentice hall international editions. Englewood cliffs, New jersey. USA, 2002.
2. Duran C and Gulek H.. Effects of the ecological factors on vegetation in river basins of northern part of mersin city (south of Turkey). Biological Diversity and Conservation. 2010;3(3): 137-152
3. Kose B. The Role and Importance of The Light and Temperature in Viticulture. Turk J Agric Res. 2004;1: 203-212
4. Kaya B and Aladad C. Precipitation, temperature and vegetation relations in the conditions of Konya. Selcuk University Journal of Social Sciences institute. 2009.;22: 465-478
5. Demirsoy H, Bilginer S. Anatomically Investigation of Graft Union in Some Compatible and Incompatible Peach/Plum Grafts Combinations. J. of Fac. of Agric. OMU. 2006;21(1):89-94
6. Ozkaynak E and Samancı B. Recent advances in environmental control in micropropagation, Batı Akdeniz Agricultural Research Institute, Derim. 2003;20 (1):450-462
7. Kadan H. and Yarılgac T.. Studies on propagation by dormant t-budding of apples and pears under van ecological conditions, Yüzüncü Yıl University, Agricultural Faculty. Journal of Agricultural Sciences (J. Agric. Sci.). 2005;15(2): 167-176
8. Atkinson C.J, Else M.A, Taylor L. and Dover CJ. Root and Stem hydraulic conductivity as determinants of growth potential in grafted trees of apple (*Malus pumila* Mill). Journal of Experimental Botany. 2003;54 (385):1221-1229


- 214 9. Moore R. A Model for graft compatibility–incompatibility in higher plants,
215 Amer.J.Bot., 1984;71(5):752-758
- 216 10. Simons RK. Graft union characteristics as related to dwarfing in apple (*Malus*
217 *domestica* Borkh). Acta Hort. 1986;160: 57-66
- 218 11. Soumelidou K, Battey NH, John P, Barnett JR. The anatomy of the developing
219 bud union and its relationship to dwarfing apple. Annals of Bot. 1994;74:605-611
- 220 12. Vachun Z. Rootstock for Apricot. The current situation and main problems. Acta
221 Hort, 1995:384:459-465.
- 222 13. Gryzb Z.S, Sitarek M. Growth and cropping of plums grafted on pixy rootstock
223 and planted in differentiated density. VI International Symposium on Plum and
224 Prune Genetics, Breeding, Pomology. 1998;478:14-16.
- 225 14. Kankaya A, Ozyigit S, Tekintas FE, Seferoglu G. Compatibility of Some Plum and
226 Apricot Cultivars on Pixy Rootstock. Third National Horticulture Congress, Ankara,
227 1999;1: 295-299.
- 228 15. Errea P, Garay J, Marin A. 2001. Early detection of graft incompatibility in apricot
229 (*Prunus armeniaca* L) using in vitro techniques. Physiologia Plantarum.
230 2001;112:135-141
- 231 16. Ikeuchi M, Sugimoto K and Iwase A. Plant callus: mechanisms of induction and
232 repression, Plant Cell. 2013;25(9): 3159–3173.
- 233 17. Stoddart FL, McCully ME. Effects of stocks and scion organs on the formation of
234 graft union in coleus, a histological study, Botanical Gazette 1980;141(4): 401-
235 412
- 236 18. Estrada-Luna A.A, Lopez-Peralta C, Cerdanes-Soriano E.. In vitro micrografting
237 and the histology of graft union formation of selected species prickly pear cactus
238 (*opuntia* spp). Sci. Hort. 2002;92:317-327
- 239 19. Sitarek M. Incompatibility problems in sweet cherry trees on dwarfing rootstock,
240 Latvian J. Of Agro. 2006;9:140-145
- 241 20. Pina A and Errea P.. Influence of graft incompatibility on gene expression and
242 enzymatic activity of UDP glucoze pyrophosphorylase. Plant Sci. 2008;174: 502-
243 509
- 244 21. Dolgun O, Tekintas FE, Ertan E.. A histological investigation on graft formation of
245 some nectarine cultivars grafted on pixy rootstock. World Journal of Agricultural
246 Sciences. 2008;4(5): 565-568.

22. Kviklis AM. Rationalization of budding methods. Horticultural Abstracts, 1986;56::8511
23. Tekintas FE, Dolgun O. An investigation on compatibility in some peach and nectarin cultivars grafted on almond seedlings. Yuzuncu Yıl Univ. Journal of Agric. Faculty. 1996; 6(1):51-54.
24. Skene D.S, Shepherd HR, Howard BH. Characteristic anatomy of union formation in T and chip-budded fruit and ornamental trees. Journal of Horticultural Science, 1983;58(13): 295-299.
25. Howard B.H, Skene DD, Coles JS. The effects of different grafting methods upon the development of one-year-old nursery apple trees. Journal of American Social and Horticultural Sciences. 1974;49(3): 287-295.
26. Stoyan I. The application of chip budding to fruit tree propagation. Horticultural Abstract. 1984;54-5098
27. Czarneck B. Comparative study of two methods of apple budding. Horticultural Abstract. 1990;60(2):3108.


274 Table 1. . Meteorological records in graft union formation period




276 ("T" Budding sections)



278 Fig 1.30 days section of Golden Delicious "T" budded on MM106




280 Fig 2. 60 days section of Granny Smith "T" budded on MM 106



281


282 Fig 3.90 days section of Mondial Gala "T" budded on MM106



283

284 Fig 4.120 days section of Red Chief "T" budded on MM106

285 (Chip Budding sections)



286

287 Fig 5.30 days section of chip budded Mondial Gala on MM106




Fig 6. 60 days section of Red Chief chip budded on MM106




Fig 7. 90 days section of Golden Delicious chip budded on MM106




Fig 8. 90 days section of Granny Smith T budded on MM 106