

Inheritance pattern of yield affecting attributes in F₂ wheat populations

ABSTRACT

Inheritance mechanism for earliness and yield influencing traits in 8 x 8 F₂ half diallel progenies was studied at The University of Agriculture, Peshawar-Pakistan during 2006 07. The experiment was planted in randomized complete block (RCB) design having four replications. Highly significant differences were observed among parental genotypes and their F₂ populations for all traits revealing greater genetic variability and thus allowing for further analysis. According to genetic analysis, Vr/Wr graphs indicated over-dominance type of gene action for all traits among parental genotypes and their F₂ populations. Traits like days to heading, plant height and spike length were controlled by recessive genes while grains per spike and biological yield were under dominant gene control. Distribution of array points along the regression line indicated that wheat cultivars Takbeer and the landrace Khattakwal possessed maximum dominant and recessive genes, respectively for plant height. However, cultivar Iqbal-2000, Ghaznavi-98, Margalla-99 and Tatara-96 possessed maximum dominant genes for days to heading, spike length, grains per spike and biological yield, respectively.

Keywords: *Diallel analysis, additive and non-additive gene action, over-dominance, inheritance, Triticum aestivum L.*

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is one of the most important cereals not only in Pakistan but in most parts of the world. It is a major staple food crop that fulfills nutritional requirement of the population in the form of protein and carbohydrates. Wheat occupies more acreage than any other crop in Pakistan due to its suitability under irrigated as well as rainfed conditions all over the country (PBS-2015-16). Wheat produced in the country fulfills most part of our requirements; however, the increasing population demands further improvement in wheat production (Jadoon *et al*, 2012)

Hayman's genetic analysis is a useful tool which has been widely used by plant breeders for constructing their desirable plant type using available genetic resources (Hayman 1954a, b). It provides useful information regarding nature of gene action and inheritance pattern of a

particular trait that directs the plant breeders to practice selection in earlier generations or delay till advanced generations. Earlier studies revealed the importance of additive as well as non-additive gene action for inheritance of various earliness and yield associated traits among different wheat populations (Kutlu and Olgun, 2015). Additive type of gene action with partial dominance for spikelets per spike was reported by Ivanovska *et al* (2000) while Sangwan and Chaudhary (1999) identified non-additive gene action for grain yield and related traits in wheat. Similarly Singh *et al* (2002) reflected additive gene control for grains per spike, number of tillers per plant and grain yield in wheat. However, Navabi *et al* (2004) reported transgressive segregation among all the F₂ wheat populations and F₅ single seed-decent lines in wheat. Moreover, Jadoon *et al* (2012), Rabbani *et al* (2009) and Akram *et al* (2008) reported significance of both additive and non-additive gene action in inheritance of various yield related traits and delay in selection for improvement of these traits. In light of past review, the present study was designed to investigate the inheritance mechanism involved in earliness and other yield related traits among eight parental genotypes and their F₂ wheat populations.

2. MATERIALS AND METHODS

2.1. Breeding material and procedure

The experimental material consisted of eight wheat cultivars/lines viz. Tatara-96, Wafaq-2001, Takbeer, Iqbal-2000, Margalla-99, Ghaznavi-98, Khattakwal and Inqalab-91, and their 28 F₂ progenies. The experiment was planted in RCBD having four replicates at The University of Agriculture, Peshawar-Pakistan during 2006-07. Each experimental unit consisted of a four row plot of five meter length with a row to row spacing of 30 cm. The Nitrogen (N) and Phosphorus (P) were applied at 120:60 NP kg ha⁻¹, respectively in form of Urea and DAP fertilizers. Half dose of Nitrogen and full dose of Phosphorus were applied at the time of seed bed preparation while the remaining half dose of Nitrogen (urea) was applied at the time of first irrigation. Recommended cultural practices including land preparation, fertilizers, irrigation and weed control were applied as and when required uniformly to all treatments in the experiment to minimize the environmental effects.

2.2. Data Collection

Ten plants from parental cultivars/lines and thirty from F₂ generations were selected at random to collect data on days to heading, plant height, spike length, grains per spike and biological yield. Days to heading were recorded as number of days taken from germination to 50% spike emergence while plant height was recorded from ground level to tip of spike excluding awns. Spike length was measured from base of 1st spikelet to tip of last spikelet,

excluding awns. Similarly, grains per spike were recorded as an average of produce of thirty spikes obtained after manual threshing and biological yield by weighing the selected plants from each sub-plot/replication in parental genotypes as well as F_2 populations after harvesting and sun drying.

2.3. Statistical analysis:

After compilation, data for all the traits were analyzed using statistical package 'SAS 6.12' following Steel *et al* (1997) to see whether significant differences existed among the genotypes for further analysis. Traits showing significant differences were further subjected to "t²" and "regression analysis" for adequacy of additive-dominance model and Hayman (1954 a, b) diallel analysis to study nature of gene action controlling these traits using computer software 'Dial-98' for half diallel F_2 wheat populations following procedures outlined by Morley-Jones (1965).

3. RESULTS AND DISCUSSION

3.1. Days to heading

According to Hayman's genetic analysis, Vr-Wr graph reflected involvement of over-dominance for days to heading as the regression line cuts the Vr-axis below the origin (Fig. 1a). Cultivars Iqbal-2000, Wafaq-2001 and Tatara-96 possessed maximum dominant genes being closest to the origin while Margalla-99 being farthest from origin possessed maximum number of recessive genes for days to heading. Genotypes Khattakwal and Margalla-99 showed maximum influence of environment for earliness as they were away from the regression line (Fig. 1a). Moreover, non-significant deviation of regression line from unit slope confirmed the absence of non-allelic interactions for days to heading. Positive correlation ($r = 0.49$) of Vr+Wr values with parental values reflected the importance of recessive gene for controlling days to heading (Fig. 1b). These results confirmed the earlier findings of Ivanovska *et al* (2000), Mishra *et al* (1994b) and Singh *et al* (2003) who reported over-dominance to be responsible for yield related traits in bread wheat population.

3.2. Plant height (cm)

Vr-Wr graph revealed that over-dominance was mainly responsible for controlling plant height as the regression line cuts the Vr-axis below origin (Fig. 2a). Earlier researcher (Ivanovska *et al* 2000, Mishra *et al* 1994b and Singh *et al* 2003) also reported over dominance for quantitative traits in F_2 wheat populations. However, Ali and Khan (1998), Joshi *et al* (2004), Singh *et al* (2002) and Zwart *et al* (2004) reported additive genetic effects for most of the traits.

Considerable variation existed among wheat genotypes for plant height as indicated by the distribution of parents along regression line inside the limiting parabola. Wheat cultivars Takbeer, Wafaq-2001, Ghaznavi-98 and Iqbal-2000, being closer to origin, possessed maximum number of dominant genes while Khattakwal possessed maximum recessive genes for plant height. Margalla-99 and Inqalab-91 being in the middle of regression line possessed equal number of dominant and recessive alleles (Fig. 2a). When V_r+W_r values were plotted against the parental values, positive correlation ($r = 0.74$) reflected involvement of recessive genes for plant height (Fig. 2b). Moreover, the position of parental genotypes along the regression line and non-significant deviation of regression line from unit slope confirmed the influence of environmental conditions on plant height and absence of non-allelic interactions for inheritance of this trait.

3.3. Spike length (cm)

Genetic analysis for spike length revealed that distribution of parental cultivars along the regression line confirmed sufficient amount of variation for spike length. The regression line cut the V_r -axis below the origin indicating that over-dominance played important role in the inheritance of spike length (Fig. 3a). Position of array points along the regression line showed that cultivars Ghaznavi-98 and Tatar-96 possessed maximum number of dominant genes while genotypes Margalla-99 and Inqalab-91 possessed maximum recessive genes for spike length. However, cultivars Khattakwal and Iqbal-2000 being in the centre of regression line possessed equal number of dominant and recessive genes. Ivanovska *et al* (2000), Mishra *et al* (1994) and Singh *et al* (2003) also reported similar results in their studies. However, contrary to present findings Khan and Ali (1998) and Singh *et al* (2002) reported additive type of gene action for yield related traits in different wheat populations. Similar type of contradictions in other past studies were reported by Abdel-Sabour *et al* (1996) and Khan *et al* (2009) which could be attributed to residual heterozygosity among the parents. Positive correlation ($r = 0.868$) of $V_r + W_r$ with parental genotypes reflected recessive gene control for spike length (Fig. 3b). Moreover, non-significant deviation of regression line depicted absence of non-allelic interaction for spike length.

3.4. Grain spike⁻¹

Hayman's genetic architecture revealed that over dominance gene action was mainly responsible for grains per spike as regression line intercepted the V_r -axis well below the origin (Fig. 4a). Majority of wheat cultivars were found in cluster near origin which indicated an excess of dominant genes for grains per spike in all studied cultivars. However, the landrace Khattakwal being farthest from origin exhibited excess of recessive genes for said trait. When V_r


+ W_r values were plotted against the parental values, negative correlation ($r = -0.96$) was observed which reflected dominant genes control for managing grains per spike (Fig. 4b). Similar results were reported by Khan and Ali (1998) and Sangwan and Chaudhry (1999) in their studies for tillers plant⁻¹, grains spike⁻¹ and grain yield. Moreover, absence of non-allelic interactions was evident from significant deviation of regression line from unit slope.

3.5. Biological yield (kg ha⁻¹)

Like other parameters, the biological yield in parental genotypes and their F_2 populations was under the control of over-dominance as the regression line cuts V_r -axis below the origin (Fig.5a). Cultivars Tatara-96, Iqbal-2000, Ghaznavi-98 and Wafaq-2001 being closer to origin possessed maximum dominant genes while genotypes Inqalab-91 and Margalla-99 being farthest from origin, possessed large number of recessive genes for biological yield. Moreover, cultivars Tatara-96, Ghaznavi-98, Inqalab-91 and Margalla-99 were closer to regression line than Iqbal-2000, Khattakwal, Wafaq-2001 and Takbeer and thus least affected by the environment (Fig.5a). Negative correlation ($r = -0.39$) among parental genotypes and V_r+W_r values suggested that inheritance pattern for this trait was controlled by dominant genes (Fig. 5b). Over-dominance type of gene action for biological yield and grain yield was also reported by earlier researchers like Asif *et al* (2000), Biranvand *et al* (2013) and Pall and Kumar (2009). However, Salehi *et al* (2014) observed that average degree of dominance was less than unity and therefore, suggested partial dominance effects for this trait in wheat genotypes.

4. CONCLUSION

Over-dominance was mainly responsible for inheritance of studied traits among F_2 wheat populations which indicated that selection would be least effective for improvement in earlier generations. Wheat cultivars Takbeer, Wafaq-2001, Ghaznavi-98 and Iqbal-2000 possessed maximum dominant genes for plant height while Iqbal-2000, Wafaq-2001 and Tatara-96 for days to heading. Similarly Tatara-96 and Ghaznavi-98 exhibited maximum dominant genes for spike length, Margalla-99 for grains spike⁻¹ while Iqbal-2000, Ghaznavi-98, Tatara-96 and Wafaq-2001 for biological yield.



172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191

192
193

194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211

Fig. 1(a).

Fig. 1(b).

Tat, Waf, Tak, Iqb, Mar, G-98, K-wal and Inq = Tatara-96, Wafaq-2001, Takbeer, Iqbal-2000, Margalla-99, Ghaznavi-98, Khattakwal and Inqalab-91, respectively.

Fig. 1(a & b). Wr-Vr & Wr+Vr/P graphs for days to heading among F₂ wheat populations at Peshawar




Fig. 2(a).




Fig. 2(b).

Tat, Waf, Tak, Iqb, Mar, G-98, K-wal and Inq = Tatara-96, Wafaq-2001, Takbeer, Iqbal-2000, Margalla-99, Ghaznavi-98, Khattakwal and Inqalab-91, respectively.

Fig. 2(a & b). Wr-Vr & Wr+Vr/P graphs for plant height among F₂ wheat populations at Peshawar




Fig. 3(a).




Fig. 3(b).

Tat, Waf, Tak, Iqb, Mar, G-98, K-wal and Inq = Tatara-96, Wafaq-2001, Takbeer, Iqbal-2000, Margalla-99, Ghaznavi-98, Khattakwal and Inqalab-91, respectively.

Fig. 3(a & b). W_r - V_r & W_r+V_r/P graphs for spike length among F_2 wheat populations at Peshawar




Fig. 4(a).




Fig. 4(b).

Tat, Waf, Tak, Iqb, Mar, G-98, K-wal and Inq = Tatara-96, Wafaq-2001, Takbeer, Iqbal-2000, Margalla-99, Ghaznavi-98, Khattakwal and Inqalab-91, respectively.

Fig. 4(a & b). W_r - V_r & W_r+V_r/P graphs for grains spike⁻¹ among F_2 wheat populations at Peshawar




Fig. 5(a).




Fig. 5(b).

Tat, Waf, Tak, Iqb, Mar, G-98, K-wal and Inq = Tatara-96, Wafaq-2001, Takbeer, Iqbal-2000, Margalla-99, Ghaznavi-98, Khattakwal and Inqalab-91, respectively.

Fig. 5(a & b). Wr-Vr & Wr+Vr/P graphs for biological yield among F₂ wheat populations at Peshawar

276 REFERENCES

- 277 Abdel-Sabour, M.S., A.M. Hassan, A.A. Abdel-Shafi, H.S. Sherif, and A.A. Hamada. Genetic
278 analysis of diallel cross in bread wheat under different environment conditions in Egypt.
279 2. F₂ and parents. Indian J. Genet. Plant Breed. 1996. 56(1): 49-61
- 280 Akram, Z., S.U. Ajmal, M. Munir and G. Shabbir. Genetic determination of yield and related
281 traits in bread wheat. Sarhad J. Agric. 2008. 24(3): 431-438
- 282 Ali Z., and A.S. Khan. Combining abilities studies of some morpho-physiological traits in bread
283 wheat (*Triticum aestivum*. L). Pak. J. Agric. Sci. 1998. 35: 1-4
- 284 Asif, M., I. Khaliq and M.A. Chowdhry. Genetic analysis for some metric traits in hexaploid
285 wheat. Pakistan J. Biol. Sci. 2000. 3(3): 525-527
- 286 Biranvand, H.P., E. Farshadfar and H. Sabakhpour. Gene action of some agronomic characters
287 in chickpea under stress and non-stress conditions. Asian J. Exp. Biol. Sci. 2013. 4(2):
288 266-272
- 289 Hayman, B.I. The theory and analysis of diallel crosses. Genet. 1954a. 39: 789-809
- 290 Hayman, B.I. The analysis of variance of diallel crosses. Biomet. 1954b. 10: 235-245
- 291 Ivanovska S., C. Stojkovski, and L. Marinkovic. Inheritance mode and gene effect on spikelets
292 number per spike in wheat. Macedonia-Agriculture sector Review. 2000. 47(1/2): 1-8
- 293 Jadoon, S.A., F. Mohammad, H. Ullah and I.H. Khalil. Gene action for pre and post-harvest
294 traits in F₂ wheat populations. Q. Science Connect. 2012. 2-5
- 295 Jones, R.M. Analysis of variance of half diallel table. Heredity. 1965. 20: 117-121.
- 296 Joshi S.K., S.N. Sharma, D.L. Singhanian and R.S. Sain. Combining ability in F₁ and F₂
297 generations of diallel cross in hexaploid wheat (*Triticum aestivum* L. em. Thell).
298 Hereditas-Lund. 2004. 141(2): 115-121
- 299 Khan A.S. and Z. Ali. General and specific combining ability estimates for plant height and some
300 yield components in wheat. Pak. J. Agric. Sci. 1998. 35: 1-4
- 301 Khan N.U., G. Hassan, K.B. Marwat, Farhatullah, M.B. Kumbhar, A. Parveen, U. Aiman, M. Z.
302 Khan and Z.A. Soomro. Diallel analysis of some quantitative traits in *Gossipium hirsutum*
303 L. Pakistan J. Bot. 2009. 41(6): 3009-3022
- 304 Kutlu, I. and M. Olgun. Determination of genetic parameters for yield components in bread
305 wheat. Int. J. Biosci. 2015). 6(12): 61-70
- 306 Mishra P.C., T.B. Singh and D.P. Nema. Combining ability analysis of grain yield and some of
307 its attributes in bread wheat under late sown condition. *Crop Research Hisar*. 1994.7(3):
308 413-423
- 309 Navabi. A., R.P. Singh, J.P. Tewari and K.G. Briggs. Inheritance of high level of plant resistance
310 to stripe rust in five spring wheat genotypes. Crop Sci. 2004. 44(4): 1156-1162.

311 Pall, D. and S. Kumar. Genetic analysis of forage yield and other traits in barley (*H. vulgare* L.).
312 Barley Genet. Newsletter. 2009. 39: 13-19

313 PBS-2015-16. Pakistan Bureau of Statistics, Government of Pakistan.

314 Rabbani, G., M. Munir, S. U. K. Ajmal, G. Shabbir and A. Mahmood. Inheritance of yield
315 attributes in bread wheat under irrigated and rainfed conditions. Sarhad J. Agric. 2009.
316 25(3): 429-439.

317 Salehi, S., S. Gholami, A. Rahmati and A.R. Golparvar. Combining ability of biological yield and
318 harvest index in diallel cross of wheat cultivars under drought and non-drought stress
319 condions. Agric. Conspec. Sci. 2014. 79(4): 221-226

320 Sangwan, V.P. and B.D. Chaudhary. Diallel analysis in wheat (*Triticum aestivum* L.). *Annals of*
321 *Biology Ludhiana*. .1999. 15(2): 181-183

322 Singh H., S.N. Sharma, R.S. Sain, and D.L. Singhania. The inheritance of production traits in
323 bread wheat by diallel analysis. Sabrao J. Breed. Gen. 2003. 35(1): 1-9

324 Singh S.P., L.R. Singh, V.K. Yadav, S. Geeta, R. Kumar, P.B. Singh and G. Singh. Combining
325 ability analysis for yield traits in bread wheat. (*Triticum aestivum* L.). Prog. Agric. 2002.
326 2(2): 119-121

327 Steel R.G.D., J.H. Torrie and D.A. Dickey. Principals and procedures of statistics: A Biometrical
328 Approach. 1997. 3rd edition. Mc Graw Hill Book Co., Inc., New York, USA

329 Zwart, R.S., J.P. Thompson, and I.D. Godwin. Genetic analysis of resistance to root-lesion
330 nematode in wheat. Plant Breed. 2004. 123(3): 209-212

331