

Original Research Article

EFFICACY OF ANACARDIUM OCCIDENTALE AND CYMBOPOGON CITRATUS AS BIOPESTICIDES AGAINST MAIZE WEEVILS (*SITOPHILUS ZEAMAI* MOTSCH.) ON STORED MAIZE

ABSTRACT

An experiment on the insecticidal potency of *Anacardium occidentale* and *Cymbopogon citratus* against maize weevils (*Sitophilus zeamais* Motsch.) on stored maize was evaluated at 1.0g, 1.5g and 2.0g doses per 20g maize grain in 300ml rearing jars. Toxicity in terms of mortality, oviposition, F1 generation emergence, grain damage, and grain weight loss were all noted. Data collected was subjected to Analysis of Variance (ANOVA) using SPSS version 16.0 at 5% level of significance ($P>0.05$). When compared with the control (untreated maize), significant difference ($P>0.05$) in the potency of the biopesticides was observed in the parameters measured. *A. occidentale* performed significantly lower than *C. citratus* as it recorded higher (18.75 ± 9.68) grain damage at 1.0g dose, as well as grain weight loss (0.58) at 1.0g and 1.5g treatment doses respectively. This suggests that although the plant products were tested to possess insecticidal potency against the maize weevils, *C. citratus* performed significantly ($P>0.05$) better than *A. occidentale* in the control of *S. zeamais* on stored maize. Hence, it could be used sustainably in the control of insect pests on stored products

KEYWORDS: *Anacardium occidentale*, Biopesticide, *Cymbopogon citrates*, Maize *Sitophilus zeamais*

INTRODUCTION

Maize (*Zea mays*) is a cereal crop widely cultivated in different parts of the world. Worldwide, about 785 million tons of maize is produced each year, with United States being the largest producer, accounting for about 42% of the overall production [1]. Maize is an important food crop in the tropics, subtropics and temperate regions where they are grown [2]. In Nigeria, and in Adamawa State, maize is one of the most important economic crops grown [3]. But unfortunately, maize production in most parts of Nigeria is restricted to only the rainy months of

the year. This however necessitates the need to store the maize grains throughout the dry months of the year.

However, the storage of maize grains and its maximum utilization is seriously affected by the activities of pests [4]. The major insect pest is *Sitophilus zeamais* Motsch (Coleoptera: Curculionidae). To enhance productivity of maize in the field and reduced its loss in storage, the control of insect pests activities is needed, including the use of chemical insecticides. The use of insecticides is very effective and reliable, and can prevent loss of maize yield. They act quickly, making them suitable for use in emergency situations [2]. But unfortunately, the intensive use of insecticides has caused problems such as residual effect and the effect on non-target organisms. Hence, safe alternatives which are also effective, non-costly and biodegradable, are needed for the control of these pests against stored product pests, especially *S. zeamais* [5]. Farmers and researchers have been trying to control stored product pests like *S. zeamais* using plant products (biopesticide) which are readily available, cheap and environmentally friendly [6, 7].

Anacardium occidentale is an evergreen tree in the family Anacardiaceae and has both medicinal and economic values [8]. *Cymbopogon citratus* is utilized for its numerous uses. It is used as fragrance, for perfume and also a source as vitamin A and E [9]. *Cymbopogon citratus* oil is used deodorants, waxes, polishes, detergent and insecticides [10].

In this study, the insecticidal potentials of leaf powders of *Anacardium occidentale* and *Cymbopogon citratus* were evaluated against the *S. zeamais* with a view of developing biopesticides against stored product pests.

MATERIALS AND METHODS

Study area

This study was carried out in Mubi town of Adamawa state, Nigeria. Mubi is located between latitude 10°12'N and longitude 13°10'E. The climate is tropical with average temperature of about 15-42°C in dry season and relative humidity 10-45% [11].

Collection of materials

Samples of lemon grass (*Cymbopogon citratus*) and cashew leaves (*Anacardium occidentale*) were obtained around the Botanical Garden, Adamawa State University, Mubi. They were cupboard-dried at room temperature in the laboratory of Biological Science, Adamawa University for four days and were subsequently pulverized into fine powder using electric blender. Synthetic chemical (Rambo-permethrin 0.60%) was obtained in an agro-chemical store in Mubi market and was used as the positive control. Clean maize cobs was also obtained from a farmer in Mubi and shelled to obtain the grains.

Insect culture

Parent stock of *S. zeamais* was obtained from infested maize grains obtained from a store in Mubi. The adult weevils obtained from the parent stock were transferred onto a clean maize grain in 500cm³ rearing jar which was allowed to stand for 48 hours. Thereafter, the insects were discarded and the exposed maize was maintained under laboratory conditions for adult emergence of *S. zeamais* [12]. This helped in raising *S. zeamais* adults of the same size and age that were used to carry out the experiment.

Experimental setup

Before the experiment was set up, the maize grains were oven-dried to a constant weight in an oven at 40-45°C for two hours and were later air-dried in the laboratory to prevent moldiness for about an hour [12, 13]. 20g of the maize grains were placed in each of the treatment cups. Three treatment doses of 1.0g, 2.0g and 3.0g were constituted and replicated 4 times. These were added individually to the 20g of maize grains in the treatment jars and mixed thoroughly to ensure a uniform coating of the grain by the treatment samples. Thereafter, 10 newly emerged adult *S. zeamais* obtained from the cultured jar were introduced into the experimental jars at the same time. The content of the jars were covered with muslin clothes with the aid of a rubber band to prevent the insects from escaping and also to allow for ventilation.

Data collection

Mortality:

Mortality count was recorded at 24 hours post exposure intervals of four days. The content of each experimental jar was emptied gently on a silver tray in the laboratory. Using a broom stick, the insects were touched one after the other. The inability of any of the insects to

respond to one touch indicated a death individual. The dead insects were subsequently removed and the remaining contents and the live insects were put back in the jar.

Oviposition:

Oviposition was determined using the acid fuchsin staining method. 10 grains of maize from each jar was randomly picked on the 10th day, immersed in warm water for two to three minutes, drained and subsequently put in 0.5% acid fuchsin stain for two to five minutes. The grains were rinsed in water and were determined for cherry red gelatinous egg plugs. The number of egg plug noticed on the 10 grains was extrapolated for the entire jar using an average number of 82 grains per jar [14].

$$\text{Oviposition} = \frac{\text{Average number of maize grain} \times \text{Number of eggs laid}}{\text{Number of maize grain picked randomly}}$$

F1 generation emergence:

The F1 progeny emergence was observed within 4-5 weeks after exposure to treatments in the course of the experiment. The newly emerged adult *S. zeamais* was counted and recorded.

Grain damage:

Ten maize grains were selected randomly from each plastic jar and were noted for grain damage. Maize grains observed with holes, cracks and any form of abrasion were considered as damaged grains. The number of damages was noted on 10 randomly selected maize grains, and was extrapolated for the entire jar using an average number of 82 grains per jar.

Weight loss:

The weight of the maize grain in each experimental jar was initially noted using a weighing balance prior to the experimental setup. Thereafter, the final weight of the grain was also noted after the experiment. Therefore, weight loss was determined by subtracting the final weight from the initial weight, as shown in the formula below:

$$\text{Weight loss} = \text{initial weight} - \text{final weight}$$

Data analysis:

Data collected were subjected to Analysis of Variance (ANOVA), and the means were separated at $P>0.05$ level of probability using Least Significant Difference (LSD).

RESULTS

Table 1 shows the toxicity effect of treatments (*C. citratus* and *A. occidentale*) in terms of mortality of *S. zeamais* reared on maize grain. Overall, the mortality counts on various treatments and the control (untreated) experiment were significantly different ($P>0.05$). Although the mortality was spread across the four days of exposure in all the treated jars, there was no mortality recorded in the control experiment throughout the exposure period. At the end of the four days of exposures, there was a significant difference ($P>0.05$) in the mortality recorded across the treatments and the control. Among the plant treatments, *A. occidentale* appeared to have recorded a significant higher mortality than the *C. citratus* (Table 1).

Table 2 shows effect of the plant treatments in suppressing egg laying and adult F1 progeny emergence of *S. zeamais*. The results revealed that there was a significant difference ($P>0.05$) in both the egg laying and the number of F1 generation emergence between the treatments and control (untreated) experiment. The treatments significantly ($P>0.05$) reduced the number of eggs laid by the adult *S. zeamais* and the number of adult emergence. The number eggs laid was significantly ($P>0.05$) lower in grains treated with *A. occidentale* (12.30) at 1.0g dose, while the highest (18.45) was recorded in *C. citratus* at 1.0g dose and *A. occidentale* at 1.5g dose respectively. As expected, the synthetic insecticide (Rambo-permethrin 0.60%) significantly ($P>0.05$) reduced the number of eggs laid to as low as 2.05 in 1.0g and 2.0g treatment dose, and no adult emerged (0.00) for F1 generation.

The effect of treatments in the protection of maize grain from damage and weight loss as a result of *S. zeamais* activities is shown in Table 3. There was a significant difference ($P>0.05$) in grain damage and subsequent weight loss by the maize grains as a result of the insect activities, when compared with the control (untreated) experiment. Here, the grains treated with *A. occidentale* significantly ($P>0.05$) recorded higher damages (18.75 ± 9.68) at 1.0g dose. The subsequent weight loss recorded was a clear reflection of the result on grain damage, as maize treated with *A. occidentale* significantly ($P>0.05$) recorded higher (0.58) at 1.0g and 1.5g

139 treatment doses respectively. As expected, the synthetic insecticide (Rambo) at all treatment
 140 doses significantly reduced grain damage as well as the subsequent weight loss by the grains.

141 **TABLE 1:** Mortality of *Sitophilus zeamais* reared on maize grain

Treatment	Dose (g)	Days of Exposure (Mean±SD)				Total
		1	2	3	4	
Control	0.00	0.00±0.00 ^a	0.00±0.00 ^a	0.00±0.00 ^a	0.00±0.00 ^a	0.00±0.00 ^a
<i>C. citrates</i>	1.0	0.00±0.82 ^a	1.00±0.82 ^b	0.25±0.50 ^a	0.00±0.00 ^a	1.25±1.26 ^{ab}
	1.5	0.00±0.00 ^a	0.25±0.50 ^{ab}	0.00±0.00 ^a	0.00±0.00 ^a	0.25±0.50 ^a
	2.0	1.00±1.15 ^a	0.25±0.50 ^{ab}	0.50±1.00 ^a	0.00±0.00 ^a	1.75±1.71 ^{ab}
<i>A. occidentale</i>	1.0	1.00±1.15 ^a	0.00±0.00 ^a	0.50±0.58 ^b	0.00±0.00 ^a	1.50±1.29 ^{ab}
	1.5	1.00±2.00 ^a	0.25±0.50 ^a	0.00±0.00 ^a	0.50±1.00 ^a	1.75±2.06 ^{ab}
	2.0	0.25±0.50 ^a	0.75±0.96 ^a	0.00±0.00 ^a	0.25±0.50 ^a	1.25±0.50 ^{ab}
Rambo	1.0	3.50±0.58 ^b	4.25±1.26 ^b	2.00±0.82 ^b	0.25±0.50 ^a	10.00±0.00 ^c
	1.5	2.50±1.00 ^{ab}	6.00±1.83 ^b	1.25±0.96 ^b	0.25±0.50 ^a	10.00±0.00 ^c
	2.0	4.00±3.39 ^b	4.75±2.63 ^b	1.00±0.82 ^{ab}	0.25±0.50 ^a	10.00±0.00 ^c

142 Values are means of four replicates. Means carrying the same superscript alphabet along the
 143 columns are not significantly different ($P>0.05$).

144

145 **TABLE 2:** Oviposition and F1 progeny emergence of *S. zeamais* reared on maize grain.

Treatment	Dose (g)	Parameters (Mean±SD)	
		Number of eggs laid	F1 generation Emerged
Control	0.0	36.9±10.59 ^c	4.50±1.29 ^c
<i>C. citrates</i>	1.0	18.45±14.00 ^b	1.50±0.58 ^b
	1.5	16.40±20.00 ^b	2.00±2.16 ^b
	2.0	18.45±12.30 ^b	2.50±1.29 ^b
<i>A. occidentale</i>	1.0	12.30±19.52 ^{ab}	2.25±1.26 ^b
	1.5	18.45±21.57 ^b	1.75±2.87 ^b
	2.0	20.50±4.74 ^b	2.50±2.52 ^b
Rambo	1.0	2.05±4.10 ^a	0.00±0.00 ^a
	1.5	6.15±7.85 ^a	0.00±0.00 ^a
	2.0	2.05±4.10 ^a	0.00±0.00 ^a

146 Values are means of four replicates. Means carrying similar alphabets along the columns are
 147 not significantly different ($P>0.05$).

148

149

150 **TABLE 3:** Damages and weight loss of maize grain.

Treatment	Dose (g)	Parameters (Mean±SD)	
		Grain Damage	Weight Loss (g)
Control	0.0	22.50±13.69 ^b	1.00±0.18 ^b
<i>C. citrates</i>	1.0	15.00±6.12 ^{ab}	0.28±0.22 ^a
	1.5	11.25±4.33 ^{ab}	0.45±0.25 ^a
	2.0	11.25±4.33 ^{ab}	0.23±0.13 ^a
<i>A. occidentale</i>	1.0	18.75±9.68 ^b	0.58±0.26 ^a
	1.5	13.12±3.75 ^{ab}	0.58±0.33 ^a
	2.0	13.12±7.18 ^{ab}	0.55±0.13 ^a
Rambo	1.0	7.50±6.12 ^a	0.33±0.10 ^a
	1.5	5.62±7.18 ^a	0.30±0.16 ^a
	2.0	7.50±6.12 ^a	0.25±0.17 ^a

151 *Values are means of four replicates. Means carrying the same superscript alphabets along the*
 152 *columns are not significantly different (P>0.05).*

153 DISCUSSION

154 In this study, two different samples of biopesticides (*Cymbopogon citratus* and
 155 *Anacardium occidentale*) were tested for the insecticidal activities on maize weevils (*Sitophilus*
 156 *zeamais*). All the treatments caused a significant mortality of adult *S. zeamais* than the control
 157 (untreated), where no mortality was recorded throughout the four days of exposure, confirming
 158 the effectiveness of the plant products (biopesticides) on *S. zeamais*. The order of their efficacy
 159 is: Rambo>*A. occidentale*>*C. citratus*. The results of this study corroborate with the findings of
 160 Dike and Mbah [15] where lemon grass products significantly protected cowpea grains in storage
 161 from insect infestation. The powder of *A. occidentale* has been reported as protectant against
 162 different storage insect pests [16]. In a similar study on the efficacy of plant products on stored
 163 maize, Wahedi [7] reported the insecticidal effect of neem seed powder on adult *S. zeamais*
 164 reared on maize grains in Mubi.

165 The biopesticides were able to reduce F1 emergence of *S. zeamais* on maize grain
 166 significantly (P>0.05). *C. citratus* and *A. occidentale* showed some insecticidal activities by
 167 reducing the number of eggs laid as well as F1 adult emergence, when compared with the control
 168 (untreated) experiment. This further confirms the insecticidal properties in the plant products.
 169 This agrees with the study conducted by Mba and Okoronkwo [2] and Wahedi [7] who reported

the efficacy of plant products in reducing egg laying as well as F1 progeny emergence of *S. zeamais*.

Similar result was reported in the grain damage and weight loss. The treatments showed some insecticidal activities by significantly protecting the maize grains from damages and subsequent weight loss possibly as a result of the activities of *S. zeamais*. This also agrees with the study performed by Wakole [17] and Wahedi [7], who reported plant products effectively protected maize grain from damage as well as weight loss.

CONCLUSION

In conclusion, the overall results have confirmed the effectiveness of *A. occidentale* and *C. citratus* as biopesticides for the control of adult *S. zeamais*. *A. occidentale* performed significantly better than *C. citratus* based on the various parameters tested.

REFERENCES

- [1] Ibrahim A, Zalkuwi J, Yusuf C. Technical efficiency of Maize production in Mubi North Local Government Area of Adamawa State, Nigeria. Int. J. of Innovative Research and Studies. 2014; 3 (1):58-67.
- [2] Mbah OI, Okoronkwo MO. An assessment of two plant product efficacy for the control of maize weevil (*Sitophilus zeamais* Motschulsky) in stored maize. African Journal of Agricultural Research. 2008; 3(7) 494-498.
- [3] Zalkuwi J, Dia YZ, Dia RZ. "Analysis of Economic Efficiency of Maize Production in Ganye Local Government Area Adamawa state, Nigeria". Nassarawa. 2010; www.sciencepub.net/report
- [4] Arannilewa ST, Odeyemi OO, Adedire CO. Effects of medicinal plant extracts and powder on the maize weevils, *Sitophilus zeamais* Motsch. (Coleopteran: Curculionidae). Annals Agric. Sci. 2002; 3:1-10.
- [5] Okonkwo EU, Okoye WI. The efficacy of four seed powders and essential oils as protectants of cowpea and maize grains infestation by *Callosobruchus maculatus* and *Sitophilus zeamais* in Nigeria. Int. J. Pet. Manag. 1996; 480pp.

- 196 [6] Ibe AC, Nwufo MI. Production of stored maize seeds against maize weevil (*Sitophilus*
197 *zeamais*) using local plant materials. In salako F.K, Lagoke S.T.O, Aina A.B.J, Enuubetine D,
198 Dipeolu A.O (eds). Proceedings of the 35th Annual conference. The Agric. Soc. of Nigeria.
199 2001; 368-370pp.
- 200 [7] Wahedi JA. Laboratory evaluation of neem (*Azadirachta indica*) seed powder and seed oil
201 for the control of *Sitophilus zeamais* on stored maize. Adamawa State University Journal of
202 Scientific Research. 2012; 2(2): 110-115.
- 203 [8] Rajesh KV, Sumathi CS, Balasubramanian V. Elementary chemical profiling and Antifungal
204 properties of Cashew Nuts. Botany Research Intl. 2009; 2:253-257.
- 205 [9] Kannan C, Sumathi S, Balasubramanian V. Elementary chemical profiling and Antifungal
206 properties of Cashew Nuts. Botany Research Intl. 2009; 2:253-257.
- 207 [10] ASNAPP. Agribusiness in sustainable natural African products (South Africa), Newsletter.
208 2008; Online: www.asnapp.org.
- 209 [11] Adebayo AA, Tukur AL. "Adamawa State in Maps" Paraclete Publisher Yola. 1999; 27-
210 31pp.
- 211 [12] Arannilewa ST, Ekrakene T, Akinneye JO. Laboratory evaluation of four medicinal plants
212 as protectants againsts the maize weevil, *Sitophilus zeamais* (Motsch.). African Journal of
213 Biotechnology. 2006; (21): 2032-2036.
- 214 [13] Adedire CO, Ladije L. Toxicity and oviposition differences of some plant extracts on
215 cowpea storage bruchid. *Callosobruchus maculatus*. J. Pl. Dis. Prot. 1999; 106: (6):647-653
- 216 [14] Wahedi JA, Ande AT, Fatoba PO. Biocidal activities of some tropical moss powders against
217 *Sitophilus zeamais* Motch. (Coleoptera: Curculionidae). Global Journal of Bio-Science and
218 Biotechnology. 2013; 2(3): 386-389.
- 219 [15] Dike MC, Mbah OI. Evaluation of lemon grass (*Cymbopogon citratus*) Products in the
220 control of *Callosobruchus maculatus* on stored cowpea. Nig. J. Pl. Prot. 1992; 4:88-91

- 221 [16] Afzal M, Nasir Z, Bashir MH, Khan BS. Analysis of list plant resistant in some genotypes
222 of maize against *Chilo partellus* (Swinhoe) (Lepidoptera: Pyralidae). Pakistan Journal Botany.
223 2009; 41:421-428.
- 224 [17] Waktale S. Effect of selected botanicals and local seed storage practices on maize insect
225 pests and health of maize seeds in Jimma Zone. Singapore Journal of Scientific Research. 2014;
226 4:19-28.