

EFFECT OF POULTRY MANURE AND UREA ON SOIL CHEMICAL PROPERTIES, NODULATION AND YIELD OF GROUNDNUT (*Arachis hypogaea*) IN AKANU IBIAM FEDERAL POLYTECHNIC, UNWANA AFIKPO EBONYI STATE

ABSTRACT

A Green House study was carried out at the Teaching and Research Farm, Department of Horticulture and Landscape Technology to investigate the influence of the integrated application of poultry manure and urea on soil nutrient properties, nodulation, growth and yield of groundnut (*Arachis hypogaea*) in Akanu Ibiam Federal Polytechnic, Unwana Southeastern Nigeria. The experiment was laid out into Complete Randomized Design with different levels of poultry manure (0, 1 and 2 tons ha⁻¹) and urea (0, 1 and 2 tons ha⁻¹) replicated three times. Results showed that poultry manure applied alone or combined with urea significantly ($P < 0.05$) improved almost all the soil nutrient properties. Relative to control, best improvements on soil fertility indices such as soil pH, available phosphorus, total nitrogen, organic carbon, total exchangeable acidity (TEB) and effective cation exchange Capacity (ECEC) were obtained in the pot treated with poultry manure 2 tons ha⁻¹ and urea 1 ton ha⁻¹. Similarly, poultry manure integrated with urea significantly ($P < 0.05$) improved number of leaves and number of pods. Application of urea without poultry manure increased number of leaves better than the lone application of poultry manure, while poultry manure and its integration with urea had significant improvement on the number of pods compared with alone application of urea. The maximum improvement on the number of leaves and pod was obtained when poultry manure was applied at 2 tons ha⁻¹ integrated with urea at 1 ton ha⁻¹. It was also observed that the number of nodules increased with poultry manure application but decreased with application of urea. Integration of urea with poultry manure is therefore recommended to improve soil fertility and grain yield of groundnut. Large amount of urea should not be used in groundnut production; rather poultry manure should be incorporated to enhance nodulation and nitrogen availability in soils of Unwana Southeastern Nigeria.

Key words: Poultry manure, urea, nodulation and groundnut.

INTRODUCTION

Groundnut is an edible legume which serves as one of the main sources of income for small holders and commercial farmers. It is mainly grown for its edible seed, and is widely grown in tropics and subtropical regions as grain legume. The seeds contain sufficient amount of protein (19%), carbohydrate (63%), fat (6.5%) and essential amino acid (Lineman 1988; Brough and Azam Ali, 1992). It is cultivated at altitude ranging from 0-150m and is generally intercropped with cereals, cassava, yam and cow pea. It fixes atmospheric nitrogen, and hence improves the soil nitrogen that play its role as a restorative crop (Mukurumbia, 1985). In Nigeria, it is cultivated in the northern regions, characterized by yellow ferratitic soil (Duchaufor, 1976). Since the nitrogen requirement for the other crop (intercropped) is met through nitrogen fixation, therefore the interest has been developed in cultivation of groundnut for its tremendous potentials both as food and in soil fertility maintenance (Somasegaran *et al*, 1992).

Crop production in Nigeria is hindered by several factors such as low soil fertility as well as restricted access to mineral fertilizer (Nyanabgara and Nyaguambo, 2010). In Nigeria, farmers are always faced with the problem of soil fertility decline which has been considered the most important biophysical constraints to crop yield and productivity (Odendo *et al.*, 2004).

Developing soil fertility management options for increasing productivity of staple food crops is a challenge in most part of Nigeria, where soils are constrained by the major mineral element such as potassium, nitrogen and phosphorus deficiencies (Jemo *et al*, 2010). Nitrogen, Phosphorus and potassium are among the limiting nutrients for legumes production (Christianson and Viek, 1991). Adequate supply of nitrogen is beneficial for carbohydrates and protein metabolism, promoting cell division and cell enlargement (Shehu *et al*, 2010).

Poultry manure promotes transformation and mineralization of less labile inorganic and organic phosphorus into labile phosphorus in the rhizosphere, which results in higher root phosphorus concentration and higher total phosphorus uptake by plants (Waldrip *et al.*, 2011). Therefore, poultry manure could be a valuable fertilizer and could serve as a suitable alternative to chemical fertilizer in Nigeria (Eneje and Azu, 2009).

Apart from the addition of both mineral and organic nitrogen fertilizers, there is another viable and ecologically friendly strategy for improving nitrogen content of soils. This process which is called biological nitrogen fixation is specifically reserved for leguminous crops (Brady and Weil 2008). A relatively small amount of nitrate or ammonia is produced by lightening, some are produced industrially by the Haber-Bosch process, but the major conversion of Nitrogen (N₂) to ammonia and hence protein is achieved by microorganisms that inhabit the root nodules of legumes in the process of nitrogen fixation (Sorensen and Sesitch, 2007). The process of biological nitrogen fixation in legume nodules is an important soil fertility improvement strategy especially nitrogen economy in many tropical ecosystem (Frankow-Lindberg, and Dahlin, 2013). Walley *et al.* (1996), stated that most of the nitrogen eventually returns to the soil when vegetation of legume dies and decomposes and this can be made use of by the upcoming crop

Even though several literatures have stated that leguminous crops can derive their maximum nitrogen requirement from biological nitrogen fixation and therefore require no external nitrogen fertilizer application, there is still need to practically investigate the effect of different fertilizer sources on nodulation of groundnut. This study was therefore designed to evaluate the effect of poultry manure and urea on soil chemical properties, nodulation and yield of groundnut (*ArachisHypogea*) in Unwana, South Eastern Nigeria.

MATERIALS AND METHODS

The study was carried out at Horticultural Farm of AkanuIbiam Federal Polytechnic Unwana. Unwana is located within the South Eastern ecological zone of Nigeria and lies at latitude 5° 48N and longitude 7° 55E (Njoku and Okpara, 2006). The climatic condition and vegetation types are generally humid tropical rain forest with mean annual rainfall of about 3,500mm and mean daily temperature of 21°C to 32 °C (Odurukwe *et al.*, 1995).

Soils sample which was collected from 0 – 20cm depth, was air dried, sieved through 2mm sieve after which 5kg of sub-samples were weighed into 10 litre capacity buckets perforated at the bottom.

Poultry manure and urea were added to each pot. The experiment was laid out into Complete Randomized Design with different levels of poultry manure (0,1 and 2 tons ha⁻¹) and urea (0, 1 and 2 tons ha⁻¹) replicated three times. Four groundnut seeds were sown in each pot after one month of treatment application to give time for the

decomposition of the poultry manure and later thinned down to two seedling per bucket after germination. Adequate watering and weeding were observed throughout the growing period.

DATA COLLECTION

At full bloom (50 days after planting) number of nodules per plant was determined from one plant in each pot. The seedlings was carefully uprooted and washed, all the soil particles was removed and then the nodules were counted (Faizah, 1980; Osodeke, 2002). At full maturity, the other plant in each pot was uprooted and both the number of leaves and pods were obtained by counting.

Post-harvest soils samples were collected from each pot and analyzed for routine chemical properties according to standard laboratory procedures. Particle size distribution was determined using the Bouyocous hydrometer method as describe by Benton (2001). The pH was determined in soil to water and soil to CaCl_2 at a ratio 1:2 soil water and soil CaCl_2 respectively using glass electrode P^{H} meter (Udo *et al.*; 2009). Organic carbon was determined by the wet oxidation method according to Pansu and Gautheyrous (2006) and converted to organic matter by multiplying by 1.792. Available P was determined using the Bray II method of Bray and Kurtz (1945) as described by Udo *et al.* (2009). The total nitrogen determination was done by the macro Kjeidahl digestion method (Simmone *et al.*, 1994). Exchangeable acidity was determined by the nikel extraction procedure as described by Kamprath (1967). Exchangeable basic cations (K^+ , Ca^{2+} , Mg^{2+} , Na^+) were determined by the ammonium acetate method (Carter and Gregoich, 2008). Ca and Mg in the extract were determined using the atomic absorption spectrophotometer, while K and Na were determined using the flame photometer. Effective cation exchange capacity (ECEC) was obtained by summation of all the exchangeable cations and exchangeable acidity as described by Udo *et al.* (2009). The base saturation was obtained mathematically with

$$\text{BS (\%)} = \frac{\text{total ctions}}{\text{ECEC}} \times \frac{100}{1}$$

STATISTICAL ANALYSES

Data from yield parameters, number of nodulation and soil chemical properties were subjected to analysis of variance (ANOVA) and the means separated using FLSD 0.005.

RESULTS AND DISCUSSIONS

The physical and chemical properties of the soil used for the study is presented in Table 1. The result showed that the sand, silt, and clay percentages were 40.13, 21.56 and 38.31% respectively. The textural class of the soil was a clay loam. These findings are in agreement with the result of Azu *et al.* (2017), who reported a clay – loam soil in Ebonyi State. Soil pH values both in H_2O and CaCl_2 , showed that the soil was acidic, which suggest the need for liming in order to improve the soil productivity. Several other researchers (Onwuka *et al.*, 2007; Eneje and Azu, 2009) have also reported acidity in most soils of South – Eastern Nigeria. Both organic carbon and organic matter moderately low (1.90 and 3.29% respectively). The total nitrogen was low (0.25%) and phosphorus was found at

low concentration (5.62mg/kg) indicating high fixation and hence low phosphorus availability. The value of phosphorus was lower than the critical value of 15mgkg⁻¹ suggested by Osodeke and Ubah (2005). Calcium was relatively high (3.18cmolkg⁻¹). The total exchangeable acidity (TEA) was high (4.01cmol.kg⁻¹) indicating the influence of aluminum and iron oxides (Azu *et al.*, 2017). The effective cation exchange capacity (ECEC) was moderate (8.27cmol/kg) and base saturation (BS) was also relatively moderate (51.51%).

Table 1. Physical and chemical properties of the soil used for the experiment

Properties	values
Sand %	40.13
Silt %	21.56
Clay %	38.31
Texture	clay-loam
pH (H ₂ O)	4.27
pH (CaCl ₂)	3.78
Organic carbon %	1.90
Organic matter %	3.29
Total Nitrogen %	0.25
Available Phosphorus (mg/kg)	5.62
Ca (cmol/kg)	3.18
K (cmol/kg)	0.06
Mg (cmol /kg)	1.00
Na (cmol/kg)	0.02
TEA (cmol/kg)	4.10
ECEC (cmol/kg)	8.27
B.S %	51.51

Most studies conducted across South Eastern Nigeria showed that most soils are acidic, low in soil organic carbon, available N, P, K, Na and some micronutrients (Azu *et al.*, 2017; Eneje and Azu 2009). These nutrient deficiencies can affect not only plants but also soil microbes including rhizobia populations (Brady and Weil, 2008) and their activities such as N₂ fixation (Osodeke, 2002). **Table 2: Results of analyzed poultry manure used for the experiment**

139	Properties	Values
140	pH (H ₂ O)	8.22
141	pH (CaCl ₂)	8.02
142	Av. P (PPM)	0.95
143	Nitrogen (%)	1.88
144	Calcium (%)	5.70
145	Magnesium (%)	1.90
146	Potassium (%)	0.51
147	Sodium (%)	0.069
148	Organic carbon (%)	4.07

149

150 **Effect of poultry manure and urea on soil chemical properties**

151 Result of the effect of poultry manure and urea on soil fertility is shown in Table 3. The application of poultry
152 manure as lone treatments generally increased soil pH both in water and in salt. There were consistent decline in soil
153 pH with increasing urea as lone treatment. This observation may be attributed to the fact that poultry manure being
154 an organic material contains large amount of basic cations which factored in the reduction of soil acidity. Urea,
155 being a mineral fertilizer with no organic component may have been responsible for increased acidity as observed.
156 This report is consistent with the results of other researcher (Azu *et al.*, 2017). The interaction between the poultry
157 manure and urea showed significant (P<0.05) improvement in the soil acidity (Adediran *et al.*, 1999; Azu *et al.*,
158 2017., FAO, 1976) who reported better performance in soil acidity when organic and inorganic fertilizers are
159 combined. The pH values ranged from 6.08 to 3.79 and the highest value was obtained at the treatment combination
160 of 2t ha⁻¹ poultry manure and 1t ha⁻¹ urea.

161 With the exception of ECEC, the lone application of poultry manure was significant (P<0.05) in improvement the
162 soil nutrient indices, but the lone application of urea was only significant in influencing the pH, N and the
163 exchangeable bases. However, the interaction between the two treatments was highly significant (P<0.05) in
164 influencing the pH, total nitrogen, exchangeable bases and exchangeable acidity.

165 **Table 3. Effect of poultry manure and urea on soil chemical properties**

166	M+UR		pH	Aval. P	Org. C	Total N	Exc. Bases	Exc. Acidity	ECEC	BS
167	(t/ha)	H ₂ O	Cacl ₂	(mg/kg)	(%)	(%)	(Cmol/kg)	(Cmol/kg)	(Cmol/kg)	%

168	0 + 0	4.13	3.85	4.68	0.89	0.19	4.17	2.97	7.14	58.40
169	0 + 1	4.00	3.69	4.99	0.76	0.23	5.00	3.23	8.23	60.75
170	0 + 2	3.79	3.44	5.55	0.90	0.29	4.94	3.60	8.54	57.85
171	1 + 0	5.40	4.71	5.63	2.60	0.21	6.63	2.21	8.84	75.00
172	1 + 1	5.64	5.08	6.02	2.60	0.28	8.92	2.33	11.25	79.29
173	1 + 2	4.32	4.24	5.10	1.99	0.39	8.71	2.42	11.13	78.26
174	2 + 0	6.07	5.65	7.15	3.02	0.34	9.76	1.00	10.76	90.71
175	2 + 1	6.08	5.76	7.39	2.93	0.42	10.81	1.09	11.90	90.84
176	2 + 2	5.59	5.09	6.48	2.77	0.44	9.54	1.11	10.65	89.58
177	Mean	5.06	4.59	5.89	2.05	0.31	7.61	2.22	9.83	75.63
178										
179	SD(0.05)U.	0.126	0.0055	0.192	0.127	0.019	0.0022	NS	NS	NS
180	SD(0.05)PM	0.121	0.0203	0.192	0.127	0.019	0.0030	0.00014	NS	2.114
181	SD (U XPM)	0.237	0.334	0.371	0.260	0.038	0.0065	0.0177	0.135	3.008


182

183 Effect of poultry manure and urea on number of nodules

184 With the exception of urea as lone treatments, the number of nodules was observed to significantly ($P < 0.05$) differ
185 among the various treatments and their combinations. As shown in figure 1, the addition of poultry manure as lone
186 treatment had significant ($P < 0.05$) increasing effect on the number of nodules relative to control and the increased
187 was proportional to the rate of amendment. These finding are similar to those reported by Panda *et al.* (2012) who
188 found that poultry manure boosts the effectiveness of rhizobium in cowpea. Consequently, the highest number of
189 nodules (196) was obtained in the pot that had poultry manure at 2t ha^{-1} . This observation can be attributed to the
190 fact that poultry manure in addition to nitrogen and other elements contains large amount of organic phosphates
191 which mineralized and enrich the soil with the availability of phosphorus which increased nodulation of legumes
192 (Osodeke, 2002; Amba *et al.*, 2013; Thedchanamoorthy and Gurung, 1999; Lomoro, 2008; Naveenkurma *et al.*,
193 2015). Therefore the nodules produced by the plant that had 2t ha^{-1} poultry manure was significantly higher than the
194 other treatments. Result also showed that the number of nodules was only higher (110.00) in plant that had 1t ha^{-1}
195 urea compare to control (84.67) and subsequently, the number of nodules decreased with increased rate of urea
196 application (54.50). Nodulation in legumes has been known to require initial nitrogen to boost the process, after
197 which continuous addition of mineral nitrogen may adversely affect the nitrogen fixing organisms. High amount of
198 nitrogen application have been reported to reduce the rate of nodulation, but little quantity have been reported to
199 enhance early vegetative growth without compromising the process of nodulation (Anteneh, 2015 ; Amba *et al.*,

200 2013). Therefore increased application of poultry manure led to corresponding increase in the number of nodules,
 201 while urea application rate was inversely proportional to the number of nodules.

202 **Figure 1. Number of nodules, growth and yield components of groundnut as affected by the addition of**
 203 **poultry manure and urea.**
 204



Treatments (Poultry manure + urea)

Effect of poultry manure and urea on number of Leaves

210 The application of poultry manure and urea both alone and combined treatments had significant ($P < 0.05$) effect on
 211 the number of leaves. Generally, these treatments increased the number of leaves compare to the control. The
 212 application of urea produced maximum number of leaves compare to lone application of poultry manure. This
 213 observation may be due to the quick release of nitrogen by urea in relation to poultry manure which takes longer
 214 time to release its nutrient content (Gunri and Nath, 2012). Nitrogen has been known to improve the vegetative
 215 growth of plants, therefore, the quicker its release the more the leave growth (Brady and Weil, 2008). It was also
 216 observed that the interaction between the two fertilizers had better improvement on the number of leaves compared
 217 to the single application. This report is in accord with the report of Naveenkurma *et al.* (2015). The highest number
 218 of leaves (110.00) was obtained in the plant that had 2t ha^{-1} poultry manure and 2t ha^{-1} urea while the least value
 219 (55.00) was obtained in the plant that had only 1t ha^{-1} urea. The most impressive report on the number of leaves
 220 observed in experimental pots that had different combinations of poultry manure and urea agreed with the recent
 221 emphasis of supplementation of mineral fertilizers with organic manures for better crop performances and
 222 sustainable soil fertility maintenance (Adediran *et al.*, 1999; Azu *et al.*, 2017).
 223 **Effect of poultry manure and urea on number of pod**

224
 225 Number of pods showed significant ($P < 0.05$) response to the application of poultry manure and urea both as single
 226 or combined treatments. Figure 1 showed that the number of pods ranges from 31 to 12. It was observed that the
 227 interaction between the two amendments consistently gave higher number of pods compared to either application
 228 of both materials as lone treatments Naveenkurma *et al.* (2015). This observation can be explained on the fact that,
 229 while nitrogen may be responsible for the leave growth, other nutrient element especially phosphorus which is in

abundance in the poultry manure might have factored more in the seed formation (Brady and Weil, 2008). Consequently, the pot that had 2t ha^{-1} poultry manure and 1t/ha urea had the highest number of pods. Gunri and Nath, (2012) have also reported greater yield pods of groundnut when poultry manure was applied at different rates with a recommended dose of mineral nitrogen fertilizer. Similarly, Lomoro, (2008) reported significant effect of poultry manure combined with mineral fertilizer on the number and weight of groundnut pods.. Previous works on organic and inorganic fertilization on the soil have found to buildup of essential plant nutrients (i.e., N, P, K, S, Ca, Mg, Zn, Fe, Mn, and B) in the soil (Dotaniya *et al.*, 2016; Ogundare *et al.*, 2012; Rezig *et al.*, 2012).

CONCLUSION

In general, organic and inorganic fertilizer applications are relevant to boost the fertility status of soils and effectiveness of *Brady rhizobium* for nodulation and yield of groundnut in degraded sandy-loam soils of southeastern Nigeria. Combined application of poultry manure and urea is needed to increase the productivity of groundnut in the soils of southeastern Nigeria. Increased application of urea may be detrimental to the process of nodulation, which may affect nitrogen economy especially residual nitrogen reserves for subsequent crop. Heavy mineral nitrogen application for groundnut production could be harmful, rather the addition of poultry manure 2 tons ha^{-1} with urea at 1ton ha^{-1} as booster dose should be emphasized in order to not only boost soil fertility and productivity, but also to enhance nodulation processes in groundnut for long lasting fertility.

RECOMMENDATION

Based on the findings of this study, the following recommendations are made

- i. Urea should be integrated with poultry manure for soil fertility improvement in groundnut production.
- ii. Heavy mineral nitrogen fertilization in groundnut production should be discouraged
- iii. Application of 2tonsha^{-1} poultry manure and 1tonha^{-1} urea should be encouraged to enhance soil fertility and nodulation of groundnut.

REFERENCES

- Adeiran, J. A., Akande, M. O., Taswo, L. B., and Sobulo, R. A.(1999). Comparative effectiveness of organic based fertilizer with mineral fertilizers on crop yield. In: Management of the soil resources for sustainable agricultural production in the 21st century, proceedings of the 25th conference of the soil science society of Nigeria (SSSN), Benin City. Pp 91-93.
- Adekiya, A.O. and Agbede, T.M.(2016). Effect of methods and time of poultry manure application on soil and leaf nutrient concentration, growth and fruit yield of tomatoes (*Lycopersiconesculentum*).Journal of Saudi Society of Agricultural Science 9(1):45-52
- Amba, A.A., Agbo, E.B. and Garba, A.(2013), Effect of nitrogen and phosphorus fertilizers on nodulation of some selected legumes in Bauchi, Northern Guinea Savannah of Nigeria, International Journal of Biosciences, 3(10):1-7.
- Anteneh Argaw (2017),Organic and inorganic fertilizer application enhances the effect of *Bradyrhizobium* on nodulation and yield of peanut (*Arachis hypogea* L.) in nutrient depleted and sandy soils of Ethiopia.Intrenational Journal of Recycling Organic Waste in Agriculture, 6(3):219-231.
- Azu, Donatus E.O., Osodeke, V.E. and Nwanja, O.U. (2017).Effect of algae on phosphorus sorption characteristics of an ultisol of Southeastern Nigeria. Direct Research Journal of Agriculture and Food Science, vol 5(2):88-95.

270 Benton, Jones J.(2001). Laboratory Guide for Conducting Soil Test and Plant Analysis.CRC Press Boca Raton
271 Washington D.C.

272 Brady N.C and Weil, R.R. (2008).The Nature and properties of soils (12thedn).Prentice Hall Inc. New Jersey.

273 Bray, R.H and Kurtz L.T. (1945). Determination of total organic and available forms of phosphorus in soils. Soil sci.
274 59:39-45.

275 Brough, S.H. and Azam-Ali, S.N. (1992). The effect of soil moisture on the proximate composition of Bambara
276 (*Vigna subterranean(L)verdc*) J. Sci. Food Agric., 60:197-203.

277 Carter, M. R. and Greforich, E. G. (2008). Soil Smapling and Methods of Analysis 2nd (ed). *Can. Soc. Soil Sci.*
278 1224p.

279 Chriatanson, C.B and Vlek, P.L.G(1991). Alleviating soil fertility constraints to food production in west Africa:
280 Efficiency of N fertilizer applied to food crops. In Mokwunye.U.(Ed), Alleviating soil fertility constraints
281 to increase crop production in west Africa. Kluwer Academic publishers, Dordrecht, Netherlands, pp. 45-
282 59.

283 Dotaniya, M.L., Datta, S.C., Biswas, D.R., Dotaniya, C.K., Meena, B.L., Rajendiran, S., Regar, K.L. and Lata, M.
284 (2016) Use of sugarcane industrial by-products for improving sugarcane productivity and soil health. Int J
285 Recycl Org Waste Agric 5:185–194

286 Duchaufour, P. (1976). AllasEcologiqu dos sold du Monde, Paris, pp. 178

287 Eneje, R. C. and Azu E. O. (2009). Algae compost effect on soil nutrient status and aggregate stability. Journal of
288 Soil Society of Nigeria, 19(2):88-95.

289 Faizah, A.W.(1980). Symbiosis rhizobium leguminensiscompetitivities.Introduction den le sol de sunches de R.
290 Melilotic.These Doctarat.Clande-Benard.University Lyon France.

291 FAO (1976) organic materials as fertilizers, soil bulletins, 27 Rome.

292 Frankow-Lindberg, B.E.andDahlin, A.S. (2013). N₂ fixation, N transfer, and yield in grassland communities
293 including a deep-rooted legume or non-legume species. *Plant and Soil*, 370, 567–581.

294 Gunri, S.K. Nath, R.(2012). Effect of organic manures, biofertilizers and biopesticides on productivity of summer
295 groundnut (*Arachishypogaeae* L.) in red and laterite zone of west Bengal. Indian Council of Agricultural
296 Research.<http://www.icar.org.in/> .

297 Jemo, M.C., Nolte, C., Tchienkoua, M.I and Abaidoo, R.C. (2010).Biological nitrogen fixation potential by
298 soyabeans in two low-p soils of southern Cameroon.Nutr.Cycl.Agrococyst, 88:49-58.

299 Kamprath, E. J. (1967). Soil acidity and response to liming tech: Bull No.4, Int. soil testing series, N. C. State
300 University.

301 Linnemann, A.R.(1988). Cultivation of Bambara groundnut in Northern Nigeria. Trop. Crops, Comm., 15:1-14.

302 Lomoro, P.M.A.(2008). The response of groundnut (*Arachishypogaea*) to phosphorus and organic fertilizer under
303 irrigation. A Thesis submitted to the University of Khartoum in partial fulfilment of the requirement for the
304 Degree of Master of Science(Agric), 74 pages.

305 Mukurumbira, L.M. (1985). Effects of the rate of fertilizer nitrogen and previous grain legume crop on maize yields.
306 Zimbabwe Agric. J., 83:177-180.

307 Naveenkumar, B.T., Malligawad, L.H., Halikatti, S.I., Hiremath, S.M., Srineevasa, M.N. and Bidari, B.I.(2015).
308 Effect of different ratios and levels of nitrogen and phosphorus fertilizers, and top dressing of nitrogen
309 fertilizers on growth and yield of groundnut.Karnatata J. Agric. Sci., 28(1):8-10.

310 Njoku, J.C., Okpara, D.A. and Asiegbu, J.A. (2006). Growth and yield response of sweet potato to inorganic
311 nitrogen and potassium in a tropical ultisol. *Nig. Agric. J.* 32:30-41.

312 Nyamangara, J. And Nyagumbo, I.(2010). Interactive effects of selected nutrient resources and tied-ridging on plan
313 growth performance in a semi-arid smallholder farming environment in central
314 Zimbabwe.Nutr.Cycl.Agroecoyst, 88: 103-109.

315 Odendo, M., and Okwosa E. (2004),. Potential for adoption of legume green manure on smallholder farms in
316 western Kenya, In: Bation A. (Ed) Managing nutrient cycles to sustain fertility in sub-saharan Africa
317 Science Publishers, Nairobi, Kenya, pp. 557-570.

318 Odurukwe, S. O., Aruebunwa, F. A., Iloka, A. W., Udeabor, A. and Ibedu, M. A. (1995).Physical environment of
319 southeast Nigeria. In: Indigenous fallow and multipurpose tree and shrub species in the farming system, southeastern
320 zone of Nigeria. A report of diagnostic survey, NRCRI, UmudikePubls.

321 Ogundare, K., Agele, S. and Aiyelari, P. (2012) Organic amendment of an ultisol: effects on soil properties, growth,
322 and yield of maize in Southern Guinea savanna zone of Nigeria. *J Recycl Org Waste Agric* 1:11

323 Onwuka, M.I., Osodeke, V.E. and Okolo, N.A.(2007). Amelioration of soil acidity using cocoa husk ash for maize
324 production in Umudike area of southeastern Nigeria .*Tropical and Subtropical Agroecosystem* 7:41-45.
325

326 Osodeke, V.E.(2002). Estimation of nitrogen fixing capacity of soybean (*Glycin max*) in soils of the forest zone of
327 southeastern Nigeria. *J. Appl. Chem Res.* Vol 8,pp 28 – 34.

328 Panda, P.K.; Nandi, A., Swain, P.K., Patnaik, S.K. and Patnaik, M. (2012) Soil amendment on growth, nodulation,
329 yield, soil health, and economics of Cowpea.*Int J Veg Sci* 18:284–297

330 Pansu, M and Gautheyrou, J. (2006). Handbook of Soil Analysis: Mineralogical Organic and Inorganic Methods.
331 Springer, p995.

332 Rezig, A.M.R., Elhadi, E.A. and Mubarak, A.R. (2012) Effect of incorporation of some wastes on a wheat–guar
333 rotation system on soil physical and chemical properties. *J Recycl Org Waste Agric* 1:1

334 Shehu, H.E, Kwari, J.D and Sandable, M.K. (2010).Effect of N.P.K by sesame (*Sesamumindicum*). *Int. J. Bool.*
335 12:845-850.

336 Simmone, E. N., Jones, J. B., Mills, A. H., Smittle, A. A. and Hussey, C. G. (1994). Comparison of analytical
337 methods for nitrogen analysis in plant tissues.*Common Soil Science*, 24:1609-1616
338

339 Somasagaran, P., Abaidao, R.C and Kumaga, F. (1992). Host-bradyrhizobium relationship and nitrogen fixation in
340 bambaragroundnut (*Vocandzieasubterranean* L.). *Trop. Agric.*, 67:137-142.

341 Soremi, A.O., Adetunji, M.T., Adeyuyigbe, C.O., Bodunde, J.G. and Azeez, J.O.(2017). Effect of poultry manure on
342 some soil chemical properties and nutrient bioavailability to soybean.*Journal of Agriculture and Ecology*
343 *Research International*, 11(3):1 – 10.

344 Sorensen, J., and A. Sessitisch. (2007). Plant –associated bacteria-Lifestyle and molecular interactions.

Thedchanamoorthy, K. and Gurung, I (1999). Scope of enhancing nodulation of groundnut through soil amendments by crop residue of wide C:N ratio. *AGRIEAST* 1(1):56-58.

Udo, E.J., Ibia, T.O. ogunwale, J.A., Ano, A.O and Esu, I.E (2009). *Manuel of Soil, Plant and Water Analysis*. Sibon Books Ltd, Lagos, Nigeria.

Waldrip, H., He, and Z. and Erich, S. (2011). Effect of poultry manure amendment on phosphorus uptake by ryegrass, soil phosphorus fractions and phosphatase acidity. *Boil. Fertile. Soils*. 47:407-418.

Walley, F.L., Tomm, G.O; Matus, A.; Slinkard, A.E., and Van Kessel, C. (1996). Allocation and cycling of nitrogen in an alfalfa-bromegrass sward. *Agronomy Journal*, 88, 834-843.

UNDER PEER REVIEW

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392