

Inducing Salinity Tolerance of Rosemary (*Rosmarinus officinalis* L.) Plants By Chitosan or Zeolite Application

ABSTRACT

Aims: Rosemary (*Rosmarinus officinalis* L.) is the foremost economically medicinal plants worldwide, its production like other crops affected negatively by environmental stresses. The adverse effects of salinity on crop production are more drastic. Application of chitosan or zeolite might considerably restore the plant productivity under the environmental stress. The present investigation aimed to evaluate the role of chitosan or zeolite on counteracting the deleterious impact of salinity on rosemary growth, oil percentage and its chemical compositions

Study design: A factorial experiment was done in a randomized complete block design system with five replications.

Place and Duration of Study: The pot experiments were done at the Agric. Botany Experimental Farm and Laboratory, Fac. of Agric., Mansoura Univ. , Egypt during 2015/2016 and 2016/2017 seasons.

Methodology: The factorial combinations of two zeolite concentrations (0, 4, 8 g/kg) as a soil additive and two chitosan foliar applications (0, 250 and 500 mg/l) and three salinity levels (0, 50, 100 mM NaCl) were considered.

Results: Shoot and root length, branches number, shoot and root dry weight were declined as a result of salinity. Similarly, the concentrations of photosynthetic pigments, minerals, and essential oil yield were decreased, whereas Na⁺ and Cl⁻ increased in each season.

Application of chitosan or zeolite counteracted the depressing effects of salinity on plant growth, photosynthetic pigments, oil %, and minerals. They declined sodium and chloride concentration in each shoot and root compared to untreated plants.

Conclusion: It could conclude that zeolite at 8 g/kg soil, showed a uniform impact in alleviating of rosemary growth inhibition and its productivity under salinity stress condition

Keywords: [Chitosan, Chlorophyll, Rosemary, Salinity, Zeolite]

1. INTRODUCTION

Medicinal plants are usually used in the manufacture of synthetic drugs based on its chemical structures [1] . Rosemary (*Rosmarinus officinalis* L.), a member of the Lamiaceae family contains flavonoids and essential oils. Rosemary essential oils are known to contain anti-microbial compounds and are therefore used in the manufacture of herbal shampoos to strengthen the hair and in the flavoring and conservation of food product [2, 3].

Salinity has drawn intensive attention worldwide like, Egypt. The deleterious effects of salinity may be attributable to osmotic and ionic effects as well as oxidative stress [4]. The free radicals induced by salinity disrupt normal metabolism through lipids peroxidation, proteins denaturation and nucleic acids [5]. Several authors studied the impact of salinity on numerous plant growth traits and obtained negative responses [6, 7].

Application of soil amendments or foliar application so as to enhance plant productivity is one amongst the foremost important approaches to overcome salinity stress. Because of its low cost and great versatility, zeolite plays a crucial role in agriculture. It can be used to improve the soils, boost the consequences impacts of fertilizers-alike, and as a component of substratum for the development of different crops [8, 9]. Zeolite application results in improving water retention capability [10] and permitting some ions and blocking others [11]. In this concern, zeolite application is that the slow release of sorbed nutrients that stop fast leaching, therefore facilitating an adequate supply of nutrients to the crops [12]. Other studies showed that zeolite served as the stabilization agent in soybean growth, hindrance of salinization [13] and induced water availability in strawberry fields [14]. Recently, Hazrati et al. [10] proved that application of zeolite at 4 and 8 g/kg soil increased leaves number per plant fresh weight of leaves, water use efficiency.

Chitosan (β -1,4-linked glucosamine, CHI) has attracted tremendous consideration as potentially vital biological resources and environmentally friendly with numerous usage in agriculture [15]. CHI has been used in agricultural systems to induce crop productivity [16], to protect plants against oxidative stress [17] and to improve plant growth [18, 19].

Therefore, the main objective of this study was to evaluate the role of zeolite or chitosan on counteracting the deleterious impact of salinity on growth characters and essential oil content as well as some biochemical characteristics of rosemary plant.

2. MATERIAL AND METHODS

Two pot experiments were done at the Agric. Botany Dept. Experimental Farm and Laboratory, Fac. of Agriculture, Mansoura University throughout the 2015/2016 and 2016/2017 years.

2.1 Plant material

Uniform terminal cuttings of *Rosmarinus officinalis* L., 10-12 cm length, were planted within the nursery under shaded conditions for rooting. Once three months, on 15th October in both seasons, the rooted cuttings were separately transplanted in plastic pots 25 cm in inner diameter containing clay loamy soil. The physio-chemical characteristics of the used soil were estimated consistent with Chapman and Pratt [20] and Presented in Table (1).

Table 1. Physiochemical characters of the experimental soil in 2015/2016 and 2016/2017 years.

Soil properties	1 st season	2 nd season	Soil properties	1 st season (meq/L)	2 nd season (meq/L)
Clay %	42.1	41.2	Cations	Calcium	1.2
Silt%	25.6	24.5		Magnesium	1.0
Fine Sand%	24.3	23.4		Sodium	2.3
Coarse Sand%	8.0	7.9		Potassium	0.4
Hygroscopic Water%	5.1	5.2	Anions	Carbonate	----
SSP%	59	60		Bicarbonate	1.3
EC dSm-1	0.53	0.56		Chloride	2.4
pH (1:1.5, soil: water)	7.46	7.44		Sulphate	1.2
					1.1

2.2 Experimental design, treatments and growth conditions

A factorial experiment was done in a randomized complete block design system with five replications. The factorial combinations of two zeolite concentrations (0, 4, 8 g/kg) as a soil additive and two chitosan foliar applications (0, 250 and 500 mg/l) and three salinity levels (0, 50, 100 mM NaCl) were considered.

The pots were held under natural environmental conditions and every pots were irrigated with tap water for 30 days, throughout that, the success of transplants took place, then the rosemary plants were subjected to salinity. Salinization was performed by adding NaCl solution to adjust the salt concentration within the soil at different examined. The water content of the pots was maintained at 75% field capacity until the end of the experiment. Zeolite concentration was mixed with the surface soil whereas; chitosan was sprayed thrice until dropping after covering the soil surface. Initial chitosan treatments occurred one week after salt treatments with 21-day intervals.

Plants were fertilized with 2.5 g ammonium sulfate (20.6% N), 3.59 g calcium super-phosphate (15.5% P₂O₅) and 1.25 g potassium sulfate (45% K₂O) per pot after 3 weeks from transplanting. All plants received traditional agricultural practices, consistent with the recommendation of ARC, Egypt, whenever they were required.

2.3 Sampling and Measurements

After 21 days from the last chitosan foliar spray (1st April), Rosemary plants were harvested and following parameters were measured.

2.3.1 Growth parameter

Plant height, number of branches per plant and both fresh and dry weights of shoots and roots were recorded.

2.3.2 Biochemical attributes

Photosynthetic pigments were extracted from the 1st mature leaf and determined spectrophotometrically as delineate by Lichtenthaler and Wellburn (1985).

Ground dried materials for the shoot and root systems were used for mineral element determination.

The dried materials were wet digested with aHClO₃/H₂SO₄ mixture (1:1 v: v) as reported by

Chapman and Pratt [20]. Total nitrogen was determined colorimetrically by Nessler's methods [21]; potassium by flame photometrically [22], and phosphorous according to the method of Cooper [23]. Moreover, sodium and chloride were extracted from dried plant materials consistent with Chaudhary et al. [24]. Sodium within the aqueous extracts was measured with a flame photometer, whereas, chloride was determined by titration with 0.001 N AgNO₃ using potassium chloride as an indicator.

2.3.3 Essential oil percentage and yield

The essential oil within the air-dried herb of each treatment was extracts determined by the water-distillation technique consistent with Egyptian Pharmacopoeia [25]. Essential oil yield (ml/plant) was estimated in proportion to the herb air-dry weight.

2.4 Statistical analysis

Data were analyzed by analysis of variance (ANOVA) technique by computer software MSTATC and significant treatment means were compared using least significance difference (LSD) test at 0.05 probability level according to Gomez and Gomez [26].

3. RESULTS AND DISCUSSION

3.1 Growth Parameters and essential oil yield

Tables (2) shows that plant growth represented as shoot and root length, and shoot and root dry weights of Rosemary were significantly declined due to salinity, and therefore the greatest reduction was assessed under high salinity. Essential oil % in rosemary shoot were enhanced by increasing salinity levels, this increase was accompanied by decreasing the essential oil yield due to decreasing shoot mass.

All plant growth and essential oil yield were significantly enhanced by the application of either zeolite or chitosan. The highest values were obtained by soil additive of 8 g/kg soil zeolite within each season comparing to control treatment

As regards to the interactions, the data given within the same table disclosed that using chitosan or zeolite under low salinity level significantly enhanced all growth parameters and essential oil yield. The highest values were obtained by adding 8 g/zeolite/kg soil. On the opposite, application of either zeolite or chitosan under severe salt stress counteracted the injuries of salinity on growth parameter that enhanced growth parameter as compared with untreated plants under high salinity levels.

Table 2. Effect of Salinity, either zeolite or chitosan , as well as their interactions on growth characters and essential oil yield of Rosemary plants

Treatments	Shoot Length (cm)		Root Length (cm)		Shoot dry weight (g)		Root dry weight (g)		Essential oil %		Essential oil yield (g/plant)	
	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd
	season	season	season	season	season	season	season	season	season	season	season	season
Salinity												
C	41.33	34.26	16.46	21.86	17.66	7.21	5.59	1.41	0.583	0.578	10.435	4.176
MS	39.40	31.60	15.80	19.73	15.29	9.01	5.69	1.39	0.669	0.661	10.708	6.386
SS	35.26	26.06	12.80	15.06	10.45	5.16	3.98	0.87	0.748	0.736	8.095	3.919
LSD 5%	2.71	1.80	1.53	2.11	2.27	0.57	0.56	0.13	0.01	0.01	1.54	0.39
Antitranspirants												
W	33.66	26.22	13.00	16.11	10.84	4.967	3.60	0.89	0.512	0.517	5.281	2.419
Ze1	41.55	33.88	15.77	20.55	16.81	7.62	5.52	1.33	0.713	0.706	11.758	5.316
Ze2	41.11	36.00	17.66	22.44	18.54	10.41	6.59	1.78	0.813	0.783	15.022	8.323
Chi1	39.33	29.66	14.55	17.55	13.65	6.63	5.07	1.11	0.676	0.676	9.090	4.454
Chi2	37.66	27.44	14.11	17.77	12.49	6.02	4.65	1.00	0.620	0.611	7.579	3.623
LSD 5%	3.506	2.336	1.977	2.726	2.931	0.741	0.726	0.176	0.015	0.022	1.997	0.504
Interaction												
CW	38.33	30.00	16.00	21.00	15.48	7.05	4.82	1.28	0.417	0.420	6.450	2.952
CZe1	43.33	36.66	16.33	22.33	20.00	7.29	5.74	1.42	0.650	0.637	12.900	4.640
CZe2	43.33	38.66	17.33	23.66	19.23	7.38	6.62	1.61	0.683	0.663	13.140	4.894
CChi1	41.66	35.33	16.33	21.00	16.96	7.28	5.587	1.40	0.607	0.617	10.273	4.495

CChi2	40.00	30.66	16.33	21.33	16.67	7.07	5.17	1.35	0.560	0.553	9.321	3.899
MSW	31.33	25.00	12.66	14.33	9.98	5.09	3.84	0.87	0.513	0.510	5.107	2.600
MSZe1	43.33	38.66	17.66	23.66	18.61	9.56	6.34	1.63	0.683	0.677	12.748	6.482
MSZe2	43.33	39.66	20.66	26.00	21.88	17.00	7.52	2.51	0.873	0.840	19.107	14.263
MSChi1	39.66	27.33	14.00	17.00	13.39	6.78	5.58	1.03	0.660	0.667	8.813	4.526
MSChi2	39.33	27.33	14.00	17.66	12.57	6.62	5.53	0.94	0.617	0.613	7.765	4.057
SSW	31.33	23.66	10.33	13.00	7.06	2.75	2.51	0.53	0.607	0.620	4.287	1.705
SSZe1	38.00	26.33	13.33	15.66	11.81	6.00	4.48	0.95	0.807	0.803	9.537	4.826
SSZe2	36.66	29.66	15.00	17.66	14.51	6.857	5.62	1.24	0.883	0.847	12.820	5.812
SSChi1	36.66	26.33	13.33	14.66	10.62	5.84	4.04	0.91	0.760	0.743	8.183	4.341
SSChi2	33.66	24.33	12.00	14.33	8.23	4.37	3.25	0.71	0.683	0.667	5.650	2.911
LSD5%	ns	4.046	ns	ns	ns	1.284	ns	0.305	0.026	0.299	ns	0.873

C, control; MS, Mild Salinity; SS, Severe salinity; W, water; Ze1, Zeolite 4g/Kg; Ze2, Zeolite 8g/Kg; Chi1, Chitosan 250 mg/l; Chi2, Chitosan 500 mg/l

The responses of the Rosemary plant to a high level of salinity were reflected by decreases in shoot and root length, as well as shoot and root dry weight. The repressing effects of salinity on these parameters add more support to the previous finding [6, 7, 27, 28]. The injurious impacts of salinity on plant growth are due to the inhibition of photosynthesis, the induction of growth inhibitor, and reduction of leaf area [29], leaf protein [27], and reduced ability to provide and utilize assimilates/photosynthates [29]. Alternatively, Romero and Maranon [30] added that the specific ion effects of salinity, caused toxicity and nutritional imbalance, leading to decline in potassium, calcium and phosphorous absorption. The deleterious impact of severe salinity concentration on growth may be ascribed to that salinity has been shown to reduce the synthesis of nucleic acid and protein in several plants that could result in disturbance in metabolic activities, cell division and elongation [31].

The positive impacts of chitosan and/or zeolite on plant growth noticed in the present investigation may be attributed to its effect on inducing antioxidant system [27, 32]. Also, chitosan or zeolite used enhanced potassium percentage (Table, 4), that may increase the chloroplast number per cell, the cells number per leaf and leaf area [31], maintaining photosynthetic activity and metabolic transport [18, 33]. Additionally, chitosan increased nitrate reductase, glutamine synthetase, and protease in the functional leaves that improved plant growth and development [34]. Khan et al. [35] found that foliar application of CHI enhanced net photosynthetic rate (PN), this increase was related with a rise in stomatal conductance (gs) and transpiration rate (E), without any effects on intercellular CO₂ concentration (Ci) that induced photoassimilate production.

It has been reported that increasing essential oil percentage within the plant may be a mechanism to adapt the plant to stress conditions [19, 36]. In rosemary, salt stress stimulated essential oil accumulation owing to the higher density of oil glands. By exerting stress, the essential oil percentage is increased, but the shoot dry weight decreased resulted in reducing essential oil [19, 37]. These results are in agreement with those reported earlier on other aromatic plant species such as *Pimpinella anisum* L. [38], *Satureja hortensis* L. [39]; *Mentha spicata* L. and *Rosmarinus officinalis* L. [40].

Chemical compounds (secondary metabolites) in essential oils of plants are directly affected by environmental factors, especially drought stress [41]. Drought stress reduces the absorption and transfer of nutrients in plants, resulting in consequences including quantitative and qualitative changes in secondary metabolites like essential oils [42]. Various studies have shown that CO₂ and

glucose are known as proper precursors for the synthesis of plant essential oils and which there's a direct correlation between photosynthetic products and essential oil production [43]. Consequently, physiological drought-induced damages like oxidative stress (ROS increase), photosynthetic pigment decomposition, alteration of membrane structure and proteins, deactivation of enzymes, stomatal closure, and dropped CO₂ under the stomata result in decreased photosynthesis (reduction of glucose as essential oil precursor) and reduced cellular growth and development, ultimately resulting in qualitative and quantitative falls within the essential oil content of plants [42]. Among the chemical compounds of thyme essential oil, only three compounds (thymol, carvacrol, and trans-caryophyllene) rose significantly with increasing drought stress levels, that is keeping with alternative studies [37], probably indicating a type of plant adaptation to stress conditions. Zaghloul et al. [9] indicate that zeolite application enhanced essential oil percentage and yield.

3.2 Photosynthetic pigments

Table (3) show that chlorophyll a, chlorophyll b and total chlorophylls markedly reduced with increasing salinity levels in both seasons. The greatest decline was obtained under high salinity. Carotenoid concentration was parallel to chlorophylls except in the first season that low salinity nonsignificantly enhanced. The ratio between chlorophyll a and chlorophyll b was reduced in the first season and enhanced in the second season as compared with control plants.

Generally, under control, application of either chitosan or zeolite in most cases significantly increased photosynthetic pigments concentration as compared with untreated control plants. The highest values of chl a, b, total chlorophylls, and carotenoids (0.919, 0.494 & 0.513, 0.274 & 1.433, 0.769 & 0.164, 0.061 mg/g FW) were obtained due to application of zeolite at 4 g/kg soil. However, application of chitosan or zeolite reduced chlorophyll a:b ratio in the 2nd season whereas increased it in the 1st season.

The interaction treatments indicated that application of either chitosan or zeolite mitigated the injuries of salinity on photosynthetic pigment concentration. The highest values of chlorophyll a in the 1st season, total chlorophyll in both seasons were obtained by application of zeolites under moderate salinity. The highest chlorophyll a and carotenoids in the 2nd season were obtained by zeolite under normal condition (Table, 3). The lowest chlorophyll a to chlorophyll b ratio was observed under the treatment of water under high salinity in the 1st season and low salinity level in the 2nd season.

Table 3. Effect of Salinity, either zeolite or chitosan, as well as their interactions on photosynthetic pigments of Rosemary plants

Treatments	Chlorophyll a		Chlorophyll b		Total chlorophyll		Total carotenoids		Chlorophyll a:b ratio	
	1 st season	2 nd season	1 st season	2 nd season	1 st season	2 nd season	1 st season	2 nd season	1 st season	2 nd season
Salinity										
C	0.972	0.428	0.508	0.250	1.481	0.679	0.162	0.058	1.913	1.700
MS	0.902	0.360	0.467	0.212	1.370	0.573	0.167	0.045	1.910	1.723
SS	0.603	0.284	0.371	0.168	0.974	0.452	0.111	0.053	1.632	1.943
LSD 5%	0.083	0.059	0.040	0.032	0.108	0.085	0.044	ns	0.173	ns

Antitranspirants										
W	0.797	0.294	0.449	0.175	1.247	0.470	0.158	0.043	1.761	2.063
Ze1	0.919	0.494	0.513	0.274	1.433	0.769	0.144	0.061	1.762	1.817
Ze2	0.812	0.315	0.426	0.197	1.238	0.513	0.154	0.043	1.894	1.617
Chi1	0.795	0.347	0.425	0.210	1.221	0.558	0.141	0.057	1.803	1.697
Chi2	0.806	0.335	0.429	0.195	1.236	0.531	0.136	0.056	1.873	1.749
LSD 5%	ns	0.077	0.051	0.042	0.139	0.110	ns	ns	ns	ns
Interaction										
CW	0.806	0.412	0.431	0.260	1.238	0.673	0.175	0.046	1.868	1.584
CZe1	0.816	0.636	0.513	0.292	1.329	0.929	0.113	0.124	1.644	2.235
CZe2	0.927	0.308	0.442	0.193	1.370	0.501	0.155	0.027	2.041	1.607
CChi1	1.235	0.416	0.614	0.259	1.849	0.676	0.159	0.049	2.013	1.585
CChi2	1.077	0.369	0.539	0.246	1.617	0.616	0.210	0.044	2.000	1.488
MSW	0.898	0.238	0.508	0.166	1.499	0.405	0.228	0.028	1.949	1.429
MSZe1	1.440	0.611	0.684	0.386	2.124	0.997	0.234	0.023	2.102	1.578
MSZe2	0.616	0.313	0.328	0.177	0.943	0.490	0.162	0.046	1.882	1.768
MSChi1	0.810	0.245	0.447	0.124	1.258	0.370	0.105	0.055	1.821	1.971
MSChi2	0.656	0.392	0.366	0.210	1.025	0.602	0.107	0.073	1.796	1.868
SSW	0.595	0.232	0.408	0.099	1.004	0.331	0.073	0.055	1.465	3.175
SSZe1	0.502	0.236	0.342	0.144	0.844	0.380	0.084	0.037	1.539	1.638
SSZe2	0.893	0.326	0.507	0.222	1.400	0.548	0.145	0.055	1.758	1.477
SSChi1	0.340	0.381	0.214	0.247	0.555	0.628	0.160	0.066	1.574	1.536
SSChi2	0.684	0.245	0.381	0.129	1.066	0.375	0.092	0.052	1.824	1.890
LSD5%	0.018	0.077	0.09	0.072	0.241	0.191	0.098	0.042	ns	ns

C, control; MS, Mild Salinity; SS, Severe salinity; W, water; Ze1, Zeolite 4g/Kg; Ze2, Zeolite 8g/Kg; Chi1, Chitosan 250 mg/l; Chi2, Chitosan 500 mg/l

The reduction in photosynthetic pigment concentration by salinity was confirmed by earlier reports [6, 7]. Salinity stress enhances the activity of chlorophyll degrading enzyme chlorophyllase and interferes with the de-novo synthesis of chlorophyll binding proteins [44]. Moreover, salinity can lead to the destruction of the fine structure of chloroplasts, instability of pigment-protein complex [45], which leads to oxidation of chlorophyll and decreased its concentration [46] and alteration in the content and composition of carotenoids. Accumulation of sodium and chloride (Table 5), could affect chlorophyll biosynthesis due to its effect on the activity of Fe-containing enzymes, cytochrome oxidase. Yeo and Flowers [47] showed that there's a negative relationship between total chlorophyll and sodium content where sodium and chloride ions penetrate the chloroplasts [48]. The hormonal imbalance under salt stress could have an effect negatively through decreasing the biosynthesis of cytokinin in salinized plant root and translocated to the shoot [45] and induced ABA accumulation leading to inducing chlorophyll breakdown [49] or inhibiting chlorophyll synthesis [50]. In addition, salt stress accelerates leaf senescence through inhibitory chlorophyll synthesis [51].

The superiority of either chitosan or zeolite in photosynthetic pigments was in harmony with [19, 52] for chitosan, and [53] for zeolite; that indicated application of chitosan considerable enhanced photosynthetic pigment concentration and accelerate carotenoids biosynthesis that protect chlorophyll from oxidation. The influence of CHI on alleviating the water stress impact on the photosynthetic pigments might be due to the fact that CHI increased the endogenous level of cytokinins that

stimulates chlorophyll biosynthesis. Chibu and Shiayama [54] proved these stimulative effects to more availability of amino compounds released from CHI. Data in the present investigation indicate that foliar application of CHI enhanced considerably either nitrogen and potassium content in the plant shoot (Table, 4) that can be taking part in a very important role in increasing the number of chloroplast per cell, the cell size and number per unit area as well as enhanced the synthesis of chlorophyll [55]. Moreover, Si form application could increase the chlorophyll concentration and chloroplast ultrastructure under stress conditions [56]. These results are in accordance with El-Saedy et al. [57] who show that application of Si enhanced total chlorophyll contents in grapevine leaves.

Data in the same table show that Chla:b ratio increased due to salinity, proved that chl b was degraded at a higher rate than chl a (Table, 3). This could be explained by the fact that the first step in the chl b degradation involves its conversion to chl a [58]. The increase in the ratio of Chla:b has been connected with lower levels of light-harvesting chlorophyll proteins; LHCPs [59]. The decline in LHCPs content is an adaptive defense mechanism of plant organs that permits them to endure the environmental stresses [60].

Photosynthesis is limited under salinity that could result in photoinhibitory injury. Photoinhibitory injury could be avoided by photorespiration, through scavenging systems that remove ROS with carotenoids [61]. Carotenoids are synthesized and accumulate in chloroplast wherever they play a crucial role in the light harvesting complex assembly and performance. Salinity induced a specific change in the level of carotenoids that reduced carotenoid concentration (Table, 3) due to degradation of β -carotene and formation of zeaxanthin, that defend the plant against photoinhibition [62], and no longer available for protection against additional damage, resulting in long-term chlorosis [63]. Thus, it looks that the decrease in carotenoids levels under saline stress could result in chlorophyll degradation [64].

3.3 Minerals percentage

Data in Tables (4 and 5) indicate that salinity stress caused disturbing in the electrolyte balance, leading to the deficiency of some nutrients and the excess of certain unwanted ions. Data presented in Table (5) prove that sodium percentage and chloride concentration were increased with salinity. This increase was accompanied by a corresponding decrease in nitrogen, phosphorous, potassium and K/Na ratio in both shoot and root system. The great reduction occurred under high salinity levels compared with control plants.

Application of either zeolite or chitosan, in special, 8 g/kg zeolite, significantly increased in most cases the K^+/Na^+ ratio, nitrogen, phosphorous and potassium percentage, but declined the percentage of sodium and chloride concentration in both shoot and root systems during the two experimental seasons. Additionally, they counteracted the injury's effects of salinity on minerals either in the shoot and root systems. The highest values of macronutrients in most cases obtained by application of 8g/kg soil zeolite under low salinity as compared with untreated control plants. Alternatively, the lowest sodium and chloride obtained because of the applying of zeolite at 8g/kg soil under normal condition.

Table 4. Effect of Salinity, either zeolite or chitosan , as well as their interactions on nitrogen, phosphorous and potassium percentage in both shoot and root of Rosemary plants

Treatments	Nitrogen in shoot		Nitrogen in root		Phosphorous in shoot		Phosphorous in root		Potassium in shoot		Potassium in Root	
	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd
	season	season	season	season	season	season	season	season	season	season	season	season
Salinity												
C	2.340	2.656	2.852	2.265	0.697	1.000	0.516	0.663	2.587	2.515	2.322	2.878
MS	2.180	2.082	2.435	2.690	0.708	0.949	0.487	0.504	2.022	2.448	1.847	2.502
SS	1.637	1.160	1.674	2.353	0.457	0.526	0.292	0.285	1.233	1.542	0.955	1.663
LSD 5%	0.161	0.056	0.206	0.119	0.063	0.077	0.053	0.054	0.333	0.240	0.177	0.173
Antitranspirants												
W	1.259	1.284	1.757	1.547	0.431	0.544	0.286	0.322	1.178	1.469	1.043	1.552
Ze1	2.250	2.335	2.543	2.572	0.736	0.962	0.514	0.569	2.627	2.478	2.134	2.702
Ze2	2.608	2.739	2.979	3.042	0.856	1.156	0.573	0.604	2.231	2.888	2.351	3.113
Chi1	2.094	1.846	2.286	2.680	0.605	0.813	0.417	0.510	1.985	2.082	1.574	2.366
Chi2	2.051	1.627	2.039	2.339	0.477	0.649	0.369	0.415	1.716	1.925	1.439	2.037
LSD 5%	0.208	0.072	0.266	0.154	0.081	0.100	0.069	0.069	0.430	0.310	0.229	0.224
Interaction												
CW	1.971	2.536	2.767	2.018	0.506	1.937	0.454	0.534	2.201	2.246	2.044	2.268
CZe1	2.274	2.794	2.895	2.370	0.738	1.054	0.533	0.669	3.232	2.717	2.762	2.851
CZe2	2.559	2.644	3.209	2.763	0.976	1.130	0.585	0.671	2.358	2.739	2.403	3.546
CChi1	2.417	2.648	2.933	2.109	0.778	1.032	0.521	0.711	2.537	2.470	2.224	2.986
CChi2	2.479	2.659	2.458	2.065	0.485	0.845	0.488	0.732	2.604	2.403	2.179	2.739
MSW	1.218	0.888	1.381	1.826	0.491	0.499	0.266	0.252	0.722	1.439	0.677	1.529
MSZe1	2.661	2.806	2.827	2.447	0.927	1.181	0.672	0.692	2.806	3.143	2.605	3.456
MSZe2	2.715	3.661	3.432	2.840	1.020	1.579	0.729	0.724	2.605	3.658	2.873	3.591
MSChi1	2.161	1.592	2.181	3.181	0.548	0.798	0.402	0.488	2.134	1.798	1.551	2.201
MSChi2	2.145	1.464	2.357	3.157	0.556	0.687	0.368	0.363	1.843	2.201	1.529	1.730
SSW	0.588	0.428	1.122	0.798	0.296	0.195	0.138	0.180	0.610	0.722	0.409	0.767
SSZe1	1.814	1.404	1.906	2.901	0.543	0.651	0.339	0.347	1.843	1.574	1.036	1.798
SSZe2	2.551	1.912	2.296	3.521	0.571	0.760	0.405	0.418	1.731	2.268	1.776	2.201
SSChi1	1.704	1.299	1.745	2.749	0.489	0.609	0.329	0.331	1.282	1.977	0.946	1.910
SSChi2	1.530	0.758	1.302	1.796	0.388	0.416	0.250	0.148	0.700	1.170	0.610	1.641
LSD5%	0.360	0.125	0.462	0.267	0.141	0.173	0.119	0.121	ns	0.538	0.397	0.387

C, control; MS, Mild Salinity; SS, Severe salinity; W, water; Ze1, Zeolite 4g/Kg; Ze2, Zeolite 8g/Kg; Chi1, Chitosan 250 mg/l; Chi2, Chitosan 500 mg/l

Table 5. Effect of Salinity, either zeolite or chitosan , as well as their interactions on sodium, chloride percentage and potassium sodium ratio in both shoot and root of Rosemary plants

Treatments	Sodium in shoot		Sodium in root		Potassium: Sodium in shoot		Potassium: Sodium in root		Chloride in Shoot		Chloride in Root	
	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd	1 st	2 nd
	season	season	season	season	season	season	season	season	season	season	season	season
	Salinity											
C	2.025	1.994	1.842	1.865	1.358	1.430	1.318	1.584	73.55	77.62	36.92	73.55
MS	3.648	2.997	2.667	3.042	0.661	0.869	0.780	0.951	136.57	139.91	90.78	136.60
SS	4.352	3.523	3.198	3.590	0.326	0.491	0.343	0.530	181.62	176.17	99.87	181.66
LSD 5%	0.293	0.178	0.217	0.155	0.150	0.169	0.124	0.106	8.003	5.242	4.942	8.002
	Antitranspirants											

W	4.502	3.841	3.403	3.886	0.380	0.449	0.411	0.512	195.30	198.16	95.131	195.32
Ze1	2.704	2.325	2.266	2.355	1.161	1.205	1.044	1.227	96.39	99.04	68.47	98.76
Ze2	2.259	2.169	1.932	2.088	1.089	1.504	1.310	1.590	90.85	88.67	63.26	88.51
Chi1	3.262	2.667	2.533	2.749	0.723	0.846	0.681	0.991	126.34	123.38	71.00	126.38
Chi2	3.982	3.187	2.712	3.083	0.555	0.647	0.623	0.788	144.02	146.89	81.41	144.05
LSD 5%	0.379	0.230	0.280	0.200	0.194	0.218	0.161	0.137	10.332	6.767	6.381	10.330
Interaction												
CW	2.466	2.712	2.132	2.177	0.911	0.832	0.958	1.042	99.40	98.54	51.59	99.40
CZe1	1.597	1.441	1.664	1.842	1.997	1.897	1.669	1.548	68.63	71.94	30.29	68.63
CZe2	1.486	1.218	1.374	1.575	1.587	2.275	1.774	2.248	51.59	61.06	19.40	51.59
CChi1	2.244	2.043	2.021	1.820	1.160	1.206	1.110	1.646	73.84	77.62	33.13	73.84
CChi2	2.333	2.556	2.021	1.909	1.137	0.940	1.082	1.437	74.31	79.04	50.17	74.31
MSW	5.476	4.094	3.849	4.562	0.127	0.362	0.179	0.338	223.36	238.08	113.1	223.41
MSZe1	2.890	2.712	2.444	2.310	0.967	1.159	1.069	1.499	88.50	91.82	82.83	95.61
MSZe2	2.355	2.712	2.065	2.311	1.093	1.348	1.404	1.581	95.61	88.04	82.83	88.51
MSChi1	2.912	2.310	2.467	2.712	0.730	0.784	0.634	0.814	130.60	130.16	82.83	130.64
MSChi2	4.606	3.158	2.511	3.314	0.390	0.694	0.615	0.520	144.80	151.46	92.30	144.84
SSW	5.565	4.718	4.228	4.918	0.103	0.153	0.096	0.156	1263.13	257.96	120.70	263.17
SSZe1	3.626	2.823	2.689	2.912	0.518	0.559	0.394	0.635	132.03	133.48	92.30	132.06
SSZe2	2.935	2.578	2.355	2.377	0.588	0.890	0.753	0.939	125.36	116.91	87.56	125.43
SSChi1	4.935	3.648	3.113	3.715	0.279	0.548	0.300	0.513	174.6	162.35	97.03	174.66
SSChi2	4.629	3.848	3.603	4.083	0.140	0.307	0.174	0.408	212.96	210.16	101.76	213.00
LSD5%	0.657	0.399	0.485	0.347	ns	0.378	ns	0.238	17.897	11.721	ns	17.895

C, control; MS, Mild Salinity; SS, Severe salinity; W, water; Ze1, Zeolite 4g/Kg; Ze2, Zeolite 8g/Kg; Chi1, Chitosan 250 mg/l; Chi2, Chitosan 500 mg/l

On the other hand, an increase in chloride accumulation accompanied by a decline in shoot nitrate concentrations of plants was observed due to the competition between chloride and nitrate that decreases the nitrate [65]. Silberbush and Ben-Asher [66] found that despite drastic reductions in leaf nitrate concentrations under salinity. These conclusions are supported [67]. Saline conditions will influence the various steps of nitrogen metabolism, reduction and protein synthesis [68]; ammonium uptake [69]. Frota and Tucker [70] expressed that the decline in nitrogen under salinity could also be due to a reduction in water absorbed and also the decrease in root permeability. Accordingly, Papadopoulos and Rendig [71], proved that the impact of salinity on leaf nitrogen and total nitrogen uptake was chiefly through the suppressing impact of salinity on root growth and water uptake.

The influence of salinity on phosphorous content in crop plants is variable and depends upon the cultivars. In our work, the results in Tables (4 and 5) indicate that increasing the salinity level promoted a reduction of phosphorous concentration in plant tissue. This negative relationship between phosphorous and salinity level could also be due to the postulation of Greenway et al. [72] who attributed the reduction in phosphorous and its uptake by tomato plants under saline conditions to a decline in the root absorption potential and to a decrease in the translocation of phosphorous upward through the root as a result of the increase in the osmotic pressure of the root medium. Champagnol [73] found that unlikely chloride and phosphate ions are competitive in terms of plant uptake. However, Papadopoulos and Rendig [71] concluded that chloride might have suppressed phosphorous accumulation in tomato. In other cases, reduction in plant phosphorous content might

result from reduced activity of phosphorous in the soil solution due to the high ionic strength of the media [74]. The decline in P concentration with increased salinity levels might have occurred attributable to decreased P transport under high salt concentrations [75]. This reduction could be due to precipitate phosphorous ions with calcium in salt-stressed soil and become unavailable to plants. Furthermore, the reduction in P accumulation under salinity may will be explained by the fact that Na ion increased the soil pH that successively declined the availability of P to the plants.

The most abundant inorganic cation in vacuoles is potassium that plays a large part in maintaining cell turgor pressure and potassium-homeostasis between cytoplasm and the vacuole [76]. Excess of sodium and chloride, creates a high ionic balance which will improve the selectivity of root membrane [77]. Sodium is often accumulated within the vacuoles wherever it will replace potassium both quantitatively and qualitatively [78]. Many glycophyte species can partially substitute sodium for potassium [79]. Several mechanisms could also be accountable for a reduction in potassium with increasing salinity, as well as the antagonism of sodium and potassium at the uptake site in roots [80]. In the present study, salinity considerably increased sodium concentration in shoot and root. This increase was accompanied by a reduction in potassium content in both shoot and root (Table, 4), indicating a visible antagonism between potassium and sodium [81]. This antagonism may be due to the direct competition between potassium and sodium at a site of ion uptake within the plasmalemma [82]. Sodium enhanced the efflux of potassium into the growth medium, attributable to disturbed membrane permeability [68], possibly due to membrane integrity [51]. Moreover, sodium could also be interference with potassium uptake or transport [83] and inhibit transport of these ions into the roots [84]. Additionally, an excess of sodium content in root media leads to a passive accumulation of this ion as shown in Table (5) results in high sodium/potassium ratio [85].

The increase in sodium content chiefly within the vacuole provides an osmotic adjustment of salt-affected plants [86]. This accumulation may be due to the vital role of sodium in reducing the osmotic pressure that facilitates absorption of water required for plants to tolerate the harmful impact on growth caused by salinity [87]. Many of the deleterious effects of sodium, however, appear to be associated with the structure change including the lose its selective permeability [88].

The role of chitosan or zeolite on inducing ion accumulation isn't totally understood, and there are few studies in this concern [9, 52, 89] under normal or stressed conditions. This increase may be due to enhanced nutrient uptake by improving membrane permeability (Table, 3) and/or giving better-developed root system due to enhanced microbial activity in addition as increased root growth which might have facilitated more efficient nutrient absorption. In this concern, Mali and Aery [90] show that application of Si form plants maintained plasma membrane fluidity, decreased the ratio of phospholipids to protein and enhanced plasma membrane H⁺-ATPase activity, thereby maintaining membrane selectivity to ion influx and enhance potassium uptake. The role of CHI on increasing ion contents may be due to its effects on stabilizing cellular membrane through increasing the antioxidant substances and saving cell membrane from oxidative stress therefore improved plant cell permeability resulting in increasing ion content. This observation was supported by the results of [17, 27, 91] who indicated that application of CHI significantly declined lipid peroxidation, due to stimulation of some antioxidants enzymes leading to decreasing membrane permeability and improved its functions. This

increase could also be caused by the amino components in chitosan and or the higher ability of the plant to absorb nitrogen from the soil when chitosan was degraded. Additionally, application of zeolite increased nitrogen percentage, due to the better use efficiency of applied nitrogen fertilizers coupled with retarded nitrification process, enabling the slow availability of applied nitrogen, leads to reduced loss of nitrogen by volatilization. Zaghloul et al. [9] and Latifah et al. [89] indicate that application of zeolite increased NPK% in thyme plants.

The increase in P accumulation as results of zeolite and CHI application could also be resulted from the prevention of P fixation within the soil and the formation of humophospho complexes, which are easily assailable by the plants [92] and this explains the more of P percentage by rosemary plant in the present study.

4. CONCLUSION

It could be concluded that application of zeolite at 8g/kg soil as a soil additive to Rosemary plants under salinity stress stimulated its growth, essential oil percentage and yield increased photosynthetic pigments and ion percentage in the herb.

REFERENCES

- Abdi GH, Khui MK, Eshghi S. Effects on natural zeolite on growth and flowering on strawberry. International Journal of Agricultural Research 2006; 1:384–389.
- Ahmad P, Mohammad AA, Alyemenia MN, Wijayaa L, Alamd P, Ashrafe M. Mitigation of sodium chloride toxicity in *Solanum lycopersicum* L. by supplementation of jasmonic acid and nitric oxide. J Plant Inter.,2018; 13 (1): 64–72. <https://doi.org/10.1080/17429145.2017.1420830>
- Alam SM. Nutrient uptake by plants under stress conditions. In: Passarakli M, editor. Handbook of plant and crop stress, Marcel Dekker, New York; 1994
- Almela L, Sachez B, Fernandez JA, Roca M, Rabe V. Liquid chromatographic mass spectrometric analysis of phenolics and radical scavenging activity of rosemary extract from different raw material. J.Chromatogr., 2006; A 1120: 221.
- Asada K, Endo T, Mano J, Miyake C. Molecular mechanism for relaxation of and protection from light stress. In: Saton K, Murata N, editors. Stress Responses of Photosynthetic Organisms. Elsevier, Amsterdam; 1998
- Awad AS, Edwards DG, Campbell LC. Phosphorous enhancement of salt tolerance of tomato. Crop Sci., 1990; 30:123-128.
- Baher ZF, Mirza M, Ghorbanil M, Rezaii MZ. The influence of water stress on plant height, herbal and essential oil yield and composition in *Satureja hortensis* L. Flavour Fragr. J. 2002; 17:275-277.
- Bahreinejad B, Razmjoo J, Mirza M. Influence of water stress on morpho-physiological and phytochemical traits in *Thymus doenensis*. Int. J. Plant Prod. 2013; 7:155-166.
- Bengtson C, Klockare B, Larssa S, Sundquist O. The effect of phytohormones on chlorophyllide, protochlorophyllide and carotenoid formation in greening dark grown wheat leaves. Phy. Plant., 1977; 40:198-204.
- Binzel ML, Hasegawa PM, Handa AK, Bressan RA. Adaptation of tobacco cells to NaCl. Plant physiology, 1985; 79:118-125.
- Bistgani ZE, Siadat SA, Bakhshandeh A, Ghasemi Pirbalouti A, Hashemi M. Interactive effects of drought stress and chitosan application on physiological characteristics and essential oil yield of *Thymus daenensis* Celak. The Crop J. 2017; 4(5): 407-415. <http://dx.doi.org/10.1016/J.cj.2017.04.003>
- Bohra JS, Doerffling K. Potassium nutrition of rice (*Oryza sativa* L) varieties under NaCl salinity. Plant and Soil 1993; 152:299-303.
- Bor M, Ozdemir F, Turkan I. The effect of salt stress on lipid peroxidation and antioxidants in leaves of sugar beet *Beta vulgaris* L. and wild *Beta maritime* L. Plant Sci.,2003; 164:77-84.

- 371 Celikta OY, Kocabas EE. Antimicrobial Activities of methanol extracts and essential oils of
372 *Rosmarinus officinalis*, depending on location and seasonal variations. Food Chem.,2007; 100: 553-
373 559.
- 374 Champagnol F. Relationships between phosphate nutrition of plants and salt toxicity. Phosphorous
375 Agric. 1979; 76:35-43.
- 376 Chapman HD, Pratt PE.. Methods of analysis of soil, plants and water. Division of Agricultural
377 Sciences, University of California, Berkeley 1978.
- 378 Charles O, Joly R, Simon JE. Effect of osmotic stress on the essential oil content and composition of
379 peppermint. Phytochemistry, 1994; 29: 2837-2840.
- 380 Chaudhary MT, Wainwright SJ, Merrett MJ. Comparative NaCl tolerance of Lucerne plants
381 regenerated from salt-selected suspension cultures. Plant Sci. 1996; 114:221-232.
- 382 Chibu H, Shibayama H. Effects of chitosan applications on the growth of several crops. In: Uragami T,
383 Kurita K, Fukamizo T, editors. Chitin and Chitosan in Life Science, Yamaguchi; 2001
- 384 Cooper TG. The tools of biochemistry. A Wiley-Interscience Pub. John Wiley and Sons, New York;
385 1977
- 386 Cova AMW, Neto ADA, Gheyi HR, Ribas RF, de Oliveira LB, Menezes RV. Gas exchange,
387 Chlorophyll Fluorescence and Pigments of Noni (*Morinda citrifolia* L.) under Salt Stress. Journal of
388 Agricultural Science, 2018; 10 (2):318-327.
- 389 Cramer GR, lauchli A, Epstein E. Displacement of Ca^{+2} by Na^{+} from the plasmalemma of root cells. A
390 primary response to salt stress. Plant Physiol., 1985; 79:207-211
- 391 Cramer GR, Epstein E, lauchli A. Na-Ca interactions in barley seedlings: Relationship to ion transport
392 and growth. Plant Cell. Environ., 1989; 12:551-558.
- 393 Delfine S, Loreto F, Pinelli P, Tognetti R, Alvino A. Isoprenoids content and photosynthetic limitations
394 in rosemary and spearmint plants under water stress. Agric. Ecosyst. Environ., 2005; 106, 243–252.
- 395 Egyptian Pharmacopoeia. Egyptian pharmacopoeia, general organization for government. Printing
396 Offices, Ministry of Health, Cairo, Egypt, pp 31-33. 1984
- 397 El-Saedy MAM, El-Sayed MEA, Hammad SE. Efficacy of boron, silicon, jojoba and four bio-products
398 on controlling *Meloidogyne incognita* infecting thompson seedless grapevines. American-Eurasian J.
399 Agric. & Environ. Sci., 2015; 15 (9): 1710-1720, 2015. DOI: 10.5829/idosi.ajeaes.2015.15.9.12779
- 400 Epstein E. Dual pattern of ion absorption by plant cells and by plants. Nature, 1966; 212:1324-1327
- 401 Fang Z, Bouwkamp J, Solomos T. Chlorophyllase activities and chlorophyll degradation during leaf
402 senescence in non-yellowing mutant and wild type of *Phaseolus vulgaris* L. J. Exp. Bot., 1998;
403 49:503–510.
- 404 Farouk S, Youssef SA, Ali AA. Induction of systemic resistance in tomato against *Alternaria solani* by
405 biostimulants and vitamins. Alex. J. Agric. Res. 2012; 57 (1): 117-129.
- 406 Farouk S, Ramadan AA. Improving growth and yield of cowpea plant by foliar application of chitosan
407 under water stress. Egyptian journal of biology, 2012; vol. 7:14-26.
- 408 Farouk S, Abdul Qados AMS. Osmotic adjustment and yield of cowpea in response to drought stress
409 and chitosan. Indian Journal of Applied Research, 2013; 3(10): Article DOI :
410 10.15373/2249555X/October2013/155
- 411 Farouk S, Abo EL-Kheer AM, Sakr MT, Khafagy MA. Osmoregulators and plant growth substances as
412 a growth inducer for pea plants under salinity levels. Int. J. of Agr. Plant Prod., 2011; 2(4):168-180.
- 413 Fedina IS, Tsonev T, Guleva EI. The effect of pre-treatment with proline on the responses of *Pisum*
414 *sativum* to salt stress. Photosynthetica 1993; 29:521-527.
- 415 Foyer CH, Lelandais M, Kunert KJ. Photooxidative stress in plants. Physiol. Plant., 1994; 92: 696-717.
- 416 Frota JNE, Tucker TC. Absorption rates of ammonium and nitrate by red kidney beans under salt and
417 water stress. Soil sci. Soc. Am. 1978; 42:752-755.
- 418 Garg N, Manchanda G. Role of Arbuscular Mycorrhizae in the alleviation of ionic, osmotic and
419 oxidative stresses induced by salinity in *Cajanus cajan* (L.) Millsp. (pigeonpea). J. Agron. Crop Sci.,
420 2009; 195: 110 - 123.
- 421 Gedroits KK. Chemical analysis of soils. Ed. National Sci Foundation, Washington Dc. By The Isreal
422 program for scientific translation, Jerusalem. 1963
- 423 Ghoname AA, EL-Nemr MA, Abdel-Mawgoud, ANR, EL-Tohamy WA. Enhancement of sweet pepper
424 crop growth and production by application of biological, organic and nutritional solutions. Res. J.
425 Agric. Biol. Sci., 2010; 6(7): 349-355.
- 426 Gomez KA, Gomez AA. Statistical procedures for agricultural research, 2 edn . John Wiley & Sons,
427 New York, NY, 680 p. 1984
- 428 Gorham J, Wyn RG, McDonnell E. Some mechanisms of salt tolerance in crop plants. Plant and Soil
429 1985; 89:15-40.

- 430 Gornik K, Grzesik M, Duda BR. The effect of chitosan on rooting of grapevine cuttings and on
431 subsequent plant growth under drought and temperature stress. J. Fruit Ornamental Plant Res, 2008;
432 16: 333-343
- 433 Grattan SR, Grieve CM. Mineral nutrient acquisition and response by plants grown in saline
434 environments. In: Pessarakli M, editor. Handbook of plant and crop stress. Marcel Dekker, Inc, New
435 York. 1994
- 436 Greenway H, Hughes PG, Klepper B. Effect of water deficit on phosphorous nutrition of tomato plants.
437 Physiol. Plant. 1969; 22:199-207.
- 438 Guan Y, Hu J, Wang X, Shao C. Seed priming with chitosan improves maize germination and
439 seedling growth in relation to physiological changes under low temperature stress. Journal of Zhejiang
440 University Science B, 2009; 10(6):427-433.
- 441 Hall HK, McWha JA. Effect of abscisic acid on growth of wheat "*Triticum aestivum*". Annals of Botany,
442 1981; 47:427-433.
- 443 Hawkins HJ, Lewis OAM. Effect of NaCl salinity, nitrogen form, calcium and potassium concentration
444 on nitrogen uptake and kinetics in *Triticum aestivum* L. cv. Gamtoos, New Phytol., 1993; 124:171-177.
- 445 Hazrati S, Tahmasebi-Sarvestani Z, Mokhtassi-Bidgolia A, Modarres-Sanavy SAM, Mohammadib
446 H, Nicolaca S. Effects of zeolite and water stress on growth, yield and chemical compositions of *Aloe*
447 *vera* L.. Agric. Water Manag., 2017; 181: 66–72
- 448 He T, Cramer GR. Abscisic acid concentrations are correlated with leaf area reductions in two salt-
449 stressed rapid-cycling *Brassica* species. Plant and Soil 1996; 179:25-33.
- 450 Heelebust JA. Osmoregulation. Annual Review of Plant Physiology, 1976;27:485-505.
- 451 Jaleel CA, Manivannan P, Lakshmanan GMA. NaCl as a physiological modulator of proline
452 metabolism and antioxidant *Phyllanthus amarus*. C R Biologies, 2007; 330: 806-813.
- 453 Kalra YP. Handbook of reference method for plant analysis. CRC Press, Washington, DC.;1998.
- 454 Kashem MA, Sultana N, Ikeda T, Hori H, Loboda T, Mitsui T. Alteration of starch-sucrose transition in
455 germinating wheat seed under sodium chloride salinity. J. Plant Biol.,2000; 43:121-127.
- 456 Khan H, Khan AZ, Khan R, Matsue N, Henmi T. Zeolite application affects vegetative phenology of
457 determinate and indeterminate soybean grown on Allophanic soil. International Journal of Agriculture
458 Research,2008; 3:148–154.
- 459 Khan MG, Srivastava HS. Changes in growth and nitrogen assimilation in maize plants induced by
460 NaCl and growth regulators. Biol. Plant., 1998; 41:93 - 99.
- 461 Khan MH, Singha KLB, Panda SK. Changes in antioxidant levels in *Oryza sativa* L. roots subjected to
462 NaCl salinity stress. Acta Physiol Plant 2002; 24:145–148.
- 463 Larcher W, Wagner J., Thammathaworn A. Effects of superimposed temperature stress on in vivo
464 chlorophyll fluorescence of *Vigna unguiculata* under saline stress. J. Plant Physiol., 1990; 136:92-102.
- 465 Latifah O, Ahmed OH, Abdul Majid NM. Enhancing nutrients use efficiency and grain yield of *Zea*
466 *mays* L. cultivated on a tropical acid soil using paddy husk compost and clinoptilolite. Bulgarian Journal
467 of Agricultural Science, 2017; 23 (3):418–428
- 468 Leopold AC, Willing RP. Evidence for toxicity effects of salt on membranes. IN: Staples RC,
469 Toenniessen GH, Editors. Salinity tolerance in plants, John Wiley and Sons, New York; 1984
- 470 Letchamo W, Xu HL, Gosselin A. Photosynthetic potential of *Thymus vulgaris* selections under two
471 light regimes and three soilwater levels. Sci. Hortic., 1999; 62:89-101.
- 472 Lichtenthaler HK, Wellburn AR. Determination of total carotenoids and chlorophylls A and B of leaf in
473 Different Solvents. Biol. Soc. Trans., 1985; 11: 591-592.
- 474 Loggini B, Scartazza A, Brugnoli E, Navari-Izzo F. Antioxidative defense system, pigment
475 composition, and photosynthetic efficiency in two wheat cultivars subjected to drought. Plant Physiol.
476 1999; 119: 1091-1099.
- 477 Lynch J, Lauchli A. Salt stress disturbs the calcium nutrition of barley (*Hordum vulgare* L.). New
478 Phytologist 1985; 99:345-354.
- 479 Mali M, Aery NC. Influence of silicon on growth, relative water contents and uptake of silicon, calcium
480 and potassium in wheat grown in nutrient solution. J. Plant Nutr. 2008; 31, 1867–1876.
- 481 Midegley JM. Drug Development from L. Flavour and Fragrance J., 1988; 20(5): 512-520.
- 482 Moosavi SM, Moosavi SG, Seghatoleslami MJ. Effect of drought stress and nitrogen levels on growth,
483 fruit and essential oil yield of fennel (*Foeniculum vulgare* Mill.). Iranian J. Medicin. Aroma. Plants,
484 2014; 30(3):453-462.
- 485 Mozafar A, Oertli JJ. Root-zone temperature and salinity: interacting effects on tillering, growth and
486 element concentration in barley. Plant and Soil 1992; 139:31-38.
- 487 Munne-Bosch S, Schwarz K, Alegre L. Enhanced formation of tocopherol and highly oxidized
488 abietane diterpene in water-stressed rosemary plants. Plant Physiology, 1999; 121:1047-1052.
- 489 Munns R, Termaat A. Whole plant responses to salinity. Aust. J. Plant Physiology, 1986; 13:143-160.

- 490 Munns R. Comparative physiology of salt and water stress. *Plant Cell Envir.*, 2002; 25: 239 - 250.
- 491 Najafi-Ghiri M. Effects of zeolite and vermicompost applications on potassium release from
- 492 calcareous soils. *Soil & Water Res.*, 2014; 9: 31–37.
- 493 Najafinezhad H, Tahmasebi-Sarvestani Z, Modarres SSAM, Naghavi H. Evaluation of yield and some
- 494 physiological changes in corn and sorghum under irrigation regimes and application of barley residue,
- 495 zeolite and superabsorbent polymer. *Arch. Agron. Soil Sci.* 2014; 61: 891–906.
- 496 New N, Chandkrachang S, Stevens WF. Application of chitosan in Myanmar's agriculture sector, in:
- 497 Proceedings of the Sixth Asia Pacific Chitin and Chitosan Symposium, May23–26, The National
- 498 University of Singapore, Singapore.2004
- 499 Papadopoulos I, Rending VV. Interactive effects of salinity and nitrogen on growth and yield of tomato
- 500 plants. *Plant Soil*, 1983; 73:47-57.
- 501 Petropoulos SA, Dimitra D, Polissiou MG, Passam HC. The effect of water deficit stress on the
- 502 growth, yield and composition of essential oils of parsley. *Sci. Hortic.*, 2008; 15:393–397.
- 503 Possingham JV. Plastid replication and development in the life cycle of higher plants. *Annu Rev Plant*
- 504 *Physiol.*, 1980; 31:113-129.
- 505 Prisco JT, O'Leary JW. Enhancement on intact bean leaf senescence by NaCl salinity. *Plant*
- 506 *Physiology*, 1972; 27:95-100.
- 507 Qin L, Kang W, Qi Y, Zhang Z, Wang N. The influence of silicon application on growth and
- 508 photosynthesis response of salt stressed grapevines (*Vitis vinifera* L.). *Acta Physiologiae Plantarum*,
- 509 2016; 38:68
- 510 Raina JN, Goswami KP. Phenolic compounds produced during decomposition of wheat straw in soils.
- 511 *J. Indian Soc. Soil Sci.*, 1988; 36: 787 -90
- 512 Ratnayake RMRNK, Daundasekera WAM, Ariyaratne HM, Ganenege MYU. Soil application of
- 513 potassium silicate reduces the intensity of downy mildew in bitter melon (*Momordica charantia* L.)
- 514 leaves. *Ceylon Journal of Science*, 2016; 45(1):23-31 DOI: <http://dx.doi.org/10.4038/cjs.v45i1.7361>
- 515 Reháková M, Čuvanová S, Dzivák M, Rimár J, Gaval'ová Z. Agricultural and agrochemical uses of
- 516 natural zeolite of the clinoptilolite type. *Curr. Opin. Solid State Mater. Sci.*, 2004; 8, 397–404.
- 517 Romero JM, Marañón T. Long-term responses of *Melilotus segetalis* to salinity. I. Growth and
- 518 partitioning. *Plant Cell Env.*, 1994; 17:1243 - 1248.
- 519 Sharma PK, Hall DO. Interaction of salt stress and photoinhibition on photosynthesis in barley and
- 520 Sorghum. I. *Plant Physiology*, 1991; 138: 614-619.
- 521 Shi Y, Zhang Y, Yao H, Wu J, Sun H, Gong H. Silicon improves seed germination and alleviates
- 522 oxidative stress of bud seedlings in tomato underwater deficit stress. *Plant Physiol. Biochem.*, 2014;
- 523 78:27–36.
- 524 Silberbush M, Ben-Asher J. The effect of salinity on parameters of potassium and nitrate uptake of
- 525 cotton. *Comm. Soil Sci. Plant Anal.*, 1987; 18:65-81.
- 526 Storey R, Pitman MG, Stelzer R. X-Ray microanalysis of cells and cell compartments of *Atriplex*
- 527 *spongiosa*. II Root. *J. Experimental Botany*, 1983; 34:1196-1206.
- 528 Taiz L, Zeiger E. Cytokinins. In: Taiz L, Zeiger E, editors. *Plant Physiology*. Benjamin/Cummings,
- 529 Redwood City, CA.; 1991.
- 530 Torres-Schuman S, Goday GA, PintorTors JA, Moreno FJ, Rodrigo RM, Garcia-Herdugo G. NaCl
- 531 effects on tomato seed germination, cell activity and ion allocation. *J. Plant Physiology* 1989; 135:228-
- 532 232.
- 533 Tuna AL, Kaya C, Higgs D, Murillo-Amador B, Aydemir S, Girgin AR. Silicon improves salinity
- 534 tolerance in wheat plants. *Environ. Exp. Bot.*, 2008; 62:10–16.
- 535 Xing K, Zhu X, Peng X, Qin S. Chitosan antimicrobial and eliciting properties for pest control in
- 536 agriculture: a review. *Agron. Sustain. Dev.*, 2015; 35:569–588.
- 537 Yeo AR, Flowers TJ. Varietal differences in the toxicity of sodium ions in rice leaves. *Physiologia*
- 538 *Plantarum*, 1983; 59:189-195.
- 539 Young A, Britton G. Carotenoids and stress. In: Alscher RG, CummingsJR, editors. *Stress responses*
- 540 *in plants, adaptation and acclimation mechanisms*. Wiley liss, New York; 1990.
- 541 Zaghloul, RA, Mohamed YFY, El-Meihy RM. Influential Cooperation between Zeolite and PGPR on
- 542 Yield and Antimicrobial Activity of Thyme Essential Oil. *International Journal of Plant & Soil Science*,
- 543 2016; 13(1): 1-18
- 544 Zanjani KE, Rad AHS, Naeemi M, Aghdam AM, Taherkhani T. Effects of zeolite and selenium
- 545 application on some physiological traits and oil yield of medicinal pumpkin (*Cucurbita pepo* L.) under
- 546 drought stress. *Curr. Res. J. Biol. Sciences*, 2012; 4(4): 462-470
- 547 Zehtab-Salmasi S, Javanshir A, Omidbaigi R, Aly-Ari H, Ghassemi-Golezani K. Effects of water
- 548 supply and sowing date on performance and essential oil production of anise (*Pimpinella anisum* L.).
- 549 *Acta Agron. Hung.*, 2001; 49, 75–81