

1 **CHARACTERIZATION OF ENGINEERING PROPERTIES**
2 **(PHYSICAL PROPERTIES) OF RUBUS FRUTICOSUS**
3

4 **ABSTRACT**

5 Some engineering properties of *Rubus Fruticosus* fruits and nuts and were characterized in order to provide fundamental information
6 about their properties that will aid in designing modern technology for their handling, processing, storage, preservation, quality
7 evaluation, distribution and marketing. The engineering properties studied are some physical property of *Rubus Fruticosus* fruits and
8 nuts, (moisture content, size and shape of the fruits and nuts) The fruits and nuts were conditioned to five and three different moisture
9 and three and four different temperature levels, respectively before testing. American Standards for Testing and Materials (ASTM)
10 and American Society of Agricultural and Bioresources Engineering (ASABE) standard procedures were used to test all the properties
11 considered. Genstat, Mathlab, JMP in SAS, Duncan in SPSS and Microsoft excel statistical packages were used to analyze the
12 generated data and the means were compared using the analysis of variance (ANOVA) at 5% level of probability. The results obtained
13 revealed that the major, intermediate and minor diameters of the fruits range 35.14 – 34.09 mm, 22.44 – 20.25 mm and 22.03 – 19.92
14 mm respectively for long and, 26.66 – 23.05 mm, 16.69 – 13.99 mm and 16.17 – 13.48 mm respectively for short fruits. The average
15 geometric mean diameter (GMD) of the fruit ranged from 25.90 to 23.95 mm for long and 19.13 to 16.27 mm for short fruits at
16 moisture range of 40.91 – 11.03% wet basis. The nuts also ranged from 36.75 – 37.36 mm, 13.19 – 13.66 mm and 12.89 – 13.18 mm
17 respectively for long nuts, 23.26 – 23.87 mm, 10.93 – 11.24 mm and 10.76 – 11.04 mm respectively for short nuts at 10.2% - 25.06 %
18 wet basis. The fruits and nuts are oblong in shape.

19 **Keywords:** Moisture content, shape and size, biomaterial, physical properties.
20

21 1.0 INTRODUCTION

22 Since man started discovering and cultivating various types of food, there has never been food without work or work for abundant
23 food without machines. The effectiveness of these machines for mechanizing agricultural production depends on adequate knowledge
24 of engineering properties of the products to be mass produced. Mechanization involves replacing human and animal labour with
25 mechanical devices in crop production, processing, storage and distribution. It reduces production cost, ensures timeliness, and
26 optimizes and protects product quality (Adamade and Jackson, 2014).

27 In handling and processing of agricultural products, some fundamental information about their characteristics is essentially needed.
28 These information can be obtained through the knowledge of engineering properties of the products which constitutes essential data in
29 designing and developing modern technologies for their production, handling, processing, storage, preservation, quality evaluation,
30 distribution and marketing.

31 Engineering properties of agricultural product are profitably used for mechanizing their planting, harvesting, drying, processing and
32 storage. They improve working efficiency of processing equipment, reduces losses and waste of constructional materials and, saves
33 time and money. It also helps to maintain quality even in adverse storage and handling conditions and offer ways in which products
34 can be utilized effectively.

35 In recent time, strong growing interest on tree crop for food, money and medicine has been ongoing. This is because there is high
36 demand of food due to effects of development and increasing population, besides, many economic tree crops are fading away without
37 being harnessed and replaced.

38 Tree crops are those perennial woody plants with a single elongated stem of about 3m high and above (Orwa *et al.*, 2009) and,
39 have head of branches and foliage on which fruits grow. The fruits of tree crops are of great interest to food scientists, food producers
40 and other scientists who work towards achieving food security. Modern agriculture has led to handling and processing of agricultural
41 products into more useful product through various unit operations like cleaning, grading, sorting, drying, dehydration, storage, milling
42 and transportation. *Rubus fruticosus* fruit is an edible fruit from *Rubus Fruticosus* tree. It is eaten boiled or fresh for its nutritional and
43 medicinal values.

44 Processing of this important fruit is still by conventional method which encourages losses of both oil and kernel, is unhygienic
45 and subjects the fruits to vagaries of heat treatment which results in poor quality oil. Olawale (2012) reported that the extraction of oils
46 from eleme pulp and kernel is not being carried out at commercial level at present, despite ready availability of the fruit in large
47 quantity in Nigeria and elsewhere in Sub-Sahara Africa. This situation would improve if data needed for the design and operation of
48 the oils' extraction plants are available. *Rubus fruticosus*
49 nuts which house the kernel are usually thrown away after eaten the mesocarp, causing environmental pollution and loss of biomass
50 resources for alternative energy generation. These are as a result of limited knowledge of engineering characteristics of this important
51 fruit and nuts that will promote mechanization of its processing into other useful products.

Engineering properties of agricultural products includes physical, thermal, electrical, optical and mechanical properties. Here physical properties – moisture content, shape and size are treated in this paper. Physical properties are those attributes that reveal the physical features of agricultural product, which can affect mechanization of their handling, processing and storage. Such properties are moisture, shape and size, solid and bulk density, porosity, surface area and volume, angle of repose and absorption which determine the behaviour of the products during handling and processing operations (Mohsenin, 1986, Stroshine, 1998). These properties do change during processing operation and should be considered during design to avoid processing failure.

The objective of this study is to determine the physical: moisture content, size and shape of *Rubus fruticosus* fruit and nuts and to establish relationship between samples size, shape, moisture content of the bio-material.

2.0 MATERIALS AND METHODS

2.1 Crop Materials (*Rubus fruticosus*)

The *Rubus fruticosus* products used for this experiment are the long and short *Rubus fruticosus* fruits and nuts. They were sourced from Ebonyi States of Nigeria. The fruits and nuts are shown in Plate 1.



Plate 1: *Rubus fruticosus* short nuts

2.2 Physical Properties of *Rubus fruticosus* Fruits And Nuts

2.3 Moisture content

The moisture content of the fruits was determined by oven drying method for the fruits and rewetting methods and oven drying for the nuts. The initial moisture content of the fruits was allowed to be uniform by placing them in a refrigerator at 5°C for about 18 hours. The moisture content of the fruits was determined at average environmental temperature and relative humidity of 38°C and 77% respectively. The fruit samples were put in a clean moisture can and weighed using digital balance (Model APX-1502, Denver Instrument US) of 0.01g accuracy before putting them in the oven. The oven was

74 set at temperature of 105°C for 8 hours. The difference between the initial and final weight of each sample was used to
75 calculate the moisture content of the sample as Eq 2.1 (Mohsenin 1986):

$$76 \quad \% M_{wb} = \frac{W_1 - W_2}{W_1} \times \frac{100}{1} \quad 1$$

77 *Where:* M_{wb} = moisture content wet basis (%), W_1 = initial weight (g). W_2 = final weight (g).

78 **2.4 Moisture content of agricultural products**

79 This is the amount of water present in any material which can be expressed as a percentage of the material weight either in wet basis
80 or dry basis. It is calculated as, Eq 1.1 (Mohsenin, 1986):

$$81 \quad \% M_{wb} = \frac{w_1 - w_2}{w_1} \times \frac{100}{1} \quad 1.1$$

82 *where:* M_{wb} = moisture content wet basis (%), w_1 = dry weight (g), w_2 = final weight (g).

83 **2.5 Shape and size of agricultural products**

84 Shape is a physical attribute of agricultural products that is used to differentiate between cultivars, study the heritage shape trait,
85 evaluate the consumer decision performance, package and transport product (Jahns *et al.*, 2001, Sadrnia *et al.*, 2007). Shape can be
86 described objectively (depending on the nature of the product) using indices calculated from geometrical characteristics of the product.
87 Such indices include roundness, roundness ratio (RR), sphericity and axial ratio (AR) (Mohsenin, 1986, Wilhelm *et al.*, 2005, Venora
88 *et al.*, 2007). Roundness measures the consistency of the product diameter at a number of different orientations). Roundness ratio is
89 non-dimensional physical parameter that uses the geometrical attribute of the product to determine its shape (Rashidi and Seyfi, 2007).
90 Sphericity is defined as the ratio of the volume of a triaxial ellipsoid with diameters equal to the major, intermediate and minor
91 diameters to the volume of sphere which circumscribed the object. Aspect ratio is also a dimensionless physical quantity that describes
92 the shape of an object (Sadrnia *et al.*, 2007).


93
94 For nuts and nutshells, a predetermined amount of distilled water was added to the sample before placed it in a refrigerator
95 set at 5 °C for about 20 hours to equilibrate. The quantity of water added was calculated as in Eq. 2.2 (Mohsenin, 1986):

$$96 \quad W_2 = W_1 \left(\frac{M_1 - M_2}{100 - M_1} \right) \quad 2$$

97 Where W_1 = sample weight, W_2 = amount of distilled water to be added, M_1 = final moisture content, M_2 = initial moisture
 98 content.
 99 Five different moisture content (M_1 , M_2 , M_3 , M_4 and M_5) wet basis were obtained for the fruits but the nuts and nutshell
 100 were conditioned to three moisture levels (M_1 , M_2 , M_3). Each moisture content was replicated three times.

101 2.4 Size and Shape of Fruits and Nuts

102 Size was determined by measuring the principal dimensions of the products such as length, width and thickness which are also known
 103 as major, intermediate and minor diameters, Fig. 1.



104
 105 Figure 1: Principal dimensions of *Rubus fruticosus* fruits (a) major diameter, (b) intermediate diameter and (c) minor diameter.

106
 107 Digital vernier caliper (Mitutoyo, Japan) of 0.01g accuracy was used to measure the principal dimensions of the fruits and nuts at
 108 various moisture levels.

109 Various mean diameters and shapes of the fruits and nuts were calculated using Eqs. 3 – 7 Mohsenin (1986) and Ehiem *et al.*(2012):

$$110 \quad GMD = \sqrt[3]{abc} \quad 3$$

$$111 \quad AMD = \frac{a+b+c}{3} \quad 4$$

$$112 \quad SMD = \sqrt{\frac{(a \times b) + (b \times c) + (c \times a)}{3}} \quad 5$$

$$EQD = \frac{SMD+GMD+AMD}{3} \quad 6$$

$$AR = \frac{b}{a} \quad 7$$

Where: a = major diameter, b = intermediate diameter, c = minor diameter, GMD = geometric mean diameter, AMD = arithmetic mean diameter, SMD = square mean diameter, EMD = equivalent diameter; AR = aspect ratio.

Roundness of the fruits were determined by tracing the edges of the fruits with sharp pencil on a white paper. The diameter of the sharp projected areas were measured with a pair of divider and digital vernier caliper while roundness of each fruit and nut was calculated as Eq 8.

$$Roundness = \frac{(A_p)}{(A_c)} \quad 8$$

Where; A_p = Largest projected area of the product when in natural rest position; A_c = Area of smallest circumscribed circle.

Roundness ration (RR) was determined as shown in Eq. 9:

$$RR = \frac{a}{\sqrt{bc}} \quad 9$$

Sphericity was determined by applying the data of principal dimensions to Eq. 10.

$$Sphericity = \sqrt[3]{abc}/a \quad 10$$

Where a = major diameter, b = intermediate diameter, c = minor diameter

The roundness and sphericity of each of the five and three moisture content studied were replicated fifty times.

3.0 RESULTS AND DISCUSSIONS

3.1 Physical Properties Of *Rubus fruticosus* Fruits And Nuts

130 3.2 Size of *Rubus fruticosus* fruits and nuts

131 The principal dimension of long and short *Rubus fruticosus* fruits and nuts is presented in Table 1. The principal dimensions of long
132 and short fruits evaluated decreased as moisture content decreased from 40.91 to 11.03% (*w.b.*). The major, intermediate and minor
133 diameters decreased by 2.65%, 15.71% and 15.24% respectively for long fruit and, 11.56%, 16.18% and 16.64% respectively for short
134 fruit. Behise *et al.* (2013) and Aghajani *et al.* (2012) observed 10.74%, 6.44% and 11.52% and, 2.76%, 13.38% and 9.29% decrease in
135 major, intermediate and minor diameters of sunflower seeds and barley fruits (Sahra) respectively as moisture content decreased from
136 18% - 10% and 42.17% - 10.12% wet basis.

137 Major, intermediate and minor diameters of the nuts increased by 1.63%, 3.51% and 2.20% respectively for long nut and 2.56%,
138 2.76% and 2.54% respectively for short nut as moisture content increased from 10.20% - 25.06% wet basis. Major, intermediate and
139 minor diameters of tiger nut, pistachio nut and dika nut were also reported to increase by 15.16%, 14.13% and 12.89% respectively;
140 18.04%, 20.10% and 5.930% respectively and, 1.21%, 6.62% and 30.81% respectively as moisture content increased from 17% - 27%
141 (wb); 3.70% - 30% (wb) and 8.74% - 13.75% (wb) respectively (Abano and Amoah, 2011, Peyman *et al.*, 2013 and Orcheba *et al.*,
142 2013 respectively). The observed increase in principal dimensions could be because the cells of the fruits and nuts expanded at higher
143 moisture content.

144 Table 1: Principal dimensions of *Rubus fruticosus*. long and short fruits and nuts at different moisture content.

M. C.	Fruits					
	Long			Short		
	a	b	c	A	b	c
40.91%	42.86 ± 0.91 (2.13)	20.05 ± 0.66 (3.27)	19.56 ± 0.65 (3.33)	26.06 ± 1.09 (4.18)	16.69 ± 0.59 (3.52)	16.17 ± 0.27 (1.98)
34.92%	42.59 ± 0.88 (2.09)	18.54 ± 0.63 (3.41)	18.06 ± 0.64 (3.56)	25.08 ± 1.37 (5.47)	15.89 ± 0.56 (3.56)	15.36 ± 1.54 (1.05)
23.44%	42.36 ± 0.93 (2.09)	18.44 ± 0.51 (2.77)	17.89 ± 0.53 (2.97)	24.20 ± 2.08 (8.62)	15.05 ± 0.78 (5.24)	14.57 ± 1.54 (1.05)
18.5%	42.02 ± 0.97 2.30	17.39 ± 2.38 (1.36)	17.18 ± 0.98 (3.74)	23.87 ± 2.24 (9.39)	14.45 ± 0.84 (5.80)	13.94 ± 1.59 (1.14)
11.03%	41.71 ± 1.36	16.90 ± 2.43	16.58 ± 1.13	23.05 ± 2.73	13.99 ± 0.96	13.48 ± 1.62

	(3.27)	(1.44)	(3.82)	(1.18)	(1.20)	(1.20)
	Nuts					
10.20%	36.75 ± 1.27 (3.46)	13.19 ± 0.45 (3.41)	12.89 ± 0.39 (3.03)	23.26 ± 0.92 (3.96)	10.93 ± 0.38 (3.48)	10.76 ± 0.35 (3.25)
17.23%	36.91 ± 1.73 (4.69)	13.26 ± 0.37 (2.79)	12.99 ± 0.36 (2.77)	23.59 ± 0.89 (3.77)	11.18 ± 0.45 (4.03)	10.98 ± 0.42 (3.83)
25.06%	37.36 ± 1.27 (3.39)	13.67 ± 0.46 (3.37)	13.18 ± 0.39 (2.96)	23.87 ± 0.92 (3.84)	11.24 ± 0.45 (4.04)	11.04 ± 0.43 (3.89)

The values in bracket are the coefficient of variation (%CV); a = major diameter (mm); b = intermediate diameter (mm); c = minor diameter (mm); M.C. = moisture content wet basis.

Lower percentage variation of principal dimensions of the nuts with moisture content when compared with that of tiger nut and pistachio nut could be because the cell of the *Rubus fruticosus* nuts have less void spaces and shrinking behavior during moisture loss than tiger nut and pistachio nut. The principal dimensions of both fruits and nuts had low coefficient of variation which indicates evenly distributed values about their means. The mean diameters also decreased as moisture content decreased from 40.91 to 11.03% (w.b.) for fruits and 25.06% - 10.20% wet basis for nuts (Table 2). The geometric mean diameter (GMD), arithmetic mean diameter (AMD), square mean diameter (SQMD) and equivalent mean diameter (EQD) of the fruits decreased by 7.73%, 6.75%, 7.03% and 7.17% respectively for long fruit and, 14.95%, 14.26%, 14.66% and 14.60% respectively for short fruit. Long and short *Rubus fruticosus* nuts also decreased in GMD, AMD, SQMD and EQD by 2.44%, 2.10%, 2.37% and 2.29% respectively and 2.65%, 2.69%, 2.64% and 2.63% respectively. Davies and Zibokere (2011) has also reported 26.79%, 22.09% and 20.11% increase in GMD of Ife Brown, IAR-339-1 and IT86G-1010 cowpea varieties, respectively as their moisture content decreased from 30% - 15% wet basis. The ANOVA in Table 3 revealed that moisture content had significant (5%) effect on the principal dimensions of both fruits and nuts while the effect of interaction is not significant (5%). Besides, the differences between the principal dimensions of long and short fruits are highly significant (5%). Regression equations of the principal dimensions in Table 4 showed high values of coefficient of determination (R^2) indicating good relationship between the parameters evaluated.

Table 2: Mean diameters of long and short *Rubus fruticosus* fruits and nuts at different moisture content.

165

M. C.	Fruits							
	GMD	Long AMD	SQMD	EQD	GMD	Short AMD	SQMD	EQD
40.91%	25.90 ± 0.64 (1.61)	26.54 ± 0.66 (1.62)	26.19 ± 0.65 (1.61)	26.21 ± 0.65 (1.61)	19.13 ± 1.09 (5.09)	19.64 ± 0.92 (4.67)	19.37 ± 0.98 (5.05)	19.38 ± 0.99 (5.10)
34.92%	25.23 ± 0.63 (2.48)	25.91 ± 0.63 (2.44)	24.54 ± 0.63 (2.47)	25.56 ± 0.63 (2.46)	18.26 ± 1.06 (3.81)	18.78 ± 0.63 (3.33)	18.49 ± 0.96 (3.18)	18.51 ± 0.83 (4.16)
23.44%	24.69 ± 0.57 (2.33)	25.41 ± 0.59 (2.31)	25.01 ± 0.58 (2.31)	25.04 ± 2.32 (2.32)	17.41 ± 1.25 (4.17)	17.94 ± 0.80 (4.47)	17.65 ± 1.18 (4.39)	17.67 ± 1.00 (4.68)
18.50%	24.36 ± 0.55 (2.25)	25.11 ± 0.55 (2.18)	24.69 ± 0.55 (2.21)	24.72 ± 0.55 (2.21)	16.84 ± 1.34 (4.92)	17.42 ± 0.87 (4.98)	17.10 ± 1.26 (4.36)	17.12 ± 1.08 (4.28)
11.03%	23.95 ± 0.54 (2.24)	24.75 ± 0.54 (2.17)	24.31 ± 0.54 (2.20)	24.33 ± 0.54 (2.19)	16.27 ± 1.47 (3.04)	16.84 ± 1.41 (3.62)	16.53 ± 1.43 (3.65)	16.55 ± 1.26 (3.62)
Nuts								
10.20%	18.42 ± 0.47 (2.55)	20.95 ± 0.57 (2.72)	19.39 ± 0.49 (2.50)	19.59 ± 0.51 (2.60)	13.98 ± 0.43 (3.08)	14.98 ± 0.48 (3.20)	14.39 ± 0.45 (3.20)	14.45 ± 0.45 (3.11)
17.23%	18.52 ± 0.53 (2.86)	21.05 ± 0.71 (3.37)	19.50 ± 0.59 (3.03)	19.69 ±0.61(3.10)	14.25 ± 0.47 (3.29)	15.25 ± 0.49 (3.21)	14.67 ± 0.48 (3.27)	14.72 ± 0.48 (3.26)
25.06%	18.88 ± 0.52 (2.75)	21.40 ± 0.61 (2.85)	19.86 ± 0.55 (2.77)	20.05 ± 0.56 (2.79)	14.36 ± 0.48 (3.34)	15.38 ± 0.51 (3.32)	14.78 ± 0.49 (3.32)	14.84 ± 0.49 (3.30)

166 The values in bracket are the coefficient of variation (%CV);

167

168 Table 3: ANOVA summary of physical parameters of *Rubus fruticosus* fruit studied at moisture range of

169 40.91% - 11.03% wet basis.

Source of variation	df	Fruits				df	Nut			
		a	b	c	5%		a	b	c	5%
Size	1	15094.87 ^{**}	713.12 ^{**}	1438.55 ^{**}	3.86	1	7796.08 ^{**}	2145.59 ^{**}	2301.78 ^{**}	3.87
Moisture content	4	23.78 [*]	79.26 ^{**}	151.40 ^{**}	2.39	2	7.57 [*]	23.56 [*]	18.28 [*]	3.03
Interaction	4	6.59 [*]	0.8593 ^{ns}	0.6308 ^{ns}	2.39	2	0.2544 ^{ns}	1.99 ^{ns}	0.4459 ^{ns}	3.03

170 *a* = major diameter (mm); *b* = intermediate diameter (mm); *c* = minor diameter (mm); ^{**} Highly significant; ^{*} Significant;
 171 *ns*

172 *Not significant*

173

174 Table 4: Equations representing relationship between principal dimensions and moisture content

Parameters	Fruit			
	Long		Short	
	Regression equation	R ²	Regression equation	R ²

a	$3.59 h + 41.39$	0.97	$9.14 h + 22.13$	0.97
b	$9.24 h + 15.92$	0.99	$8.71 h + 13.01$	0.99
c	$8.48 h + 15.70$	0.89	$8.64 h + 12.51$	0.99
Nut				
a	$4.14 h + 36.28$	0.95	$4.09 h + 22.86$	0.99
b	$3.27 h + 12.80$	0.87	$0.3533 \ln(h) + 11.76$	0.94
c	$1.96 h + 12.67$	0.98	$0.3184 \ln(h) + 11.50$	0.98

a = major diameter (mm); b = intermediate diameter (mm); c = minor diameter (mm); h = moisture content wet basis

175

176

177

In design and development of processing and handling equipment for *Rubus fruticosus* products, moisture content of the fruits and nuts should be specified to avoid waste of money and material due to over /under design.

3.3 Shape of *Rubus fruticosus* fruits and nuts

3.4 Sphericity, roundness and aspect ratio of the fruits and nuts

The sphericity, roundness and aspect ratio of *Rubus fruticosus* long and short fruits decreased with increase in moisture content (40.91% to 11.03% wet basis) while the long and short nuts increased with increase in moisture content (10.20% - 25.06% wet basis). The sphericity, roundness and aspect ratio of short and long fruits reduced by 5.27%, 3.80% and 45.10% and, 4.65%, 6.98% and 4.62%, respectively (Table 5 and Fig 3). Fathollahzadeh *et al.* (2008) and Razavi *et al.* (2009) also reported that the sphericity of barberry fruits and canola seeds (*SLM spp.*) reduced by 40.43% and 2.29% as moisture content reduced from 89.23 – 11.64% and 21 – 6% wet basis respectively. At 40.91% wet basis, the sphericity and aspect ratio values showed that short fruits are more spherical and oblong than long fruit; hence, short fruit can roll more easily than the long fruit

Aspect ratio (AR), sphericity and roundness ratio (RR) of the nuts at 10.20% moisture content wet basis ranged from 2.8 – 3.09, 1.69 – 1.88 and 0.4700–0.5300 respectively for long nut and, 1.88 – 2.32, 1.20 – 1.45 and 0.5700–0.6600 respectively for short nut. The mean values of shape parameters shown in Table 5 and Fig 3 revealed that AR for both nuts are within the range for oblong shape (1.85 – 2.31) as classified by Şehrali (1988) and Kara *et al.* (2013). The values of RR and sphericity being slightly higher than 50% confirmed the shape. *Rubus fruticosus* nuts had similar shape with Aras-98, Elkoca-05, Noyanbey-98 and Yakutiye-98 bean cultivars as reported by Kara *et al.* (2013) while Tenera and Dura palm nuts are ovate in shape (Akinoso and Raji, 2010).

Table 5: Aspect ratio of long and short *Rubus fruticosus* fruits and nuts at different moisture content

Sizes	Moisture content (wb)				
	40.91 %	34.92 %	23.44 %	18.50 %	11.03 %
Long fruit	46.78 ± 1.67 (3.57)	45.23 ± 1.43 (3.16)	43.47 ± 1.43 (3.29)	41.36 ± 5.72 (3.38)	40.56 ± 5.94 (3.02)
Short fruit	64.09 ± 2.14 (3.34)	63.52 ± 2.97 (4.68)	62.49 ± 4.27 (4.83)	60.86 ± 4.32 (0.09)	60.71 ± 4.88 (4.73)

	10.02%	17.23%	25.06%	NS	NS
Long nut	2.79 ± 0.09 (3.23)	2.77 ± 0.13 (4.69)	2.74 ± 0.09 (3.29)	NS	NS
Short nut	2.13 ± 0.09 (4.23)	2.11 ± 0.08 (3.79)	2.11 ± 0.09 (4.27)	NS	NS

The values in bracket are the coefficient of variation (% CV); The fruits aspect ratio values are in percentage. NS = not studied

F-LSD (5%) of difference between two sizes of Rubus fruticosus fruit and nuts means respectively = 0.0252 and 0.0418

F-LSD (5%) of difference between moisture levels of Rubus fruticosus fruit nuts means respectively means = 0.0398 and 0.0625




Fig. 3a




Fig. 3b

Figure 3: The plot of (a) sphericity and (b) roundness of *Rubus fruticosus* fruits against moisture content.




Fig. 4a




Fig. 4b

Figure 4: The plot of (a) sphericity and (b) roundness of *Rubus fruticosus* nuts against moisture content.

All the shape parameters for short fruits had non-linear relationship with moisture content except RR. Long and short fruits are significantly different (5%) in terms of sphericity and aspect ratio (AR) while the nuts differ in terms of AR and RR (Table 4). Moisture content had no effect on the shape parameters of the nut but significantly (5%) affected the AR of the fruit only. This result could be because the cells of the fruits and nuts are inelastic and could have less pore spaces which resulted in low shrinkage effect during moisture loss. The interaction of both fruits and nuts are not significant (5%). Roundness had low values for both fruits and nuts studied, showing that *Rubus fruticosus* fruits and nuts are not round, therefore, the shape of screen opening for separation purposes must not be round. These results mean that the shape of outlet channel of storage facility for *Rubus fruticosus* fruits and nuts should be oblong or rectangular. The rolling nature of the fruits is an indication that agitator device is not needed in the design.

4.1 Conclusions

Some engineering properties of *Rubus fruticosus* fruits, nut and nutshell were studied and the following conclusions were made:

1. The major, intermediate and minor diameters of the fruits range 35.14 – 34.09 mm, 22.44 – 20.25 mm and 22.03 – 19.92 mm respectively for long and, 26.66 – 23.05 mm, 16.69 –

13.99 mm and 16.17 – 13.48 mm respectively for short fruits at moisture range of 40.91 to 11.03% (wet basis).

2. The nuts also ranged from 36.75 – 37.36 mm, 13.19 – 13.66 mm and 12.89 – 13.18 mm respectively for long nuts, 23.26 – 23.87 mm, 10.93 – 11.24 mm and 10.76 – 11.04 mm respectively for short nuts at 10.2% - 25.06 % wet basis.

3. Both fruits and nuts are oblong in shape and denser than water (with average values of 1.05 g/cm³ and 1.69 g/cm³ respectively).

REFERENCES

Abano, E. E. and Amoah, K. K. (2011). Effect of moisture content on the physical properties of tiger nuts (*Cyperus Esculentus*). *Asian Journal of Agricultural Research*, 4(1): 1 – 11.

Adamade, C.A. and Jackson, B.A. (2014). Agricultural mechanization: a strategy for food sufficiency. *Journal of Agricultural Science* 4(3): 152-156.

Aghajani, N., Ansaripour, E. and Kashaninejad, M (2012). Effect of moisture content on physical properties of barley seeds. *Journal of Agricultural. Sci. Tech.* 14: 161-172.

Akinoso, R. and Raji, A.O. (2010). Physical properties of fruit, nut and kernel of oil palm. *Int. Agrophys.*, 25, 85-88.

Barbosa-Canovas, G.V., Ortega-Rivas, E., Juliano, P., Yan, H. (2005). *Food Powders: Physical Properties, Processing, and Functionality*. New York, NY: Kluwer Academic/Plenum Pub.

Bäumler E, Cuniberti A., Nolasco S.M., Riccobene I.C. (2006). Moisture dependent physical and compression properties of safflower seed. *Journal of Food Eng.* 73: 134-140.

Behise S., Kaur A., Manikantan M. R. (2013): Moisture dependant physical properties of sunflower seed (PSH 569). *International Journal of Engineering and Science*, 2: 23-27.

Bilbao-Sainz, C., Andres, A. and Fito, P. (2005). Hydration kinetics of dried apple as affected by drying conditions. *Journal of Food Engineering*, 68, 369 – 376.

Botelho, F. M., Correa, P. C., Martins, A. R., Botelho, S. C. C. and Oliveira, G. H. H. (2013). Effect of mechanical damage on water absorption process by corn kernel. *Journal of Food Science and Technology* (Campinas), 33 (2): 119 – 125.

Boyd, C. E. (1999). *Water quality: An Introduction*. The Netherlands. Kluwer Academic Publishers Group. ISBN 0-7923-7853-9.

Ehiem, J. C., Simonyan, K. J. (2012). Physical properties of wild mango (*Irvingia spp.*) fruits and nuts. *International Agrophysics*, 26: 95 – 98.

- 289 Ela, W. P. (2007). *Introduction to Environmental Engineering and Science*, Prentice Hall,
290 3rd ed. Pp.153 – 159.
- 291 Fathollahzadeh, H. and Rajabipour, A. (2008). Some mechanical properties of barberry.
292 *Int. Agrophysics*, 22: 299-302
- 293
294 Gowen, A. Abu-Ghannam, A, Frias, J. and Oliveira, J. (2007). Influence of pre-blanching on
295 the water absorption kinetics of soyabean. *Journal of Food Engineering*, 78 (3): 965 –
296 971.
- 297
298 Hassan, H. E. (2002). Study of sorting and grading operations of Egyptian mature
299 oranges using visible laser. Ph.D. Th., Nat. Inst. of Laser In Enhanced Sc. (NILES), Cairo
300 U.: 12-15.
- 301
302 Irtwange, S. V. and Igbeka, J. C. (2004). Thermal conductivity of African yam bean
303 (*Sphenostlis Stenocarpa*) as affected by bulk density and moisture content. *Journal of*
304 *Applied Science, Engineering Technology*, 4(2): 1-6.
- 305
306 Jideani, V. A. and Mpotokwana, S. M. (2009). Modeling of water absorption of Botswana
307 Bambara variety using Peleg's equation. *Journal of Food Engineering*. 92 (2): 182 – 188.
- 308
309 Kabukey, A., Herák, D. and Sedláček, A. (2011). Behaviour of different moisture
310 contents of *Jatropha curcas* L. seeds under compressive loading. *Res. Agri. Eng.* 57(2): 72
311 – 77.
- 312 Kara, M., Sayinci, B., Elkoca, E., Öztürk, İ. and Özmen, T. B. (2013). Seed size and
313 shape analysis of registered common bean (*Phaseolus vulgaris* L.) cultivars in turkey
314 using digital photography. *Journal of Agricultural Sciences* 19: 219-234.
- 315
316 Kashaninejad, M and Kashiri, M. (2007). Hydration kinetics and changes in some
317 physical properties of wheat kernel. *Iranian Food Science and Technology Research*
318 *Journal*, 5(1): 47 – 59.
- 319
320 Kawamura S., M. Natsuga, K. Takekura, and K. Itoh (2003). Development of an
321 automatic rice-quality inspection system. Ag. Process Eng. Lab, Grad. Sch. Ag. Sc.,
322 Hokkaido U., Sapporo 060-8589, Japan. *Computers and Electronics in Agriculture* 40:
323 115- 126.
- 324
325 Konak, M., Carman, K. and Aydin, C. (2002). Physical properties of chick pea seeds.
326 *Biosystem Engineering*. 182(1): 73-78.
- 327
328 Licht, S. (2004). Chem. Comm. 4635.
- 329
330 Masoumi A. A., Rajabipoor A., Tabil L., and Akram A. A., (2006), Studies on physical
331 attributes of a garlic, *Journal of Agricultural Science Technology*, 8: 15-23.
- 332
333 Midilli, A., and Kucuk, H. (2003). Mathematical modeling of thin layer drying of

- pistachio by using solar energy. *Energy conversion and Management*, 44(7): 1111-1122.
- Mohsenin, N. N. (1986). *Physical properties of plants and animals materials*. 2nd edition (revised). Gordon and Breach Science publishers. New York.
- Montanuci, F. D., Jorge, L. M. and Jorge, R. M. M. (2013). Kinetic, thermodynamic properties and optimization of barley hydration. *Journal of Food Science and Technology*, Campinas, 33(4): 690 – 698.
- Nelson S. O, Guo W, Trabelsi S, Kays S. (2007). Dielectric spectroscopy of watermelons for quality sensing. *Measurments science Technology*; 18: 1887–1892. 15.
- Ndirika, V.I.O. and Oyeleke, O.O. (2006). Determination of some selected physical properties and their relationship with moisture content for millet (*Pennisetum Glaucum L.*). *Applied Engineering in Agric*. ASABE, vol. 22(2):
- Orwa, C., A. Mutua, R. Kindt, R. Jamnadass, and A. Simons. (2009). Agroforestry Database: a tree reference and selection guide version 4.0. <http://www.worldagroforestry.org/af/treedb/>
- Orcheba, B. A, Idah, P. A, Adebayo, S. E. and Nwankwo, C. C. (2013). Determination of some engineering Properties of dika nut (*Irvingia gabonensis*) at two moisture content levels as relevant to its processing. *International Journal of Engineering Research and Applications (IJERA)*, 3(2), 182-188.
- Olawale, A. S. (2012). Solid-liquid extraction of oils of African elemi's (*Rubus Fruticosus schweinfurthii*'s) fruit. *Agri Eng Int: CIGR Journal*, 14 (2): Manuscript No. 2083.
- Oni, K, C. (2011). Man, machine and food insecurity. The ninety-fourth inaugural lecture, University of Ilorin. Library and Publication Committee, University of Ilorin, Nigeria. 1-70.
- Peleg, M. (1988). An empirical model for the description of moisture sorption curves. *Journal of Food Science*, 53(4): 1249–1251.
- Peyman, L., Mahmoudi, A. and Ghaffari, H. (2013). Some physical properties of pistachio nuts. *International Journal of Agriculture and Crop Sciences*, 5 (7), 704-711.
- Razavi S.M.A., Yeganehzad S., Sadeghi A. (2009). Moisture dependent physical properties of canola seeds. *Journal Agricultural Science Technology*, 11: 309-322.
- Resio, A. C., Aguerre, R. J. and Suarez, C. (2006). Hydration kinetics of amaranth grain. *Journal of Food Engineering*, 72(3): 247 – 253.
- Shafaei, S. M., Masoumi, A. A. and Roshan, H. (2014). Analysis of water absorption of bean and chickpea during soaking using Peleg model. *Journal of the Saudi Society of Agric. Science*, 1 – 10.

- Simonyan, K. J., Yiljep, Y. D., Oyatoyan, O. B. and Bawa, G. S. (2009). Effect of moisture on some physical properties of *Lablab purpureus* (L.) sweet seeds. *Agricultural Engineering International: the CIGR Ejournal Manuscript* 1279. Vol. XI.
- Sirisomboon, P., Potncheloeampong, P. and Romphophek, T. (2007). Physical properties of greensoya bean: Citrus fo sorting. *Journal of Food Engineering*, 79: 18-22.
- Solomon, W.K. and Lewdu, A.D. (2009). Moisture dependent physical properties of Niger (*Guiconia abyssinica* Cass.) seed. *Ind Crops and Prod* 29:165-170
- Turhan, M., Sayar, S. and Gunasekaran, S. (2002). Application of Peleg's model to study water absorption in chickpea during soaking. *Journal of Food Engineering*, 53 (2): 153 – 159.
- Vega-Gálvez, A., Notte-Cuello, E., Lemus- Mondaca, R., Zura, L. and Miranda, M. (2009). Mathematical Modelling of Mass Transfer during Rehydration Process of Aloe vera (*Aloe barbadensis* Miller). *Food Bioprod.Process.*, **87**: 254-260.
- Vishwakarma, R. K., Shivhare, U. S. and Nanda, S. K. (2013). Water absorption kinetics of guar seeds and unhulled guar splits. *Food Bioprocess Technol.* (2013) 6:1355–1364.
- Yaldiz, O., Ertekin, C. and Uzun, H.B. (2001). Mathematical modelling of thin layer solar drying of sultana grapes. *Energy*, 26: 457-465.