

**AMINO ACID PROFILE AND MINERAL CONTENTS OF  
*BALANITES AEGYPTIACA* KERNEL**

**ABSTRACT**

Study on the evaluation of nutritional quality of poorly utilized plant products is of immense importance. This research work therefore evaluate the amino acid profiles and mineral contents of *Balanites aegyptiaca* kernel using standard analytical methods. The amino acid profile analysis revealed the presence of essential and the non-essential amino acids of which Total sulphur and Total aromatic amino acids contents were 2.86g/100g protein and 7.21g/100g protein respectively. Analysis of the mineral content showed the presence of the following minerals in the order; Magnesium > Sodium > Calcium > Potassium > Iron > Zinc > Manganese > Lead > Copper > Nickel > Chromium > Cadmium. The kernel could therefore be used as a good source of protein and minerals in supplementing other sources in order to reduce food insecurity and hence malnutrition from lack of protein diets.

**Keywords:** *Amino acid profiles, Mineral contents, Balanites aegyptiaca, and Kernel.*

**1.0 INTRODUCTION**

*Balanite aegyptiaca* (Linn.) is a genus of flowering plants in the caltrop family, commonly known as desert date. *Banalites aegyptiaca* is an important multipurpose tree found in most African countries [1]. It is a woody evergreen xylophylic tree with a height of 10m grown in various ecological conditions mainly distributed in semi-arid zones in tropical Africa especially in Senegal, Sudan and also Asian countries such as India [2]. In

Arabic, it is known as lalob, hidjihi, inteishit and heglig. In Hausa, it is called Aduwa and in Swahili and Amharic, it is respectively called mchunju and bedena [3].

*Balanite aegyptiaca* is used in treatment of various ailments such as jaundice, intestinal worm infection, malaria, syphilis, epilepsy, dysentery, constipation, haemorrhoid among others [4]. The seed is about 1.5-3cm long, light brown, fibrous and hard, making up to 50-60% of the fruit with about 500-1500 dry clean seeds/kg. These seeds were reported to contain a cytostatic saponins "balanitins" [5], deltonin and isodeltonin which are used as molluscicidal agents [6]. In addition, various reports on the nutritional and anti-nutritional profile of *Balanites aegyptiaca* seeds powder has shown that the seed powder contains a relatively high amount of protein and lipids [7] and some amount of anti-nutritional factors such as tannins, oxalate and phytic acid as compared to other plant products [8]. Antinutritional factors such as phytic acid, tannins, saponins, oxalic acid, have adverse effect on health through inhibition of protein digestion, growth, iron and zinc absorption [9, 10]. Due to the presence of high protein content in *Balanites aegyptiaca* seed kernel as mentioned by [11], there is a need for determination of the amino-acid profile and mineral contents of this kernel.

This study could go a long way in proving the nutritional content of kernel obtained in Kano state, northwest Nigeria. Thus, providing a scientific basis of the use of this seed as source of food in drought areas, war zones as well as commercialization of *Balanites aegyptiaca* which could bring about boosting the economy, encouraging its cultivation and hence solving the problem of malnutrition and food insecurity.

## **2.0 MATERIALS AND METHODS**

## 2.1 Sample Collection and Preparation

*Balanites aegyptiaca* fruits were obtained from Gwammaja market, Dala Local Government Area, Kano state Nigeria. It was authenticated at department of plant biology Bayero University Kano, with an accession number BUKHAN0359. The fruit was processed by soaking in water for 48 hours, de-pulped. The seed obtained were hard cracked using a hammer to obtain its kernel. The kernels were air-dried for 12 hours and pulverized using a mortar and pestle. The pulverized sample was stored in plastic containers.

## 2.2 Analysis of amino acid profile of *Balanites aegyptiaca* Kernel

Amino Acid profile in the sample was determined using the methods of [12]. The sample was dried to constant weight, defatted, hydrolyzed, evaporated in a rotary evaporator and loaded into the Technicon Sequential Multi-Sample Amino Acid Analyzer (TSM).

## 2.3 Analysis of Mineral contents of *Balanites aegyptiaca* Kernel

Elemental analysis of *Balanites aegyptiaca* kernel involves ashing of the solid sample, digestion, dilution and quantitative analysis with atomic absorption spectrophotometer and UV-Visible spectrophotometer [13].

## 3.0 RESULTS

Table 1 shows the amino acid profile of *Balanites aegyptiaca* kernel. The result showed the presence of both essential and non-essential amino acids.

**Table 1: Amino acid profile of *Balanites aegyptiaca* kernel**

Amino acid	Abbreviation	g/100g Protein
Lysine*	Lys	3.52
Histidine*	His	2.19

Arginine	Arg	6.38
Aspartic acid	Asp	8.71
Threonine*	Thr	2.98
Serine	Ser	3.58
Glutamic acid	Glu	13.57
Proline	Pro	2.90
Glycine	Gly	4.40
Alanine	Ala	3.94
Cystine	Cys	1.52
Valine*	Val	4.02
Methionine*	Met	1.34
Isoleucine*	Ile	3.29
Leucine*	Leu	6.69
Tyrosine	Tyr	2.98
Phenylalanine*	Phe	4.23

\*Essential amino acids and their corresponding percentages when compared to the total amino acids analyzed.

Table 2 shows the amino acid score for the *Balanites aegyptiaca* kernel calculated from the WHO ideal protein value in both children and adults.

**Table 2: Amino acid score for *Balanites aegyptiaca* kernel**


Amino acid	Amino acid (g/100g protein)	WHO ideal [(%Aminoacid/ideal)×100]			
		Children	Adult	Children	Adult
Isoleucine	3.29	2.8	1.3	117.50	253.08
Leucine	6.69	8.3	6.6	80.60	101.36
Lysine	3.52	4.2	5.8	83.81	60.69
Histidine	2.19	1.9	1.6	115.26	136.88
Valine	4.02	4.4	3.5	91.36	114.86
Threonine	2.98	3.0	3.4	99.33	87.65
Total Sulphur amino acid	2.86	1.6	2.5	178.75	114.40
Total Aromatic amino acid	7.21	7.4	6.3	97.43	114.44

Table 3 showed the calculated total amino acids (TAA), total non-essential amino acids (TNEAA), total essential amino acids (TEAA), their respective percentages, the predicted protein efficiency ratio (P-PER), Leu/Ile ratio value for *Balanites aegyptiaca* kernel.

**Table 3: Various Amino acids parameters of *Balanites aegyptiaca* kernel**

Amino acid	Abbreviation	g/100g Protein
Total amino acid	TAA	76.24
Total non-essential amino acid	TNEAA	50.17
Total essential amino acid	TEAA	26.07
-With His		28.26
-No His		26.07
%TNEAA		65.80
%TEAA		34.19
-With His		37.07
-No His		34.19
Total neutral amino acid	TNAA	41.87
%TNAA		54.92
Total acidic amino acid	TAAA	22.28
%TAAA		29.22
Total basic amino acid	TBAA	12.09
%TBAA		15.86
Total sulphur amino acid	TSAA	2.86
%TSAA		3.75
% Cys in TSAA		53.15
Total aromatic amino acid	TArAA	7.21
% TArAA		9.56
Predicted protein efficiency ratio	P-PER	2.26
Leu/Ile ratio		2.03
Leu-Ile (difference)		3.40
% Leu-Ile		4.46

Figure 1 presented the mineral content of *Balanites aegyptiaca* kernel. The result showed a very high quantity of Sodium and Magnesium. The micronutrients (Cu, Cd, Cr, Mn, Ni, Pb, Zn) were quite low except for Fe which is a bit high.



**Figure 1: Mineral contents of *Balanites aegyptiaca* kernel**

#### 4.0 DISSCUSSION

The amino acids profile of *Balanites aegyptiaca* revealed the presence of only seventeen amino acids out of the common twenty amino acids found in proteins. This may be due to the complete destruction of tryptophan during acid hydrolysis and the conversion of the amide glutamine and asparagine to their corresponding amino acids [14]. i.e glutamate and aspartate respectively [15]. The result showed that the percentage of non-essential amino acid is higher than that of essential amino acids which were 50.17% and 26.07% respectively, this may be due to the fact that proteins from plant sources tend to have a relatively lower concentration of protein by mass in comparison to protein from animal sources. They are nevertheless "complete" in that they contain trace amounts of all of the amino acids that are essential in human nutrition in adequate quantities [16]. The non-essential amino acids which are higher than essential amino acids in *Balanites aegyptiaca*

kernel play several important roles in human body along with these essential amino acids. They are incorporated into new proteins needed by cells and can undergo chemical conversion to ultimately create glucose, for use as a fuel source, or fatty acids, for storage of excess calories. Therefore, *Balanites aegyptiaca* kernel is a good source of both non-essential and essential amino acids used as building blocks of protein that boost tissue growth and repair, immunity, red blood cell formation, hormone synthesis among others.

To evaluate the nutritional quality of the kernel, the respective percentages of the essential, total aromatic (TArAA) and total sulphur (TSAA) containing amino acids in the sample were compared with the reference standard amino acids profile established for both adults and preschool children by [17]. the results showed that the TArAA and TSAA of the kernel were all slightly higher than that of the reference standard amino acids profile established for both adults and preschool children by [18] indicating its high nutritional quality. Sulphur containing amino acids, aromatic amino acids and arginine are always required by infants and growing children in order to enhance brain function, growth and boost their immunity against infections. The amino acid score for the kernel was calculated from the WHO ideal protein value in both children and adults. From the calculations done, TSAA and TArAA were found to be 114.40 and 114.44 in adults and 178.75 and 97.43 in children respectively showing that it is a good source of TSAA and TArAA since the TSAA and TArAA are above 100% except for TArAA in children which is slightly below the ideal value. Hence, with proper processing the kernel could meet up with the WHO ideal protein value for both children and adults.

Threonine, Leucine, Valine were found to be close to the WHO ideal protein value in both adults and children of which Isoleucine and lysine with calculated value of 3.29 and

3.52g/100g of protein were only closer to the value of that of children. However, Histidine with calculated value of 2.19g/100g of protein was found to be higher than the WHO ideal protein value in both adults and children. It could therefore be deduced that *Balanites aegyptiaca* kernel could be used in weaning and general food formula for infants and children after adequate processing. In addition, adults especially the elderly could be given the food products of *Balanites aegyptiaca* kernel.

From the various parameters presented, total amino acids (TAA) in the kernels is 76.24g/100g as compared with the TAA in its leaves (71.67g/100g of protein) [19]. The total non-essential amino acids (TNEAA) for the kernel, 50.17g/100g was found to be higher than that of its leaves, 44.95g/100g of protein and their respective percentages are 65.80% and 62.71%. However, the TEAA in *Balanites aegyptiaca* kernel which was 26.07g/100g of protein and its percentage TEAA with His 37.07% was found to be lower than that of soybean (44.4g/100g protein) [20], melon (53.4g/100g protein) and gourd oilseeds (53.6g/100g protein) [21]. Therefore, as expected that for a food substance to be acceptable for consumption its non-essential amino acids has to be higher than that of the essential amino acids as some of these essential amino acids could be harmful when in excess, *Balanites aegyptiaca* kernel could be used as a good source of essential amino acids.

In addition, the predicted protein efficiency ratio (P-PER) for the kernel was 2.27 which is comparable in value with reported values for true digestible protein of whole dried honey bees (*Apis mellifera* L.), 2.47 and 2.50 for casein [22]. Due to the fact that the experimentally determined P-PER usually ranged from 0.0 for a very poor protein to a



maximum possible value of just over 4 [23], the protein from the kernel could be efficiently utilized in human body.

Furthermore, the Leu/Ile ratio value of the kernel was found to be 2.03 of which Ile was less than half that of Leu. It has been suggested that an amino acid imbalance from excess leucine might be a factor in the development of pellagra due to sorghum consumption [24]. High Leu in the diet impairs tryptophan and niacin metabolism and is responsible for niacin deficiency in sorghum eaters [25]. This leads to the hypothesis that excess Leu in sorghum is etiologically related to pellagra in sorghum-eating populations [26]. In addition, a study was carried out showing that Leu/Ile balance is more important than the dietary excess of Leu alone in regulating the metabolism of Tryptophan and niacin and hence the disease process [27]. However, in dogs, experiments have shown that animals fed with sorghum proteins having Leu value to be less than 11.0g/100g protein did not suffer from nicotinic acid deficiency. From some of the above findings, *Balanites aegyptiaca* kernel could therefore be recommended as a good source of protein in supplementing other sources in order to reduce food insecurity and hence malnutrition from lack of protein diets [28].

Figure 1 shows some of the mineral contents of *Balanites aegyptiaca* kernel with Magnesium and Sodium having higher values than other mineral contents. Sodium content was also found to be higher than those in the flower and the recommended daily allowance for Sodium in adults [29]. Otori and Mann [30] shows that *Balanites aegyptiaca* kernel is a good source of Sodium. Magnesium which is important in connection with circulatory diseases and calcium metabolism in bone [31]. Its content in the kernel is higher than that of the [32]. However, Calcium contents of *Balanites*

*aegyptiaca* kernel was found to be lower than those in the flower as reported by [33]. The kernel is therefore not a good source of calcium and could be made up to the recommended daily allowance of calcium though, enough to increase the shelf-life of the kernel. In addition, most of the micronutrients (trace elements) Zinc, Copper, Lead, Cadmium, Chromium, Manganese and Nickel were found to be of low quantities. This indicates that the kernel is a good source of essential nutrients in moderate quantities needed by animals for various activities especially in metabolic processes acting as cofactors.

## 5.0 CONCLUSION

The study revealed that *Balanites aegyptiaca* kernel is of high nutritional value due to its high contents of minerals, essential and non-essential amino acids.

## Reference:

- [1]. Clement, A.O., Jacob, G.A., James, M.K., Refaat, A. and Ahmed, A. (2011). Harvesting and Processing of *Balanite aegyptiaca* Leaves and Fruits for Local Consumption by Rural Communities in Uganda. *Journal of Food Technology* 9 (2):83-90.
- [2]. Chothani and Vaghasiya, A.O., Jacob, G.A., James, M.K., Refaat, A. and Ahmed, A. (2011). Harvesting and Processing of *Balanite aegyptiaca* Leaves and Fruits for Local Consumption by Rural Communities in Uganda. *Journal of Food Technology* 9 (2):83-90.
- [4]. Daya, L. C. and H. U. Vaghasiya (2011). A review on *Balanite aegyptiaca* Del (Desert date): phytochemical constituent, traditional uses and pharmacological activity. *Pharmacogen Review* 5 (9):55-62.
- [5]. Yadav, J.P. and Manju, P. (2010). *Balanites aegyptiaca* (L.) Del. (Hingot): A review of its traditional uses, phytochemistry and pharmacological properties. *International Journal of Green Pharmacolgy* 4:140-6.
- [6]. Gnoula, C., Guisso, P., Duez, P., Frederich, M. and Dubois J. (2007) Nematocidal compounds from seeds of *Balanitesa egyptiaca* Isolation and structure Elucidation. *International Journal of Pharmacology* 3:280-4.

- [7]. Lohlum, S.A., Forcados, E.G., Agida, O.G., Ozele, N. and Gotep J.G. (2012) Enhancing the Chemical Composition of *Balanites aegyptiaca* Seeds through Ethanol Extraction for Use as a Protein Source in Feed Formulation. Biochemistry Division, National Veterinary Research Institute Vom, Jos. Nigeria. URL:<http://dx.doi.org/10.5539/sar.v1n2p251>. Retrieved, July 28, 2017.
- [8]. Chothani and Vaghasiya, A.O., Jacob, G.A., James, M.K., Refaat, A. and Ahmed, A. (2011). Harvesting and Processing of *Balanite aegyptiaca* Leaves and Fruits for Local Consumption by Rural Communities in Uganda. *Journal of Food Technology* 9 (2):83-90.
- [9]. Liener, I.E. and Kkade, M.L. (1980). Protease inhibitor, Toxic constituents of plants foodstuffs. Academic Press. New York. Pp. 7-71.
- [10]. Larsgon, M., Rossande- Hulthen, L., Sandstome, B. and Sandberg, A. (1996). Impoved iron and zinc absorption from breakfast meals containing malted oats with reduced phytate content. *Journal of Nutrition* 76:677-688.
- [11]. Benitez, L. V. (1989). Amino Acid and fatty acid profiles in aquaculture nutrition studies, p. 23- 35.in S.S. De Silva (ed.) Fish Nutrition Research in Asia. Proceedings of the Third Asian Fish Nutrition Network Meeting. *Asian fish Society Special Publication*. 4: 166.
- [12]. AOAC (2006). Official Method of Analysis of the Association of Official Analytical Chemists, Washington, D. C.. 18<sup>th</sup> Edition, Pp 810-931.
- [13]. Wathelet, B. (1999) athelet, B., 1999. Nutritional analysis of proteins and amino acids in beans (*Phaseolus* spp.).*Biotech. Agronomy Soc. Environ.* 3: 197-200.
- [14]. Alhassan, A.J. and Sule, M.S. (2011). Characterization and nutritional quality of degutted white grubs' protein. *Biological and Environmental Sciences Journal for the Tropics*. 8(3): 26-30.
- [15]. McDougall J. (2002). Plant foods have a complete amino acid composition. *Circulation* 105(25):197.
- [16]. WHO/FAO/UNU (1985). Report: Energy and protein requirements: WHO technical report series No.724, pp. 220.
- [17]. WHO/FAO/UNU (1985). Report: Energy and protein requirements: WHO technical report series No.724, pp. 220.
- [18]. Kubmarawa, D., Andenyang, I.F.H. and Magomya, A.M. (2008). Amino acid profile of two non-convetional leafy vegetables, *Sesamum indicum* and *Balanites aegyptiaca*. *African Journal of Biotechnology* 7 (19): 3502–3504.

240  
 241 [19]. Kuri, Y.E., Sundar, R.K., Kahuwi, C., Jones, G.P. and Rivett, D.E. (1991).  
 242 Chemical composition of *Momordica charantia* L. fruits. *Journal of Agricultural and*  
 243 *Food chemistry*. 39: 1702-1703.  
 244

245 [20]. Olaofe, O., Adeyemi, F.O. and Adediran, G.O. (1994). Amino acid and mineral  
 246 compositions and functional properties of some oil seeds. *Journal of Agricultural and*  
 247 *Food Chemistry*. 42: 878-881.

248 [21]. Ozimek, L., Saver, W.C., Kzokwoski, V., Ryan, J.K., Jorgensen, H. and Jelen, P.  
 249 (1985). Nutritive value of protein extracted from honey bees. *Journal of Food Sciences*.  
 250 50: 1332.  
 251

252 [22]. Muller, H.E. and Tobin, G. (1980). *Nutrition and Food Processing*. London: Crown  
 253 Helm. Pp. 450.  
 254

255 [23] FAO (1995). Sorghum and millets in human nutrition. Rome, Italy: FAO Food  
 256 Nutrition Series, No. 27: 23-29.  
 257

258 [24]. Guinand, Y. and Dechassa, L. (2009) Wild-Food Plants in Southern Ethiopia:  
 259 Reflections on the role of 'famine-foods' at a time of drought" UN-OCHA Report.  
 260

261 [25]. Belavady, B., Srikantia, S.G. and Gopalan, C. (1963). The effect of oral  
 262 administration of leucine on metabolism of tryptophan. *Biochemical Journal* 87: 652-  
 263 655.

264 [26]. FAO (1995). Sorghum and millets in human nutrition. Rome, Italy: FAO Food  
 265 Nutrition Series, No. 27: 23-29.  
 266

267 [27]. Belavady, B. and Udayasekhara-Rao, P. (1979). Leucine and Isoleucine content of  
 268 jowar and its pellagragenicity. *Indian Journal of Experimental Biology*. 17: 659-661.

269 [28]. Belavady, B. and Udayasekhara-Rao, P. (1979). Leucine and Isoleucine content of  
 270 jowar and its pellagragenicity. *Indian Journal of Experimental Biology*. 17: 659-661.

271 [29]. Umar, K.J., Abubakar, L., Alhassan, B., Yahaya, S.D., Hassan, L.G, Sani, N.A.,  
 272 Muhammad, M.U (2014). Nutritional Profile of *Balanites aegyptiaca* Flower. *Studia*  
 273 *Universitatis "Vasile Goldiş", Seria Ştiinţele Vieţii* (24) Pp 169-173.  
 274

275 [30]. Umar, K.J., Abubakar, L., Alhassan, B., Yahaya, S.D., Hassan, L.G, Sani, N.A.,  
 276 Muhammad, M.U (2014). Nutritional Profile of *Balanites aegyptiaca* Flower. *Studia*  
 277 *Universitatis "Vasile Goldiş", Seria Ştiinţele Vieţii* (24) Pp 169-173.  
 278

- [31]. Otori, A.A. and Mann, A. (2014).Determination of chemical composition, minerals and antinutritional factors of two wild seeds from Nupeland, North Central Nigeria. *AmericanJournal of Chemistry and Application* 1(1): 20-26.
- [32] FAO (1995). Sorghum and millets in human nutrition. Rome, Italy: FAO Food Nutrition Series, No. 27: 23-29.
- [33]. Umar, K.J., Abubakar, L., Alhassan, B., Yahaya, S.D., Hassan, L.G, Sani, N.A., Muhammad, M.U (2014).Nutritional Profile of Balanites aegyptiaca Flower.Studia Universitatis “Vasile Goldiș”, Seria Științele Vieții (24) Pp 169-173.