

## Original Research Article

# PHYSICOCHEMICAL PROPERTIES OF *ALCHORNEA CORDIFORLIA*, *CYPERUS ESCULENTUM* AND *IRVINGIA GABONENSIS* SEED OILS AND THEIR APPLICATIONS

## ABSTRACT

The physicochemical properties of oils extracted from three locally available plant seeds in Nigeria namely: *Alchomea cordiforlia*, *Cyperus esculentum* and *Irvingia gabonensis* using n-Hexane were determined. The results of the analysis revealed that their % yield were 37.00, 27.50 and 33.00 for *A. cordiforlia*, *C. esculentum* and *I. gabonensis* respectively. Their odour was non-offensive and their colour bright and attractive. The specific gravity of the oils was 0.91, 0.94 and 0.92 for *A. cordiforlia*, *C. esculentum* and *I. gabonensis* respectively. Their flash points in °C were also 155, 159 and 229 respectively, indicating that *I. gabonensis* is the most thermally stable oil and suitable for frying. The chemical properties for *A. cordiforlia*, *C. esculentum*, and *I. gabonensis* respectively were as follows: Acid values in mEqKg<sup>-1</sup> were 24.67, 5.33 and 3.73. Peroxide values in mKqKg<sup>-1</sup> were 7.26, 9.86 and 2.96. Saponification values were 162.13, 179.52 and 238.43. Iodine values were 24.62, 11.68 and 3.38. These results indicates that the three seeds are viable sources of oil based on their % yield. They are good for both domestic and industrial use based on their acid, saponification and iodine values. Their properties in most cases compete favorably with palm kernel oil (PKO) which is currently being used for many domestic and industrial purposes in Nigeria especially for the making of paints, soap, cosmetics, lubricant and varnishes.

**Keywords:** Physiochemical Properties, Seed oils, Applications

## INTRODUCTION

Oil from plants is used for both domestic and industrial purposes all over the world based on its physical and chemical properties. Palm kernel oil is one of the most commonly used vegetable oils because its properties

35 have been studied and known especially by researchers like Akubugwo  
36 and Ugborgu, stating that is good for soap making and the production of  
37 cosmetics, paints, varnishes industrially based on its saponification  
38 value, iodine value etc[1]. According to Unilever [2], vegetable oils from  
39 plants like oil palm tree, groundnut, olive, beniseed (*sesame*), soya  
40 beans, coconut, castor seed, linseed etc. plays an important role in our  
41 diet as a source of fat and oil, a major class of food required for  
42 warmth and energy in the body. Besides, some of these oils are used for  
43 the production of commodities like soaps, cosmetics, paints, varnishes,  
44 lubricants plastics, while others are used for cooking or are prepared  
45 and eaten in form of butter or margarine. Moreover, some vegetable oils  
46 are now used as substitutes for petrol or diesel as fuel in automobiles in  
47 the form of biodiesel or bioethanol [3]. Medicinally, Ihesie have reported  
48 that sesame seed oil can be used to treat health problems like chronic  
49 constipation in elders, round worms in children, dysmenorrhea (painful  
50 menstruation) in women, amenorrhea, asthmatic symptoms, coughs and  
51 hiccoughs, and insufficient flow of breast milk in nursing mothers by the  
52 oral intake of the oil up to two tea spoons at a time [4]. Akpe have also  
53 studied the physiochemical properties of *Carica papaya*, *Citrus paradisi*  
54 and *Croton zambesicus* seed oils and reported that they compete  
55 favorably with palm kernel oil used for several industrial purposes in

56 Nigeria [5]. Based on the facts so far, the importance of vegetable oils  
57 to man cannot be over emphasized and their economic value  
58 unquantifiable. However, one can observed that the oil plants or crops  
59 mentioned above are a small percentage of the several hundreds of  
60 plants in nature whose oil potentials have not been discovered, even  
61 some that have been identified as oil seed crops are being underutilized  
62 because their oil properties and potentials have not been properly  
63 studied to ascertain their suitability for use domestically and or  
64 industrially. Consequently, this study is aimed at determining the  
65 physicochemical properties of oils extracted from *Alchornea cordifolia*  
66 (Christmas bush), *Cyperus esculentum* (Tiger nut) and *Irvingia*  
67 *gabonensis* (Bush mango) which are found locally in Obudu Area of  
68 Cross River State and many other parts of Nigeria and have not been  
69 properly studied. This is with a view to determine their potentials and  
70 properties as sources of vegetable oil for domestic and industrial uses,  
71 and also compare their properties with the established properties of  
72 palm kernel oil from *Elaeis guinensis* (palm tree) which is popularly in  
73 use now.

## 74 **MATERIALS AND METHOD**

75 **Sample collection and preparation:** Viable or healthy seeds of  
76 *Alchornea cordifolia* (Christmas bush), *Cyperus esculantum* (Tiger nut)  
77 and *Irvingia gabonensis* (Bush mango) were collected locally from  
78 Obudu Area of Cross River State of Nigeria between February and  
79 March, and were taken to the Department of Botany, University of  
80 Calabar for identification of botanical names and labelling. The samples  
81 were then taken to the Chemistry Department of the same University  
82 where they were shelled or dehauled (where applicable), sun dried for  
83 several days, wrapped in polythene bags and kept for use within one  
84 month. Each of sample was crushed or ground into a paste using a  
85 manual grinding machine. 100 g of the paste of each sample was  
86 packed in an ashless filter paper and placed in the thimble of a Soxhlet  
87 apparatus (extractor) and extracted using N-hexane as the extracting  
88 solvent. At the end of the continuous extraction for about 5 to 6 hours,  
89 the extracting solvent was evaporated off leaving the oil sample for  
90 analysis. The percentage yield of the oil extract of each sample was  
91 determined thus:

$$\% \text{ yield} = \frac{\text{weight of oil}}{\text{weight of sample}} \times 100\%$$

92 **Sample analysis:** The specific gravity of the oils was determined  
93 according to the method reported by Onwuka [6] thus: A 50 mL

pycometer bottle was washed with water and detergent, rinsed and dried. The bottle was filled with distilled water and weighed. After drying the bottle of water, it was filled with the oil sample and weighed. The specific gravity was calculated thus:

$$\text{Specific gravity} = \frac{\text{weight of 50mL of oil}}{\text{weight of 50mL of water}}$$

The colour, state at room temperature and the odour were observed and perceived using the human sense organs. The flash point of the oil samples was determined following the procedure reported by [6] thus: 10 mL of the oil was poured into an evaporating dish and placed on a source of heat. A thermometer was suspended at the centre of the dish ensuring that its bulb dips inside the oil without touching the bottom of the dish. The temperature of the oil was raised gradually by regulating the source of heat. The point at which the oil began to give off a thin bluish smoke continuously (i.e. smoke point), a flame was applied using a match-stick. The temperature at which the oil started flashing when the flame is applied without supporting combustion was noted as the flash point of the oil.

The acid value was determined following the method of AOAC [7] as reported by [6] thus: 1.0 g each of the oils was dissolved in a mixture

obtained by mixing 25 mL diethylether and 25 mL ethanol, and titrated with 0.1M NaOH using phenolphthalein as an indicator, shaking till a pink colour end point which persisted for 15 seconds was observed. The acid value and % free fatty acids were calculated as follows:

$$\text{Acid Value} = \frac{\text{Titre volume (mL)} \times 56.1 \times M}{\text{weight of sample}}$$

Where M is the molarity of NaOH (0.1M).

Acid value is expressed in milli equivalent per kilogramme ( $\text{mEqkg}^{-1}$ ).

$$\% \text{ free fatty acid} = \frac{1}{2} \times \text{Acid value}$$

The saponification value was determined using the method of [7] as described by [6] thus: 1 gram of the oil was weighed into a round bottom flask and 24 mL of alcoholic potassium hydroxide solution was added. A reflux condenser was attached to the flask and heated on a sandbath for 1 hour shaking frequently. One mL of phenolphthalein (1%) solution was added and titrated while hot with 0.5M HCl to a colourless end point. A blank titration was also carried out the volume at end point recorded. The saponification value was calculated thus.

$$\text{Saponification value} = \frac{(X - Y) \times 56.1 M}{\text{Weight of sample}}$$

Where X = volume (mL) of test solution titration

128 Y = volume (mL) of blank titration

129 M = Molarity of HCl (0.5)

130 The peroxide value was determined using the method of [7] as described by  
 131 [6] thus: 1 ml of potassium iodide (KI) was added to 20 mL of a solution of  
 132 mL of (2:1) volumes of glacial acetic acid and chloroform. The result out  
 133 solution was added to 1.0 g of the oil sample in a clean dry conical flask. The  
 134 mixture was left in a dark for about 2 minutes and 30 mL of distilled water  
 135 was added and titrated with 0.02M sodium thiosulphate solution using 5 mL  
 136 starch as indicator. A blank titration was also carried out. The peroxide value  
 137 was calculated thus:

$$\text{Peroxide value} = (100M (Va - Vb))/W$$

138 Where W = weight of oil sample

139 Va = volume in Ml of thiosulphate used in test solution

140 Vb = volume in mL of thiosulphate used in blank solution

141 M = molarity of sodium thiosulphate (0.02).

142 The iodine value was determined using Wij's method as described by [6]  
 143 thus: 0.5 g of the oil samples were poured into a beaker and 10ml of carbon  
 144 tetrachloride was added, 20 mL of Wij's solution was added and a stopper  
 145 previously moisten with potassium iodide was inserted and allowed to stand  
 146 in the dark for 30 minutes. 15mL of potassium iodide solution (10%) was

added and titrated with 0.1M thiosulphate solution using starch as indicator.

A blank titration was also carried out. The iodine value was calculated thus:

$$Iodine\ value = \frac{(b - a) \times 12.69M}{weight\ of\ sample}$$

Where a = volume in mL of test titration

b = volume in mL of blank titration

M = molarity of thiosulphate (0.1)

**RESULTS:** The results of the physical and chemical properties are presented in Table 1 and 2 respectively

**Table 1: Physical properties of *Alchornea cordifolia*, *Cyperus esculentum* and *Invingia gabonensis* seed oils.**

	% yield	Specific gravity	Flash point (°C)	State at 25°C (Room temperature)	Colour	Odour
<i>Alchornea cordifolia</i>	37.00±1.50	0.91±0.01	155.00± 2.00	Semi-solid	Reddish	Non-offensive
<i>Cyperus esculentum</i>	27.50± 2.01	0.94± 0.02	159.00± 2.50	Liquid	Light yellow	Non-offensive
<i>Invingia gabonensis</i>	33.00± 2.00	0.92± 0.02	229.00± 2.00	Waxy solid	Milky white	Non-offensive

Values reported in mean ± SD, with N = 3



**Table 2: Chemical properties of *Alchornea cordifolia*, *Cyperus esculentum* and *Invingia gabonensis* seed oils**

Seed oil	Acid value in mEqkg <sup>-1</sup>	% free fatty acids	Peroxide value	Saponification value	Iodine value
<i>Alchornea cordifolia</i>	24.67±1.25	12.34±1.30	7.26± 0.12	162.13± 2.50	24.62± 1.50
<i>Cyperus esculentum</i>	5.33± 0.15	2.67± 0.10	9.86± 0.15	179.52± 3.00	11.68± 1.25
<i>Invingia gabonensis</i>	3.73± 0.13	1.87± 0.11	2.96± 0.02	238.43± 2.50	3.38± 0.15

Values reported in mean ± SD, with N = 3

## DISCUSSION

The percentage yield in Table 1 revealed that *A. cordifolia* (37.00%), *C. esculentum* (27.50%) and *I. gabonensis* (33.00 %). This shows that *A. cordifolia* has the highest yield, followed by *I. gabonensis* while *C. esculentum* has the lowest % yield. However, these values compete favorably with palm kernel oil with a % yield of 28% as reported by [1], with *A. cordifolia* and *I. gabonensis* even better as their values are higher than that of palm kernel oil that is commonly used. Thus, the 3 seed plants can be used as good sources of vegetable oil.

The flash point is the temperature at which volatile evolving from the heated oil will flash but not support combustion. It measures the thermal stability of the oil [6]. It is also an indicator for the suitability of the oil for frying [5]. The results in Table 1 shows that the flash points in

181 °c were 155, 159 and 229 for *A. cordifolia*, *C. esculentum* and *I.*  
182 *gabonensis* respectively. This results indicate that *I. gabonensis* oil is a  
183 better frying oil and a more thermally stable oil than *C. esculentum* and  
184 *A. cordifolia* respectively.

185 The specific gravity (relative density) of the oils was 0.91, 0.94  
186 and 0.92 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively,  
187 all higher than 0.88 reported for palm kernel oil by [1], which is  
188 commonly used industrially. All the oil samples were non-offensive in  
189 their odour, *A. cordifolia* was reddish in colour, *C. esculentum* was light  
190 yellow and *I. gabonensis* milky white in colour. This makes the oil  
191 attractive and appealing. Their state at room temperature was semi-  
192 solid, liquid and waxy solid for *A. cordifolia*, *C. esculentum* and *I.*  
193 *gabonensis* respectively. This also compete with palm kernel oil (PKO)  
194 which is semi-solid as reported by [1].

195 The chemical properties of the studied oils are reported in Table 2.  
196 The results showed that the acid values of the oils were 24.67, 5.33 and  
197 3.73 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively.  
198 Also, the % free fatty acids were 12.34, 2.67 and 1.87 for *A. cordifolia*,  
199 *C. esculentum* and *I. gabonensis* respectively. All these values are less  
200 than PKO with an acid value of 14.04 [1]. Acid value is an indicator for

edibility of oil and suitability for use in the paint industry. *C. esculentum* and *I. gabonensis* oil are edible going by their free fatty acid value of less than 3 [8] as cited by [5]. They can also compete favorably with sesame, soya bean, sun flower and rapeseed oils with acid value of about 4 as reported by [9]. Thus, the two oils with the least acid value can be consume directly (i.e. *C. esculentum* and *I. gabonensis*).

The peroxide values of the oils were 7.26, 9.86 and 2.96 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. It is an indicator for the deterioration of oils. Fresh oils have values less than 10 mEqkg<sup>-1</sup> and rancid oils have values ranging from 20 to 40 [6]. It is also an indicator for longer and shorter shelf life during storage, as fresh oils last longer [10]. Thus, all the 3 seed oils are fresh oils and compete favorably with 2.12 mEqKg<sup>-1</sup> reported for PKO by [1].

The saponification value is an indication that the oils have potential for use in the industry when values are high especially for soap and cosmetics [11]. Its values for the oils were 162.13, 179.52 and 238.43 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. *I. gabonensis* with the highest value has the best potential for industrial use and compete with PKO that has a value of 246. 60 [1].

The Iodine values of the oils were 24.62, 11.68 and 3.38 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. These values indicates that all the three oils are non-drying oils because their values are less than 100, those with values between 100 and 150 are semi-drying oils while those greater than 150 are drying oils [12]. This non-drying character qualifies them for use in the paint industry [13]. However, the oils compete favorably with PKO which is also non-drying oil with an iodine value of 18.30 as stated by [1]. Based on their iodine values, the storage procedure should ensure protection from oxidative rancidity or deterioration as they contain appreciable level of unsaturated bonds.

## CONCLUSION

At the end of this study, the 3 oil seeds can be classified as high yielding based on the % yield. *C. esculentum* and *I. gabonensis* are suitable for direct consumption by their free fatty acid value. Their iodine and saponification values shows they are suitable for the industrial production of soaps, cosmetics, paints etc. Their colours are bright and attractive while their odours are non-offensive. Most of the physicochemical properties of the three seed oils studied compete favorably with palm kernel oil (PKO) and conventional seed oils like

groundnut oil, soya bean, rapeseed, castor seed etc. One can therefore recommend that the 3 seed oils have potentials for development and use for domestic and industrial purposes.

## REFERENCES

- Akubuguo, I. E and Ugbogu, A. E. Physiochemical studies on oils from five selected Nigerian plant seeds. *Pakistan Journal of Nutrition* (2007) 6(1): 75-78.
- Unilever Limited. Vegetable oil and fats. A Unilever educational booklet; Revised ordinary series, (1968) 2:1-5
- Ekpa, O. D. Entrepreneurial Chemistry: A Chemist's approach to wealth creation in a depressed Economy. 43<sup>rd</sup> Inaugural lecture University of Calabar, Calabar, (2008).
- Ihesie, G. Medicinal Uses of Sesame oil. *Saturday Vanguard*, November, (2008) 29, 44-45.
- Akpe, M. A., Oyo-Ita, I. and Dosunmu, M. I. Analysis of the physiochemical properties of Carica Papaya Citrus paradisi and croton zambesicus seed oils and their applications. *World Journal of Research and Review* (2018) 6(2): 6-8.
- Onwuka, G.A. Food Analysis and instrumentation; Theory and practice. Napthali Prints, Lagos, Nigeria, (2005).
- AOAC (1984). Official Methods of Analysis: 4<sup>th</sup> ed. Association of Official analytical chemists; V.A. Arlington.
- Eckey, E.W. Vegetable fats and oils. Reinhold; New York, 1954.
- Pearson, D. The Chemical Analysis of food. Churchill, Livingstone, (1976).
- Eka, O. U. Proximate Composition of Bush Mango tree and some properties of Dika fat. *Nig. J. Nutri. Sci.* (1980); 1:33-36
- Amoo, I. a Eleyinmi, A.F., Oilelaboye, N. A and Akoja, S. S. Characteristics of Oil extract from gourd (Cucurbita Maxima) Seed Food, *Agriculture and Environment*; (2004), 2:38-39

- 269 Schwartz, T. A., Bunce, M. D., Silberman, R. G., Stanizki, C. I., Starlton, W. J.  
270 and Zipp, A. P. *Chemistry in context*. 2<sup>nd</sup> Edition: U.S.A, McGraw-  
271 Hill companies Inc, (1997).
- 272 Dosunmu, M.J. and Ochu, C. Physiochemical properties and fatty acid  
273 composition of lipids extracted from some Nigeria fruits and seeds.  
274 Global J. Pure and applied Sci. (1995), 1:45-47.