

NUTRITIONAL COMPOSITIONS OF SELECTED GREEN LEAFY VEGETABLES IN OYO STATE, NIGERIA

ABSTRACT

The nutritional compositions of selected green leafy vegetables obtained from major towns in Oyo State, Nigeria, (March and April) were determined using standard analytical methods for proximate analysis. The following nutrients in percentage were determined; moisture contents, ash, fat, crude fibre and crude protein. The green leafy vegetable used are *Talium triangulare*, *Amaranthus hybridus*, *Launaena taraxacifolia* *Ocimum gratissimum*, *Celosia argentea*, *Cucuribita maxima*, *Abelimoschus esculentus*, *Solanum macrocarpon*, *Vernonia amygdalina* and *Sesamum indicum*. All nutrients were present in appreciable quantities. Moisture contents ranged from $5.33 \pm 0.06\%$ - $8.33 \pm 0.06\%$, ash ($23.07 \pm 0.06\%$ - $61.27 \pm 0.06\%$), fat ($1.13 \pm 0.06\%$ - $3.37 \pm 0.06\%$), crude fibre ($2.43 \pm 0.12\%$ - $22.03 \pm 0.06\%$), and crude protein ($18.50 \pm 0.10\%$ - $55.23 \pm 0.06\%$). The functional properties of vegetables were close in term of high protein level indicating that they are more nutritious. Also, the level of their ash content showed that the vegetables are very rich in essential minerals for healthy life when compared with one another and recommended dietary allowance (RDA). Thus, there is a need for farmers in the area to engage in dry season vegetable production so as to ensure availability of leafy vegetables throughout the year.

Keywords: Oyo state, proximate, nutritional composition, green leafy vegetables.

INTRODUCTION

Vegetables are the fresh and edible portions of herbaceous plants, which can be eaten raw, or cooked [1, 2]. They contain valuable food ingredients which can be successfully utilized to

27 build up and repair the body. Vegetables are valuable in maintaining alkaline reserve of the
28 body. Vegetables can be grouped into edible roots, stems, leaves, fruits or seed. Each group
29 contributes to diet in its own way [2]. Leafy vegetables are regular ingredient in the diet of
30 average Nigerian with their level of consumption; they can provide appreciable amounts of
31 nutritive minerals [3]. *Amarathus hybridus*, *Celusia argentea*, *Abelmoschus esculentus*,
32 *Talinum triangulare*, *Vernonia amygdalina* and *corchorus olitorious* are popular edible
33 vegetables in Nigeria. *Corchorus olitorius* is usually recommended for pregnant women and
34 nursing mothers because it is believed to be rich in iron [4].

35 Most developing countries depend on starch-based food as the main staple food for the
36 supply of both energy and protein. This account in part for protein deficiency which prevails
37 among the populace as recognized by Food and Agricultural Organization [5].

38 Apart from the variety which they add to the menu [6, 7], they are valuable sources of
39 nutrients especially in rural areas where they contribute substantially to protein, minerals,
40 vitamins, fibers and other nutrients which are usually in short supply in daily diets [8]. It is
41 worthwhile to note that consumption of numerous types of edible plants as sources of food
42 could be beneficial to nutritionally marginal population especially in developing countries
43 where poverty and climate is causing havoc to the rural populace [9]. African indigenous
44 leafy vegetables (ALVs) are the cheapest and most readily available sources of important
45 proteins, vitamins, especially the pro-vitamin A [9] and essential amino acids. Vegetables
46 rank higher in production than all other crops; they are known to provide 80% of the vitamin
47 A in diet [10]. Indigenous vegetables are reported to play a very important role in income
48 generation and subsistency [11]. They are important commodities for poor households
49 because their prices are relatively affordable when compared to other food items. Vegetables
50 provide very important sources of employment for those outside the formal sector in urban
51 areas because of their generally short, labour intensive production systems, low levels of

investment and high yield [11]. A large number of African indigenous leafy vegetables have long been known and reported to have health protecting properties and uses [12]. It is reported that the roots, leaves and twigs, as well as the bark of the tree of Moringa plants (*Moringa oleifera*) are used in traditional medicine [13].

The WHO recommended a minimum daily intake of 400g of fruits and vegetables [14]. However, it is not clear from the report what proportion of this total daily intake should go to vegetables. Nevertheless, according to the Kobe framework document and an FAO report, the recommended total daily intake is equivalent to five (5) servings of 80g each of fruits and vegetables [15, 16]. Vegetables are full of water, especially when eaten raw, and when eaten, the body does not need to use some of its own water to digest them. This means less pressure is put on the digestive systems [17]. Green leafy vegetables like cabbage, lettuce, dandelion, and Moringa may be eaten raw, boiled or dried. Perhaps the most common use in all parts of the world is boiled vegetable leaves. This process eliminates potential pathogens, sometimes poisonous or irritating substances are neutralized and spoilage is brought to a halt [18].

In Nigeria, as in most other tropical countries of Africa where the daily diet is dominated by starchy staple foods, vegetables are the cheapest and most readily available sources of important proteins, vitamins minerals and essential amino acids [2]. Vegetables also act as buffering agents for acidic substances produced during the digestion process [2].

Traditional African leafy vegetables are better adapted to the environment than the introduced exotic vegetables and also provide low-cost quality nutrition for large parts of the population in both rural and urban areas [19]. Inadequate information on these vegetable species is causing gradual neglect of some of the useful ones that have been used for food over the years. Vegetables are a vital constituent of West African diet, and traditional vegetable species are highly important yet, many species are poorly known, being used only locally [20]. The objective of this study is to evaluate the nutritional value of some leafy vegetables

77 consumed in Oyo State South West of Nigeria to give more information that are lacking on
78 the importance of these vegetables on the nutrient they supply.

79 **MATERIALS AND METHODS**

80 Eleven leafy vegetables were collected from different locations within the study area (Ibadan,
81 Ogbomoso, Oyo, Igboora, Iseyin, and Saki all in Oyo State) between months of March and
82 April 2016. Vegetable collected includes *Talium triangulare*, *Amaranthus hybridus*,
83 *Launaena taraxacifolia* *Ocimum gratissimum*, *Celosia argentea*, *Cucuribita maxima*,
84 *Abelimoschus esculentus*, *Solanum macrocarpon*, *Vernonia amygdalina* and *Sesamum*
85 *indicum* (English/Common and Local name shown in Table 1). They were identified and
86 authenticated at department of Botany, University of Ibadan, Nigeria. The vegetables were
87 air-dried at room temperature (25°C) and ground to fine powder, using a laboratory mill and
88 stored in air-tight containers for laboratory analysis. The nutritional compositions in terms of
89 proximate analysis were carried out to determine Moisture contents, crude protein, crude
90 fibre, fat, and ash. All analysis was carried out in triplicates.

91

92

93 **NUTRITIONAL ANALYSIS**

94 The moisture content of the vegetable samples was determined after drying at 105 °C in an
95 electric oven (model: UNISCOPE5M9053) until a constant weight was attained [21]. The
96 micro-Kjeldahl method was employed to determine the total nitrogen and the crude protein
97 (N×5.95) [21]. A dry ashing method was used to determine the ash content [21]. The samples
98 were ashed in a furnace (model: SXL) at 550 °C. Crude fat was determined by Soxhlet
99 extraction and crude fiber by incineration method after acid and base digestion.

STATISTICAL ANALYSIS

Three determinations were carried out for each analysis. The mean value and standard deviation were separated using Duncan's Multiple Range Test (DMRT) at $p \geq 0.05$.

RESULTS AND DISCUSSION

Table: 1. Botanical, English/Common and Local name of the vegetable samples

Botanical name	English name	Local name
<i>Amaranthus hybridus</i>	Smooth amaranth	Tete
<i>Talinum triangulare</i>	Water leaf	Gure
<i>Launaena taraxacifolia</i>	Wild lettuce	Yanrin
<i>Ocimum gratissimum</i>	African basil	Efinrin
<i>Cucuribita maxima</i>	Winter squash	Gboro
<i>Ocimum canum</i>	African mint	Marugbo
<i>Celosia argentea</i>	Cocks comb	Soko
<i>Solanum macrocarpon</i>	African eggplant	Igbo/Gbagba
<i>Abelmoschus esculentus</i>	Okro leave	Ilasa
<i>Vernonia amygdalina</i>	Bitter leaf	Ewuro
<i>Sesamum indicum</i>	Sesame	Ekuuku/Morogbo

Table: 2. Proximate Composition of green leafy Vegetable Samples

Leafy Vegetables	Moisture (%)	Ash (%)	Fat (%)	Crude Fiber (%)	Crude Protein (%)
<i>Amaranthus hybridus</i>	5.50 ± 0.10^b	27.53 ± 0.06^a	3.37 ± 0.06^a	10.07 ± 0.06^a	55.23 ± 0.06^a
<i>Talinum triangulare</i>	7.70 ± 0.10^b	40.10 ± 0.10^b	3.23 ± 0.06^a	20.07 ± 0.12^b	25.30 ± 0.10^b
<i>Launaena taraxacifolia</i>	5.33 ± 0.06^a	23.07 ± 0.06^a	1.30 ± 0.00^a	20.10 ± 0.10^a	50.10 ± 0.10^b
<i>Ocimum gratissimum</i>	7.73 ± 0.06^a	46.2 ± 0.00^c	1.37 ± 0.06^a	10.10 ± 0.10^b	32.63 ± 0.06^a
<i>Cucuribita maxima</i>	6.17 ± 0.06^a	50.27 ± 0.06^a	1.13 ± 0.06^a	10.07 ± 0.12^a	30.47 ± 0.06^a
<i>Ocimum canum</i>	6.77 ± 0.06^a	23.13 ± 0.06^a	1.20 ± 0.00^a	20.13 ± 0.12^a	47.13 ± 0.12^b

<i>Celosia argentea</i>	6.33 ± 0.06 ^a	40.13 ± 0.06 ^a	2.67 ± 0.06 ^a	22.03 ± 0.06 ^a	20.20 ± 0.17 ^c
<i>Solanum macrocarpon</i>	6.27 ± 0.06 ^a	54.03 ± 0.06 ^a	2.20 ± 0.10 ^b	2.43 ± 0.12 ^b	23.10 ± 0.10 ^b
<i>Abelmoschus esculentus</i>	6.23 ± 0.06 ^a	61.27 ± 0.06 ^a	3.40 ± 0.10 ^b	7.60 ± 0.10 ^b	20.10 ± 0.00 ^c
<i>Vernonia amygdalina</i>	7.27 ± 0.06 ^a	58.33 ± 0.15 ^c	2.47 ± 0.06 ^a	10.17 ± 0.06 ^a	19.20 ± 0.17 ^b
<i>Sesamum indicum</i>	8.33 ± 0.06 ^a	58.17 ± 0.06 ^a	2.33 ± 0.06 ^a	10.13 ± 0.06 ^a	18.50 ± 0.10 ^a

Mean values ± Standard deviation values. Values carrying different letter within a column are significantly different at P>0.05.




Figure: 1. Graphical comparison of vegetables nutritional composition in percentage.

The proximate composition of the eleven green leafy vegetables and Graphical comparison of vegetables nutritional composition in percentage are shown in the table 2 and figure 1 respectively and Botanical, English/Common and Local name in table 1 [22].

The eleven leafy vegetables showed moisture content ranges from 5.33% to 8.33% (Table 2 and **fig.1**). The results obtained were close to that reported for *Basella alba* (11.57%) and *Amaranthus hybridus* (10.00%) by Asaolu *et al.* [23], *Cleome gynandra* (15.90%) reported by Clement [19] and 15.6% for *Celosia argentea* by Onwordi *et al.* [2]. In this study, it was

observed that a closer relationship occurred between the moisture content of the leafy vegetables under study, but a great difference was observed when compared with the one reported by Idris *et al.* [24] for *Abelimoschus esculentus* (82.53%) and 79.98% reported for *Corchorus olitorious* by Adeniyi *et al.* [25]. Also, as it has been reported in the work of Kwenin *et al.* [17] that leafy vegetables have high moisture content ranging from 72.93% to 91.83%, the significant difference observed now may be due to the cultivation conditions that influence the water level of vegetables [26]. George [27] stated that moisture content makes an important contribution on the texture of the leaves and helps in maintaining the protoplasmic content of the cells; it also makes them perishable and susceptible to spoilage by microorganisms.

The highest result was recorded for *A. esculentus* (61.27%) while *Launaena taraxacifolia* had the least value of 23.07%. Ash, according to Mc Clement [28] is the inorganic residue which provides a measure of the total amount of minerals present in food. There were significant interactions among the samples used in this study and a great difference was obtained when compared with the ash content range of 10.0% - 12.5% recorded for *Cleome gynandra* by Clement [19].

Therefore the small difference observed when these vegetables are compared with one another, and the immense difference observed when compared with the vegetables studied by another scientist above may be of the fact that, there is different concentration of minerals in the soil onto which they were planted. Crude fibre ranged from 2.43% (*S. macrocarpon*) to 22.03% (*Celosia argentea*) (Table 2 and fig.1), this fell within the range (8.50% - 20.90%) reported by Isong and Idiong [29] for some Nigerian vegetables. Dietary fibre helps to prevent constipation, bowel problems and piles. High crude fibre in the vegetable according to CFW [30] could also help in blood cholesterol attenuation, as well as blood glucose attenuation when consumed. The fibre content recorded in this study were in line with 6.0g/

100g to 6.33g/100g reported by Hassan *et al.* [31] and also in line with the result obtained for crude fibre content of Asaolu *et al.* [23] which ranges from 8.05% to 12.08%. Therefore, *Celosia argentea*, *Launaena taraxacifolia*, *Talinum triangulare* and *Ocimum canum* are good source of crude fibre as suggested by this study which could be of immense health benefit to their consumer which follows Clement [19] reports that an increased intake of dietary fibre appears to be useful in treatment of both obesity and diabetes.

The crude fat content in this study ranges from 1.20% to 3.40% (Table 2 and fig. 1), where high values were observed for *Abelmoschus esculentus*, *Amaranthus hybridus* and *Talinum triangulare*. Asaolu *et al.* [23] study, reported 3.51% to 14.02% range for crude fat in *Amaranthus hybridus*, *Indian spinach* and *Telfaira occidentalis*. Vegetables with high Dietary fats represent the most compact chemical energy available to man [32].

The protein content of the vegetables ranged from 18.50% to 55.23% with *Amaranthus hybridus* showing the highest value followed by *Launaena taraxacifolia* and *Ocimum canum* among others (Table 2 and fig. 1). It is also in accordance to the result reported by Asaolu *et al.* [23] which ranges from 46.56% to 66.60%. Plant foods that provide more than 12% of their calorific value from protein have been shown to be good source of protein [33]. This shows that all the leafy vegetables investigated are all good sources of protein. Protein help in building and maintaining all tissue in the body forms an important part of enzymes, fluid and hormones of the body and also helps form antibodies to fight against inaction and supplies energy [34]. Proteins help in building and maintaining all tissues, forms an important part of enzymes, fluids, hormones of the body and form antibodies (immunoglobulin) that fight against infections and supplies energy [19]. The protein content of vegetables in this study was high, showing that they are more nutritious.

The level of protein in the vegetables generally indicates that they are very important for human health and are good supplements for people living below the poverty level, especially in the rural areas.

CONCLUSION

The result of this research work showed that all the vegetables used in this study are more nutritious because they are very good source of protein. Also, their fibre contents were a bit low but when consumed, could correct body abnormalities such as obesity and diabetes. Also, their low fat content level indicated that, they are good for human health because they will not easily provide additional calories to the body when ingested. High ash contents indicate that, they are vital source of minerals (Na, K, Fe, Zn, Cu, Ca and P), particularly Ca could be helpful in building up strong teeth and bones, and also prevent haemophilia in blood.

Farmers should continue engaging in vegetable production and marketing. The farmers should be motivated by the government, especially the local government with provision of effective measures that could guide against any factor which may hinder the productivity of the vegetables. In addition, for sustenance of poor people in some rural areas, leafy green vegetables are very important and should therefore be an effective and efficient means of transportation to other parts of the country where productivity is low.

REFERENCES

1. Dhellot, J. R., Matouba, E., Maloumbi, M. G., Nzikou, J. M., Safou-Ngoma, D. G., Linder, M., Desobry S., Parmentie, M. 2006. Extraction, Chemical Composition and Nutritional Characterization of Vegetable Oils: Case of *Amaranthus hybridus* (Var 1 and 2) of Congo Brazzaville. *Afr. J. Biotechnol.* 5(11): pp. 1095-1101.
2. Onwordi, C. T, Anuoluwa, M. O and Adedoja, D. W. 2009. The Proximate and Mineral Compositions of three Leafy Vegetables Commonly Consumed in Lagos, Nigeria. *African Journal of Pure and Applied Chemistry.* 3 (6): pp. 102-107.

- 196 3. Ajewole, K. 1999. Analysis of the Nutritive Elements in Some Nigerian Leafy
197 Vegetables Proceedings 23rd Annual NIFST Conference, pp. 25-27 October, Abuja.
198
- 199 4. Oyedele, D. J., Asonugbo, C., Awotoye, O. O. 2006. Heavy Metals in Soil and
200 Accumulated Edible Vegetable after Phosphate Fertilizer Application. *Electron. J.*
201 *Environ. Agric. Food Chem.* 5(4): pp. 1446-1453.
202
- 203 5. Ladeji, O., Okoye, Z. S. and Ojobe, T. 1995. Chemical Evaluation of the Nutritive
204 Value of Leaf of Fluted Pumpkin (*Telferia occidentalis*). *Food Chem.*, 53: pp. 353-
205 355.
206
- 207 6. Mepba, H. D., Eboh, L. and Banigo, D. E. B. 2007. Effects of Processing Treatments
208 on the Nutritive Compositions and Consumer Acceptance of Some Nigerian Edible
209 Leafy Vegetables. *African Journal of Food, Agriculture, Nutrition and Development*
210 7(1): pp.1-18.
211
- 212 7. Sobukola, O. P., Dairo, O. U., Odunewu, A. V., and Fafiolu, B. O. 2007. Thin-layer
213 Drying Process of Some Leafy Vegetables under Open Sun. *Food Science and*
214 *Technology*, 13(1), pp. 35-40.
215
- 216 8. Mohammed, M. I., and Sharif, N. 2011. Mineral Compositions of Some Leafy
217 Vegetables Consumed in Kano, Nigeria. *Nigerian Journal of Basic and Applied*
218 *Science*, 19(2), pp. 208-211.
219
- 220 9. Martin F.W. and Ruberte Meitner L.S, (1998). *Edible Leaves of the Tropic*,
221 Educational concerns for Hunger organization, INC, pp 1 – 8.
222
- 223 10. Bosland, P. W. and Votava, E. J. 2000. Peppers: Vegetable and Spices *Capsicums*.
224 Crop Production Science in Horticulture Series No. 2, CABI Publishing, United
225 Kingdom.
226
- 227 11. Schippers, R. R. 2000. Economic and Social Importance of Indigenous African
228 Vegetables: African Indigenous Vegetables, An Overview of the Cultivated species.
229 Natural Resource Institute/ACP-EU Technical Centre for Agriculture and Rural
230 Cooperation. Chatham, U.K.
231
- 232 12. Okeno, J. A., Chebet, D. K., Mathenge, P. W. 2003. Statues of indigenous vegetables.
233 *Acta Hort. Culturae*, 621: pp. 95-100.
234
- 235 13. Smith, F. I. and Eyzaguirre, P. 2007. *African Leafy Vegetables: their Role in the*
236 *World Health Organization's Global Fruit and Vegetables Initiative* AJFAND on line,
237 7(3) ISSN 1684-5374.
238
- 239 14. WHO. 2003. Diet, Nutrition and the Prevention of Chronic Diseases. *Report of a Joint*
240 *FAO/WHO. Expert Consultation, WHO Technical Report series*, Geneva.
241
- 242 15. FAO and WHO. 2004. *Vitamin and Mineral Requirements in Human Nutrition*. 2nd
243 Edition. World Health Organization and Food and Agriculture Organization of the
244 United Nations.
245

- 246 16. FAO. 2003. United Nations Food and Agriculture Organization. Increasing fruit and
247 vegetable consumption becomes a global priority. Available from:
248 <http://fao.org/english/newsroom/focus/2003/fruitveg1.htm>
249
- 250 17. Kwenin, W. K. J, Wolli, M. and Dzomeku, B. M. 2011. Assessing the Nutritional
251 value of some African Indigenous Green Leafy Vegetables in Ghana. *Journal of*
252 *Animal and Plant Sciences*, 10,(2): pp. 1300- 1305.
253
- 254 18. Martin, E. A. and Farrell, D. J. 1998. Strategies to Improve the Nutritive Value of
255 Rice Brawn in Poultry Diets. II Changes in Oil Digestibility, Metabolisable Energy
256 and Attempts to Increase the Digestability of the Oil Fraction in the Diets of Chickens
257 and Ducklings. *Br. Poult. Sci.*, 39(4): pp. 555-559.
258
- 259 19. Clement, A. 2011. Assessment of some Traditional Leafy Vegetables of Upper East
260 Region and Influence of Stage of Harvest and Drying Method on Nutrients Content of
261 Spider Flower (*Cleome gynandra* l.). B. Sc. (Hon). Thesis submitted to the
262 Department of Horticulture, Kwame Nkrumah University of Science and Technology,
263 Kumasi, Ghana.x+136.
264
- 265 20. Shei, L. 2008. An Evaluation of Native West African Vegetables. *Agriculture and*
266 *Rural Development*.
267
- 268 21. AOAC. 2000. “*Official Methods of Analysis*”. 17th Edition. *Association of*
269 *Official Analytical Chemists*. Washington, D. C., U. S. A.
270
- 271 22. Aiyeloja A.A. and Bello O.A. 2006. Ethnobotanical Potentials of Common Herbs in
272 Nigeria:A case study of Enugu state. *Educational research and review*.1 (1); pp. 16-
273 22.
274
- 275 23. Asaolu, S. S, Adefemi, O. S, Oyakilome, I. G., Ajibulu, K. E. and Asaolu, M. F. 2012.
276 Proximate and Mineral Compositions of Nigerian Leafy Vegetables. *Journal of Food*
277 *Research*, 1(3) ISSN 1927-0887.
278
- 279 24. Idris, S.,Yisah, J. and Itodo, A. U. 2009. Proximate and Mineral Composition of the
280 Leaves of *Abelmoschus esculentus*: *International Journal of Tropical Agriculture and*
281 *food system* 3(2): pp. 116-120.
282
- 283 25. Adeniyi,S.A.,Ehiagbonare,J.E. andNwangwu, S.C.O. 2012. Nutritional Evaluation of
284 Some Staple Leafy Vegetables in Southern Nig. *Int’ Journal of Agric and Food*
285 *Science*, 2(2): pp. 37-43.
286
- 287 26. Florkowski, J. W., Robert, L. S., Bernhard, B. and Stanley, E. P. 2009. *Postharvest*
288 *Handling: A Systems Approach*, Second Edition. Elsevier Inc. Academic Press.
289
- 290 27. George, P. M. 2003. Encyclopedia of Foods. Volume 1. Humane Press; Washington.
291 p. 526.
292
- 293 28. Mc clements, D. J. 2003. *Analysis of Food Products*. Chenoweth Lab, Room 238.
294

- 295 29. Isong, E. and Idiong. 1997. Nutrient Content of the Edible Leaves of Seven Wild
296 Plants from Nigerian. *Plant foods for human nutrition*, 51: pp. 79-84.
297
- 298 30. C. F. W. 2003. All Dietary Fiber is Fundamentally Functional. C. F. W. Report.
299 Publication no. W-2003-0407-010. 128 / May-June, 2003. 48(3).
300
- 301 31. Hassan, S. W., Umar, R. A., Maishanu, H. M., Matazu, I. K., Faruk, U. Z. and Sani
302 A. A. 2007. The Effects of Drying Method on the Nutrients and Non-nutrients
303 Composition of Leaves of *Gynandropsis gynandra* (Capparaceae). *Asian Journal of*
304 *Biochemistry* 2 (5): pp. 349-353.
305
- 306 32. Kummerow, F. A. 2007. *Fats in Human Nutrition*. Department of Food Technology,
307 University of Illinois, Urbana, Illinois.
308
- 309 33. Ali, A. 2009. Proximate and Mineral Composition of the Marchubeh (*Asparagus*
310 *officinalis*). *World Dairy and Food Science*, 4(2): pp. 142-149.
311
- 312 34. Johnson, R. S. 1996. Key Nutrients. Cooperative extension service, Iowa State
313 University of Science and Technology, Ames, Iowa.
314