

PHYSICOCHEMICAL PROPERTIES OF *ALCHORNEA CORDIFOLIA*, *CYPERUS ESCULENTUM* AND *IRVINGIA GABONENSIS* SEED OILS AND THEIR APPLICATIONS

ABSTRACT

*The physicochemical properties of oils extracted from three locally available plant seeds in Nigeria namely: *Alchomea cordifolia*, *Cyperus esculentum* and *Irvingia gabonensis* using n-Hexane were determined. The results of the analysis revealed that their % yield were 37.00, 27.50 and 33.00 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. Their odour was non-offensive and their colours were reddish, light yellow and milky white for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively, making them bright and attractive. The specific gravity of the oils at 25°C was 0.91, 0.94 and 0.92 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. Their flash points in °C were also 155, 159 and 229 respectively, indicating that *I. gabonensis* is the most thermally stable oil and suitable for frying. The chemical properties for *A. cordifolia*, *C. esculentum*, and *I. gabonensis* respectively were as follows: Acid values in mgKOH/g were 24.67, 5.33 and 3.73. Peroxide values in mEqKg⁻¹ were 7.26, 9.86 and 2.96. Saponification values in mgKOH/g were 162.13, 179.52 and 238.43. Iodine values in mEqKg⁻¹ were 24.62, 11.68 and 3.38. These results indicate that the three seeds are viable sources of oil based on their % yield. They are good for both domestic and industrial use based on their acid, saponification and iodine values. Their properties in most cases compete favorably with palm kernel oil (PKO) which is currently being used for many domestic and industrial purposes in Nigeria especially for the making of paints, soap, cosmetics, lubricant and varnishes.*

Keywords: Physiochemical Properties, Seed oils, Applications

INTRODUCTION

Oil from plants is used for both domestic and industrial purposes all over the world based on its physical and chemical properties. Palm kernel oil

36 is one of the most commonly used vegetable oils because its properties
37 have been studied and known especially by researchers like Akubugwo
38 and Ugborgu, stating that is good for soap making and the production of
39 cosmetics, paints, varnishes industrially based on its saponification
40 value, iodine value etc[1]. According to Unilever [2], vegetable oils from
41 plants like oil palm tree, groundnut, olive, beniseed (*sesame*), soya
42 beans, coconut, castor seed, linseed etc. plays an important role in our
43 diet as a source of fat and oil, a major class of food required for
44 warmth and energy in the body. Besides, some of these oils are used for
45 the production of commodities like soaps, cosmetics, paints, varnishes,
46 lubricants plastics, while others are used for cooking or are prepared
47 and eaten in form of butter or margarine. Moreover, some vegetable oils
48 are now used as substitutes for petrol or diesel as fuel in automobiles in
49 the form of biodiesel or bioethanol [3]. Medicinally, Ihesie have reported
50 that sesame seed oil can be used to treat health problems like chronic
51 constipation in elders, round worms in children, dysmenorrhea (painful
52 menstruation) in women, amenorrhea, asthmatic symptoms, coughs and
53 hiccoughs, and insufficient flow of breast milk in nursing mothers by the
54 oral intake of the oil up to two tea spoons at a time [4]. Akpe have also
55 studied the physiochemical properties of *Carica papaya*, *Citrus paradisi*
56 and *Croton zambesicus* seed oils and reported that they compete

57 favourably with palm kernel oil used for several industrial purposes in
58 Nigeria [5]. Based on the facts so far, the importance of vegetable oils
59 to man cannot be over emphasized and their economic value
60 unquantifiable. However, one can observed that the oil plants or crops
61 mentioned above are a small percentage of the several hundreds of
62 plants in nature whose oil potentials have not been discovered, even
63 some that have been identified as oil seed crops are being underutilized
64 because their oil properties and potentials have not been properly
65 studied to ascertain their suitability for use domestically and or
66 industrially. Consequently, this study is aimed at determining the
67 physicochemical properties of oils extracted from *Alchornea cordifolia*
68 (Christmas bush), *Cyperus esculentum* (Tiger nut) and *Irvingia*
69 *gabonensis* (Bush mango) which are found locally in Obudu Area of
70 Cross River State and many other parts of Nigeria and have not been
71 properly studied. This is with a view to determine their potentials and
72 properties as sources of vegetable oil for domestic and industrial uses,
73 and also compare their properties with the established properties of
74 palm kernel oil from *Elaeis guinensis* (palm tree) which is popularly in
75 use now.

76 MATERIALS AND METHOD

77 **Sample collection and preparation:** Viable or healthy seeds of
78 *Alchornea cordifolia* (Christmas bush), *Cyperus esculentum* (Tiger nut)
79 and *Irvingia gabonensis* (Bush mango) were collected locally from
80 Obudu Area of Cross River State of Nigeria between February and
81 March, and were taken to the Department of Botany, University of
82 Calabar for identification of botanical names and labelling. The samples
83 were then taken to the Chemistry Department of the same University
84 where they were shelled or dehauled (where applicable), sun dried for
85 several days, wrapped in polythene bags and kept for use within one
86 month. Each of sample was crushed or ground into a paste using a
87 manual grinding machine. 100 g of the paste of each sample was
88 packed in an ashless filter paper and placed in the thimble of a Soxhlet
89 apparatus (extractor) and extracted using N-hexane as the extracting
90 solvent. At the end of the continuous extraction for about 5 to 6 hours,
91 the extracting solvent was evaporated off leaving the oil sample for
92 analysis. The percentage yield of the oil extract of each sample was
93 determined thus:

$$\% \text{ yield} = \frac{\text{weight of oil}}{\text{weight of sample}} \times 100\%$$

94 **Sample analysis:** The specific gravity of the oils was determined
95 according to the method reported by Onwuka [6] thus: A 50 mL

pycometer bottle was washed with water and detergent, rinsed and dried. The bottle was filled with distilled water and weighed. After drying the bottle of water, it was filled with the oil sample and weighed, at the room temperature of 25°C. The specific gravity was calculated thus:

$$\text{Specific gravity} = \frac{\text{weight of 50mL of oil}}{\text{weight of 50mL of water}}$$

The colour, state at room temperature and the odour were observed and perceived using the human sense organs. The flash point of the oil samples was determined following the procedure reported by [6] thus: 10 mL of the oil was poured into an evaporating dish and placed on a source of heat. A thermometer was suspended at the centre of the dish ensuring that its bulb dips inside the oil without touching the bottom of the dish. The temperature of the oil was raised gradually by regulating the source of heat. The point at which the oil began to give off a thin bluish smoke continuously (i.e. smoke point), a flame was applied using a match-stick. The temperature at which the oil started flashing when the flame is applied without supporting combustion was noted as the flash point of the oil.

The acid value was determined following the method of AOAC [7] as reported by [6] thus: 1.0 g each of the oils was dissolved in a mixture

114 obtained by mixing 25 mL diethylether and 25 mL ethanol, and titrated
115 with 0.1M NaOH using phenolphthalein as an indicator, shaking till a
116 pink colour end point which persisted for 15 seconds was observed. The
117 acid value and % free fatty acids were calculated as follows:

$$\text{Acid Value} = \frac{\text{Titre volume (mL)} \times 56.1 \times M}{\text{weight of sample}}$$

118 Where M is the molarity of NaOH (0.1M).

119 Acid value is expressed in milligram KOH per gram (mgKOH/g).

$$\% \text{ free fatty acid} = \frac{1}{2} \times \text{Acid value}$$

121 The saponification value was determined using the method of [7] as
122 described by [6] thus: 1 gram of the oil was weighed into a round bottom
123 flask and 24 mL of alcoholic potassium hydroxide solution was added. A
124 reflux condenser was attached to the flask and heated on a sandbath for 1
125 hour shaking frequently. One mL of phenolphthalein (1%) solution was
126 added and titrated while hot with 0.5M HCl to a colourless end point. A
127 blank titration was also carried out the volume at end point recorded. The
128 saponification value was calculated thus.

$$\text{Saponification value} = \frac{(X - Y) \times 56.1 M}{\text{Weight of sample}}$$

129 Where X = volume (mL) of test solution titration

130 Y = volume (mL) of blank titration

131 M = Molarity of HCl (0.5)

132 The peroxide value was determined using the method of [7] as described by
133 [6] thus: 1 ml of potassium iodide (KI) was added to 20 mL of a solution of
134 mL of (2:1) volumes of glacial acetic acid and chloroform. The result out
135 solution was added to 1.0 g of the oil sample in a clean dry conical flask. The
136 mixture was left in a dark for about 2 minutes and 30 mL of distilled water
137 was added and titrated with 0.02M sodium thiosulphate solution using 5 mL
138 starch as indicator. A blank titration was also carried out. The peroxide value
139 was calculated thus:

$$\text{Peroxide value} = (100M (Va - Vb))/W$$

140 Where W = weight of oil sample

141 Va = volume in Ml of thiosulphate used in test solution

142 Vb = volume in mL of thiosulphate used in blank solution

143 M = molarity of sodium thiosulphate (0.02).

144 The iodine value was determined using Wij's method as described by [6]
145 thus: 0.5 g of the oil sample was poured into a beaker and 10ml of carbon
146 tetrachloride was added, 20 mL of Wij's solution was added and a stopper
147 previously moisten with potassium iodide was inserted and allowed to stand
148 in the dark for 30 minutes. 15mL of potassium iodide solution (10%) was

149 added and titrated with 0.1M thiosulphate solution using starch as indicator.

150 A blank titration was also carried out. The iodine value was calculated thus:

$$Iodine\ value = \frac{(b - a) \times 12.69M}{weight\ of\ sample}$$

151 Where a = volume in mL of test titration

152 b = volume in mL of blank titration

153 M = molarity of thiosulphate (0.1)

154 **RESULTS:** The results of the physical and chemical properties are
155 presented in Table 1 and 2 respectively

156 **Table 1: Physical properties of *Alchornea cordifolia*, *Cyperus***
157 ***esculentum* and *Invingia gabonensis* seed oils.**

158

	% yield	Specific gravity	Flash point (°C)	State at 25°C (Room temperature)	Colour	Odour
<i>Alchornea cordifolia</i>	37.00±1.50	0.91±0.01	155.00± 2.00	Semi-solid	Reddish	Non-offensive
<i>Cyperus esculentum</i>	27.50± 2.01	0.94± 0.02	159.00± 2.50	Liquid	Light yellow	Non-offensive
<i>Invingia gabonensis</i>	33.00± 2.00	0.92± 0.02	229.00± 2.00	Waxy solid	Milky white	Non-offensive

159 Values reported in mean ± SD, with N = 3

160

161

162

163

164

Table 2: Chemical properties of *Alchornea cordifolia*, *Cyperus esculentum* and *Invingia gabonensis* seed oils

Seed oil	Acid value in mgKOH/g	%free fatty acids	Peroxide value in mEqKg ⁻¹	Saponification value in mgKOH/g	Iodine value in mEqKg ⁻¹
<i>Alchornea cordifolia</i>	24.67±1.25	12.34±1.30	7.26± 0.12	162.13± 2.50	24.62± 1.50
<i>Cyperus esculentum</i>	5.33± 0.15	2.67± 0.10	9.86± 0.15	179.52± 3.00	11.68± 1.25
<i>Invingia gabonensis</i>	3.73± 0.13	1.87± 0.11	2.96± 0.02	238.43± 2.50	3.38± 0.15

Values reported in mean ± SD, with N = 3

DISCUSSION

The percentage yield in Table 1 revealed that *A. cordifolia* (37.00%), *C. esculentum* (27.50%) and *I. gabonensis* (33.00 %). This shows that *A. cordifolia* has the highest yield, followed by *I. gabonensis* while *C. esculentum* has the lowest % yield. However, these values compete favorably with palm kernel oil with a % yield of 28% as reported by [1], with *A. cordifolia* and *I. gabonensis* even better as their values are higher than that of palm kernel oil that is commonly used. Thus, the 3 seed plants can be used as good sources of vegetable oil.

The flash point is the temperature at which volatile evolving from the heated oil will flash but not support combustion. It measures the thermal stability of the oil [6]. It is also an indicator for the suitability of the oil for frying [5]. The results in Table 1 shows that the flash points in

183 °C were 155, 159 and 229 for *A. cordifolia*, *C. esculentum* and *I.*
184 *gabonensis* respectively. These results indicate that *I. gabonensis* oil is a
185 better frying oil and a more thermally stable oil than *C. esculentum* and
186 *A. cordifolia* respectively.

187 The specific gravity (relative density) of the oils was 0.91, 0.94
188 and 0.92 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively,
189 all higher than 0.88 reported for palm kernel oil by [1], which is
190 commonly used industrially. All the oil samples were non-offensive in
191 their odour, *A. cordifolia* was reddish in colour, *C. esculentum* was light
192 yellow and *I. gabonensis* milky white in colour. This makes the oil
193 attractive and appealing. Their state at room temperature was semi-
194 solid, liquid and waxy solid for *A. cordifolia*, *C. esculentum* and *I.*
195 *gabonensis* respectively. This also competes with palm kernel oil (PKO)
196 which is semi-solid as reported by [1].

197 The chemical properties of the studied oils are reported in Table 2.
198 The results showed that the acid values of the oils were 24.67, 5.33 and
199 3.73 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively.
200 Also, the % free fatty acids were 12.34, 2.67 and 1.87 for *A. cordifolia*,
201 *C. esculentum* and *I. gabonensis* respectively. All these values are less
202 than PKO with an acid value of 14.04 [1]. Acid value is an indicator for

edibility of oil and suitability for use in the paint industry. *C. esculentum* and *I. gabonensis* oil are edible going by their free fatty acid value of less than 3 [8] as cited by [5]. They can also compete favorably with sesame, soya bean, sun flower and rapeseed oils with acid value of about 4 as reported by [9]. Thus, the two oils with the least acid values of 5.33 and 3.37 can be consumed directly (i.e. *C. esculentum* and *I. gabonensis* respectively). Besides, these values compete favourably with CODEX Standard Acid values for cold pressed palm oil and virgin palm oil which are 4.0 and 10.0 mgKOH/g respectively, and are consumed directly.

The peroxide values of the oils were 7.26, 9.86 and 2.96 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. It is an indicator for the deterioration of oils. Fresh oils have values less than 10 mEqkg⁻¹ and rancid oils have values ranging from 20 to 40 [6]. It is also an indicator for longer and shorter shelf life during storage, as fresh oils last longer [10]. Thus, all the 3 seed oils are fresh oils and compete favourably with 2.12 mEqKg⁻¹ reported for PKO by [1].

The saponification value is an indication that the oils have potential for use in the industry when values are high especially for soap and cosmetics [11]. Its values for the oils were 162.13, 179.52 and

238.43 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively.
I. gabonensis with the highest value has the best potential for industrial
use and compete with PKO that has a value of 246. 60 [1].

The Iodine values of the oils were 24.62, 11.68 and 3.38 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. These values indicates that all the three oils are non-drying oils because their values are less than 100, those with values between 100 and 150 are semi-drying oils while those greater than 150 are drying oils [12]. This non-drying character qualifies them for use in the paint industry [13]. However, the oils compete favorably with PKO which is also non-drying oil with an iodine value of 18.30 as stated by [1]. Based on their iodine values, the storage procedure should ensure protection from oxidative rancidity or deterioration as they contain appreciable level of unsaturated bonds.

CONCLUSION

At the end of this study, the 3 oil seeds can be classified as high yielding based on the % yield. *C. esculentum* and *I. gabonensis* are suitable for direct consumption by their free fatty acid value. Their iodine and saponification values shows they are suitable for the industrial production of soaps, cosmetics, paints etc. Their colours are

bright and attractive while their odours are non-offensive. Most of the physicochemical properties of the three seed oils studied compete favorably with palm kernel oil (PKO) and conventional seed oils like groundnut oil, soya bean, rapeseed, castor seed etc. One can therefore recommend that the 3 seed oils have potentials for development and use for domestic and industrial purposes.

REFERENCES

1. Akubuguo, I. E and Ugbogu, A. E. Physiochemical studies on oils from five selected Nigerian plant seeds. *Pakistan Journal of Nutrition* (2007) 6(1): 75-78.
2. Unilever Limited. Vegetable oil and fats. A Unilever educational booklet; Revised ordinary series, (1968) 2:1-5.
3. Ekpa, O. D. Entrepreneurial Chemistry: A Chemist's approach to wealth creation in a depressed Economy. 43rd Inaugural lecture University of Calabar, Calabar, (2008).
4. Ihesie, G. Medicinal Uses of Sesame oil. *Saturday Vanguard*, November, (2008) 29, 44-45.
5. Akpe, M. A., Oyo-Ita, I. and Dosunmu, M. I. Analysis of the physiochemical properties of Carica Papaya Citrus paradisi and croton zambesicus seed oils and their applications. *World Journal of Research and Review* (2018) 6(2): 6-8.
6. Onwuka, G.A. Food Analysis and instrumentation; Theory and practice. Naphthali Prints, Lagos, Nigeria, (2005).
7. AOAC. Official Methods of Analysis: 4th ed. Association of Official analytical chemists; V.A. Arlington, 1984.
8. Eckey, E.W. Vegetable fats and oils. Reinhold; New York, 1954.
9. Pearson, D. The Chemical Analysis of food. Churchill, Livingstone, (1976).
10. Eka, O. U. Proximate Composition of Bush Mango tree and some properties of Dika fat. *Nig. J. Nutri. Sci.* (1980); 1:33-36.

- 273 11. Amoo, I. a Eleyinmi, A.F., Oilelaboye, N. A and Akoja, S. S.
274 Characteristics of Oil extract from gourd (*Cucurbita Maxima*) Seed Food,
275 Agriculture and Environment; (2004), 2:38-39.
- 276 12. Schwartz, T. A., Bunce, M. D., Silberman, R. G., Stanizki, C. I., Starlton,
277 W. J. and Zipp, A. P. *Chemistry in context*. 2nd Edition: U.S.A, McGraw-
278 Hill companies Inc, (1997).
- 279 13. Dosunmu, M.J. and Ochu, C. Physiochemical properties and fatty acid
280 composition of lipids extracted from some Nigeria fruits and seeds.
281 Global J. Pure and applied Sci. (1995), 1:45-47.