

PHYSICOCHEMICAL PROPERTIES OF *ALCHORNEA CORDIFOLIA*, *CYPERUS ESCULENTUM* AND *IRVINGIA GABONENSIS* SEED OILS AND THEIR APPLICATIONS

ABSTRACT

The physicochemical properties of oils extracted from three locally available plant seeds in Nigeria namely: Alchomea cordifolia, Cyperus esculentum and Irvingia gabonensis using n-Hexane were determined. The results of the analysis revealed that their % yield were 37.00, 27.50 and 33.00 for A. cordifolia, C. esculentum and I. gabonensis respectively. Their odour was non-offensive and their colours were reddish, light yellow and milky white for A. cordifolia, C. esculentum and I. gabonensis respectively, making them bright and attractive. The specific gravity of the oils at 25°C was 0.91, 0.94 and 0.92 for A. cordifolia, C. esculentum and I. gabonensis respectively. Their flash points in °C were also 155, 159 and 229 respectively, indicating that I. gabonensis is the most thermally stable oil and suitable for frying. The chemical properties for A. cordifolia, C. esculentum, and I. gabonensis respectively were as follows: Acid values in mgKOH/g were 24.67, 5.33 and 3.73. Peroxide values in mEqKg⁻¹ were 7.26, 9.86 and 2.96. Saponification values in mgKOH/g were 162.13, 179.52 and 238.43. Iodine values in g/100g were 24.62, 11.68 and 3.38. These results indicate that the three seeds are viable sources of oil based on their % yield. They are good for both domestic and industrial use based on their acid, saponification and iodine values. Their properties in most cases compete favorably with palm kernel oil (PKO) which is currently being used for many domestic and industrial purposes in Nigeria especially for the making of paints, soap, cosmetics, lubricant and varnishes.

Keywords: Physiochemical Properties, Seed oils, Applications

INTRODUCTION

Palm kernel oil is one of the most commonly used vegetable oils because its properties have been studied and known especially by

36 researchers like Akubugwo and Ugborgu, stating that is good for soap
37 making and the production of cosmetics, paints, varnishes industrially
38 based on its saponification value, iodine value etc[1]. According to
39 Unilever [2], vegetable oils from plants like oil palm tree, groundnut,
40 olive, beniseed (*sesame*), soya beans, coconut, castor seed, linseed etc.
41 plays an important role in our diet as a source of fat and oil, a major
42 class of food required for warmth and energy in the body. Besides,
43 some of these oils are used for the production of commodities like
44 soaps, cosmetics, paints, varnishes, lubricants plastics, while others are
45 used for cooking or are prepared and eaten in form of butter or
46 margarine. Moreover, some vegetable oils are now used as substitutes
47 for petrol or diesel as fuel in automobiles in the form of biodiesel or bio-
48 ethanol [3]. Medicinally, Ihesie have reported that sesame seed oil can
49 be used to treat health problems like chronic constipation in elders,
50 round worms in children, dysmenorrhea (painful menstruation) in
51 women, amenorrhea, asthmatic symptoms, coughs and hiccoughs, and
52 insufficient flow of breast milk in nursing mothers by the oral intake of
53 the oil up to two tea spoons at a time [4]. Akpe have also studied the
54 physiochemical properties of *Carica papaya*, *Citrus paradisi* and *Croton*
55 *zambesicus* seed oils and reported that they compete favourably with
56 palm kernel oil used for several industrial purposes in Nigeria [5]. Thus,

57 oils from plants are used for both domestic and industrial purposes all
58 over the world based on their physical and chemical properties. Nyam
59 and others have also studied the physicochemical properties and
60 bioactive compounds of oil extracted from bitter melon, Kalahari melon,
61 kenaf, pumpkin roselle seeds and found that the oils were rich in
62 tocopherols, with γ -tocopherol as the major component in all the oil
63 samples, they also contain a great number of compounds that have a
64 potential high value as food and for production of non-food products [6].
65 Neagu and his team have also worked on the physicochemical properties
66 of sunflower oil, corn oil, rapeseed oil, and peanut oil, and reported that
67 their properties were within the requirements of food domain and their
68 density correlated with other properties of edible oils [7]. Kalias and his
69 group have researched on the physicochemical properties of soybean,
70 castor, groundnut, cotton seed oils and their blends, and the results
71 showed that higher viscosity and lower acid value of castor oil blends in
72 soybean oil suggest that they can be directly used as an alternative for
73 lubricant for mineral oil based lubricants [8]. Serjouie and his team have
74 also studied the effect of vegetable-based oil blends on physicochemical
75 properties of oils during deep-fat frying and stated that they properties
76 of the oils were significantly influenced by the type and concentration of
77 the component oils [9]. Zahir and his group also investigated the

78 physicochemical properties of edible oils and evaluated their frying
79 quality using FT-IR spectroscopy and the results revealed that due to
80 the temperature change in the oils, there is a notable difference in the
81 spectral band which showed that the proportions of fatty acids were
82 changed [10]. Angaye and Maduelosi studied the physicochemical
83 properties of some brands of edible oils namely; Turkey, Grand, Gino,
84 Tropical, Power, Mammador, local vegetable oil and raw/locally
85 extracted groundnut oil sold in some super markets and open markets in
86 Port-Harcourt, Nigeria, and compare the results with standards and
87 found that oils sold in super markets are better protected from light-
88 induced oxidation than those sold in open markets [11]. Based on the
89 facts so far, the importance of vegetable oils to man cannot be over
90 emphasized and their economic value unquantifiable. However, one can
91 observed that the oil plants or crops mentioned above are a small
92 percentage of the several hundreds of plants in nature whose oil
93 potentials have not been discovered, even some that have been
94 identified as oil seed crops are being underutilized because their oil
95 properties and potentials have not been properly studied to ascertain
96 their suitability for use domestically and or industrially. Consequently,
97 this study is aimed at determining the physicochemical properties of oils
98 extracted from *Alchornea cordifolia* (Christmas bush) which is a shrub

99 belonging to the family *Euphorbiaceae*, *Cyperus esculentum* (Tiger nut
100 or yellow nutsedge) which is a crop of the family *Cyperaceae* and
101 *Irvingia gabonensis* (Bush, wild or African mango) which is a tree of the
102 family *Irvingiaceae*, all of which are found locally in Obudu Area of
103 Cross River State and many other parts of Nigeria, Africa and some
104 other countries of the world, especially the tropics and have not been
105 properly studied. This is with a view to determine their potentials and
106 properties as sources of vegetable oil for domestic and industrial uses,
107 and also compare their properties with the established properties of
108 palm kernel oil from *Elaeis guinensis* (palm tree) which is popularly in
109 use now.

110 MATERIALS AND METHOD

111 **Sample collection and preparation:** Viable or healthy seeds of
112 *Alchornea cordifolia* (Christmas bush), *Cyperus esculentum* (Tiger nut)
113 and *Irvingia gabonensis* (Bush mango) were collected locally from
114 Obudu Area of Cross River State of Nigeria between February and
115 March, and were taken to the Department of Botany, University of
116 Calabar for identification of botanical names and labeling. The samples
117 were then taken to the Chemistry Department of the same University
118 where they were shelled or de-hauled (where applicable), sun dried for

119 several days, wrapped in polythene bags and kept for use within one
120 month. Each of the samples was crushed or ground into a paste using a
121 manual grinding machine. 100 g of the paste of each sample was
122 packed in an ash less filter paper and placed in the thimble of a Soxhlet
123 apparatus (extractor) and extracted using N-hexane as the extracting
124 solvent. At the end of the continuous extraction for about 5 to 6 hours,
125 the extracting solvent was evaporated off leaving the oil sample for
126 analysis. The percentage yield of the oil extract of each sample was
127 determined thus:

$$\% \text{ yield} = \frac{\text{weight of oil}}{\text{weight of sample}} \times 100\%$$

128 **Sample analysis:** The specific gravity of the oils was determined
129 according to the method reported by Onwuka [12] thus: A 50 mL
130 pycometer bottle was washed with water and detergent, rinsed and
131 dried. The bottle was filled with distilled water and weighed. After drying
132 the bottle of water, it was filled with the oil sample and weighed, at the
133 room temperature of 25°C. The specific gravity was calculated thus:

$$\text{Specific gravity} = \frac{\text{weight of 50mL of oil}}{\text{weight of 50mL of water}}$$

134 The colour, state at room temperature and the odour were
135 observed and perceived using the human sense organs. The flash point

of the oil samples was determined following the procedure reported by Onwuka [12] thus: 10 mL of the oil was poured into an evaporating dish and placed on a source of heat. A thermometer was suspended at the centre of the dish ensuring that its bulb dips inside the oil without touching the bottom of the dish. The temperature of the oil was raised gradually by regulating the source of heat. The point at which the oil began to give off a thin bluish smoke continuously (i.e. smoke point), a flame was applied using a match-stick. The temperature at which the oil started flashing when the flame is applied without supporting combustion was noted as the flash point of the oil.

The acid value was determined following the method of AOAC [13] as reported by Onwuka [12] thus: 1.0 g each of the oils was dissolved in a mixture obtained by mixing 25 mL diethylether and 25 mL ethanol, and titrated with 0.1M NaOH using phenolphthalein as an indicator, shaking till a pink colour end point which persisted for 15 seconds was observed. The acid value and % free fatty acids were calculated as follows:

$$\text{Acid Value} = \frac{\text{Titre volume (mL)} \times 56.1 \times M}{\text{weight of sample}}$$

Where, M is the molarity of NaOH (0.1M).

Acid value is expressed in milligram KOH per gram (mgKOH/g).

$$\% \text{ free fatty acid} = \frac{1}{2} \times \text{Acid value}$$

The saponification value was determined using the method of AOAC [13] as described by Onwuka [12] thus: 1 gram of the oil was weighed into a round bottom flask and 24 mL of alcoholic potassium hydroxide solution was added. A reflux condenser was attached to the flask and heated on a sandbath for 1 hour shaking frequently. One mL of phenolphthalein (1%) solution was added and titrated while hot with 0.5M HCl to a colourless end point. A blank titration was also carried out the volume at end point recorded. The saponification value was calculated thus.

$$\text{Saponification value} = \frac{(X - Y) \times 56.1 \text{ M}}{\text{Weight of sample}}$$

Where X = volume (mL) of test solution titration

Y = volume (mL) of blank titration

M = Molarity of HCl (0.5)

The peroxide value was determined using the method of AOAC [13] as described by Onwuka [12] thus: 1 ml of potassium iodide (KI) was added to 20 mL of a solution of mL of (2:1) volumes of glacial acetic acid and chloroform. The result out solution was added to 1.0 g of the oil sample in a clean dry conical flask. The mixture was left in a dark for about 2 minutes

and 30 mL of distilled water was added and titrated with 0.02M sodium thiosulphate solution using 5 mL starch as indicator. A blank titration was also carried out. The peroxide value was calculated thus:

$$\text{Peroxide value} = (100M (Va - Vb)/W$$

Where W = weight of oil sample

Va = volume in Ml of thiosulphate used in test solution

Vb = volume in mL of thiosulphate used in blank solution

M = molarity of sodium thiosulphate (0.02).

The iodine value was determined using Wij's method as described by Onwuka [12] thus: 0.5 g of the oil sample was poured into a beaker and 10ml of carbon tetrachloride was added, 20 mL of Wij's solution was added and a stopper previously moisten with potassium iodide was inserted and allowed to stand in the dark for 30 minutes. 15mL of potassium iodide solution (10%) was added and titrated with 0.1M thiosulphate solution using starch as indicator. A blank titration was also carried out. The iodine value was calculated thus:

$$\text{Iodine value} = \frac{(b - a) \times 12.69M}{\text{weight of sample}}$$

Where a = volume in mL of test titration

b = volume in mL of blank titration

189

M = molarity of thiosulphate (0.1)

190 **RESULTS:** The results of the physical and chemical properties are
 191 presented in Table 1 and 2 respectively

192 **Table 1: Physical properties of *Alchornea cordifolia*, *Cyperus***
 193 ***esculentum* and *Invingia gabonensis* seed oils.**

194

	% yield	Specific gravity	Flash point (°C)	State at 25°C (Room temperature)	Colour	Odour
<i>Alchornea cordifolia</i>	37.00±1.50	0.91±0.01	155.00± 2.00	Semi-solid	Reddish	Non-offensive
<i>Cyperus esculentum</i>	27.50± 2.01	0.94± 0.02	159.00± 2.50	Liquid	Light yellow	Non-offensive
<i>Invingia gabonensis</i>	33.00± 2.00	0.92± 0.02	229.00± 2.00	Waxy solid	Milky white	Non-offensive

195 Values reported in mean ± SD, with N = 3

196

197

198

199

200 **Table 2: Chemical properties of *Alchornea cordifolia*, *Cyperus***
 201 ***esculentum* and *Invingia gabonensis* seed oils**

202

Seed oil	Acid value in mgKOH/g	%free fatty acids	Peroxide value in mEqKg ⁻¹	Saponification value in mgKOH/g	Iodine value in g/100g
<i>Alchornea cordifolia</i>	24.67±1.25	12.34±1.30	7.26± 0.12	162.13± 2.50	24.62± 1.50
<i>Cyperus esculentum</i>	5.33± 0.15	2.67± 0.10	9.86± 0.15	179.52± 3.00	11.68± 1.25
<i>Invingia gabonensis</i>	3.73± 0.13	1.87± 0.11	2.96± 0.02	238.43± 2.50	3.38± 0.15

203 Values reported in mean ± SD, with N = 3

DISCUSSION

The percentage yield in Table 1 revealed that *A. cordifolia* (37.00%), *C. esculentum* (27.50%) and *I. gabonensis* (33.00 %). This shows that *A. cordifolia* has the highest yield, followed by *I. gabonensis* while *C. esculentum* has the lowest % yield. However, these values compete favorably with palm kernel oil with a % yield of 28% as reported by Akubugwo and Ugborgu [1], with *A. cordifolia* and *I. gabonensis* even better as their values are higher than that of palm kernel oil that is commonly used. Thus, the 3 seed plants can be used as good sources of vegetable oil.

The flash point is the temperature at which volatile evolving from the heated oil will flash but not support combustion. It measures the thermal stability of the oil [12]. It is also an indicator for the suitability of the oil for frying [5]. The results in Table 1 show that the flash points in °C were 155, 159 and 229 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. These results indicate that *I. gabonensis* oil is a better frying oil and more thermally stable oil than *C. esculentum* and *A. cordifolia* respectively.

The specific gravity (relative density) values of the oils were 0.91, 0.94 and 0.92 for *A. cordifolia*, *C. esculentum* and *I. gabonensis*

respectively, all higher than 0.88 reported for palm kernel oil by Akubugwo and Ugborgu [1], which is commonly used industrially. All the oil samples were non-offensive in their odour, *A. cordifolia* was reddish in colour, *C. esculentum* was light yellow and *I. gabonensis* milky white in colour. This makes the oil attractive and appealing. Their state at room temperature was semi-solid, liquid and waxy solid for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. This also competes with palm kernel oil (PKO) which is semi-solid as reported by Akubugwo and Ugborgu [1].

The chemical properties of the studied oils are reported in Table 2. The results showed that the acid values of the oils were 24.67, 5.33 and 3.73 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. Also, the % free fatty acids were 12.34, 2.67 and 1.87 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. All these values are less than PKO with an acid value of 14.04 [1]. Acid value is an indicator for edibility of oil and suitability for use in the paint industry. *C. esculentum* and *I. gabonensis* oil are edible going by their free fatty acid value of less than 3 [14] as cited by Akpe [5]. They can also compete favorably with sesame, soya bean, sun flower and rapeseed oils with acid value of about 4 as reported by Pearson [15]. Thus, the two oils with the least

acid values of 5.33 and 3.37 can be consumed directly (i.e. *C. esculentum* and *I. gabonensis* respectively). Besides, these values compete favourably with CODEX Standard Acid values for cold pressed palm oil and virgin palm oil which are 4.0 and 10.0 mgKOH/g respectively, and are consumed directly [16].

The peroxide values of the oils were 7.26, 9.86 and 2.96 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. It is an indicator for the deterioration of oils. Fresh oils have values less than 10 mEqkg⁻¹ and rancid oils have values ranging from 20 to 40 [12]. It is also an indicator for longer and shorter shelf life during storage, as fresh oils last longer [17]. Thus, all the 3 seed oils are fresh oils and they compete favourably with 2.12 mEqKg⁻¹ reported for PKO by Akubugwo and Ugborgu [1].

The saponification value is an indication that the oils have potential for use in the industry when values are high especially for soap and cosmetics [18]. Its values for the oils were 162.13, 179.52 and 238.43 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. *I. gabonensis* with the highest value has the best potential for industrial use and compete with PKO that has a value of 246.60 [1].

The Iodine values of the oils were 24.62, 11.68 and 3.38 for *A. cordifolia*, *C. esculentum* and *I. gabonensis* respectively. These values indicates that all the three oils are non-drying oils because their values are less than 100, those with values between 100 and 150 are semi-drying oils while those greater than 150 are drying oils [19]. This non-drying character qualifies them for use in the paint industry [20]. However, the oils compete favorably with PKO which is also non-drying oil with an iodine value of 18.30 as stated by Akubugwo and Ugborgu [1]. Based on their iodine values, the storage procedure should ensure protection from oxidative rancidity or deterioration as they contain appreciable level of unsaturated bonds.

Finally, the variations in the physical and chemical properties of the oils were already expected because the plants used belong to different families or genera. However, the purpose of the study was to determine their oil properties and compare them based on their suitability for domestic or industrial applications, and not the oils generally.

CONCLUSION

At the end of this study, the 3 oil seeds can be classified as high yielding based on the % yield. *C. esculentum* and *I. gabonensis* are suitable for direct consumption by their free fatty acid value. Their

iodine and saponification values show they are suitable for the industrial production of soaps, cosmetics, paints etc. Their colours are bright and attractive while their odours are non-offensive. Most of the physicochemical properties of the three seed oils studied compete favorably with palm kernel oil (PKO) and conventional seed oils like groundnut oil, soya bean, rapeseed, castor seed etc. One can therefore recommend that the 3 seed oils have potentials for development and use for domestic and industrial purposes.

REFERENCES

1. Akubugwo, I. E and Ugborugu, A. E. Physiochemical studies on oils from five selected Nigerian plant seeds. *Pakistan Journal of Nutrition* (2007) 6(1): 75-78.
2. Unilever Limited. Vegetable oil and fats. A Unilever educational booklet; Revised ordinary series, (1968) 2:1-5.
3. Ekpa, O. D. Entrepreneurial Chemistry: A Chemist's approach to wealth creation in a depressed Economy. 43rd Inaugural lecture University of Calabar, Calabar, (2008).
4. Ihesie, G. Medicinal Uses of Sesame oil. *Saturday Vanguard*, November, (2008) 29, 44-45.
5. Akpe, M. A., Oyo-Ita, I. and Dosunmu, M. I. Analysis of the physiochemical properties of Carica Papaya Citrus paradisi and croton zambesicus seed oils and their applications. *World Journal of Research and Review* (2018), 6(2): 6-8.
6. Nyam, K.L., Tan, C.P., Lai, O.M., Long, K. and Che Man, Y.B. Physicochemical properties and bioactive compounds of selected seed oils. *LWT-Food Science and Technology* (2009), 42:1396-1403.
7. Neagu, A.A., Nita, N., Botez, E. and Geaca, S. A physicochemical study for some edible oils properties. *Ovidius University Annals of Chemistry* (2016), 27(1).

8. Kalias, M.T., Mahajan, D.T., and Masand, V.H. Study on physicochemical properties of vegetable oils and their blends use as possible ecological lubricant. *Journal of Chemical and Pharmaceutical Research* (2012), 4(12): 5139-5144.
9. Serjouie, A., Tan, C.P., Mirhosseini, H. and Che Man, Y.B. Effect of vegetable-based oil blends on physicochemical properties of oils during deep-fat frying. *American Journal of Food Technology* (2010), 5(5):310-323.
10. Zahir, E., Saeed, M.A. and Yousuf, A. Study of physicochemical properties of edible oil and evaluation of frying oil quality by Fourier Transform Infrared (FT-IR) spectroscopy. *Arabian Journal of Chemistry* (2014), 10(2): S3870-S3876.
11. Angaye, S.S. and Maduelosi, N.J. Comparative study of the physicochemical properties of some refined vegetable oils sold in Mile one Market and some Departmental stores in Port-Harcourt, River State, Nigeria. *Food Science and Quality Management* (2015), 39: 16-19.
12. Onwuka, G.A. *Food Analysis and instrumentation; Theory and practice*. Naphthali Prints, Lagos, Nigeria, (2005).
13. AOAC. *Official Methods of Analysis*: 4th ed. Association of Official analytical chemists; V.A. Arlington, 1984.
14. Eckey, E.W. *Vegetable fats and oils*. Reinhold; New York, 1954.
15. Pearson, D. *The Chemical Analysis of food*. Churchill, Livingstone, (1976).
16. Codex-Stan210. *Codex Standard for fats and oils from vegetable sources*. Codex Stand.-Food and Agriculture Organisation, 1999.
17. Eka, O. U. Proximate Composition of Bush Mango tree and some properties of Dika fat. *Nig. J. Nutri. Sci.* (1980); 1:33-36.
18. Amoo, I. a Eleyinmi, A.F., Oilelaboye, N. A and Akoja, S. S. Characteristics of Oil extract from gourd (*Cucurbita Maxima*) Seed Food, *Agriculture and Environment*; (2004), 2:38-39.
19. Schwartz, T. A., Bunce, M. D., Silberman, R. G., Stanizki, C. I., Starlton, W. J. and Zipp, A. P. *Chemistry in context*. 2nd Edition: U.S.A, McGraw-Hill companies Inc, (1997).
20. Dosunmu, M.J. and Ochu, C. Physiochemical properties and fatty acid composition of lipids extracted from some Nigeria fruits and seeds. *Global J. Pure and applied Sci.* (1995), 1:45-47.