

1
2
3 Study on variation of non polar metabolites in
4 *Gossypium hirsutum* L. under water stress
5 condition using gas chromatography-mass
6 spectroscopy technique
7

8
9 **Abstract**
10

11 Present work was aimed at the study on variation of non polar metabolites content in
12 *Gossypium hirsutum* L. under water stress condition. The variation of non-polar metabolites
13 was observed by gas chromatography-mass spectrometry (GC-MS) technique. Total 17 non-
14 polar metabolites were detected in control and water stressed *G. hirsutum* leaf. The major
15 metabolites were quinoline derivative (26.37±0.29%), 2- methylhexadecan-1-ol
16 (7.47±0.07%), phytol (7.71±0.02%), myristic acid (5.94±0.04%), hexadecanol
17 (14.30±0.94%), nonadecane (1.67±0.05%) and palmitic acid (3.20±1.39%). Total 14
18 metabolites were detected in control and water stressed *G. hirsutum* stem. The major
19 metabolites were dodecene (1.67±0.11%), L-lysine (0.65±0.06%), dibutylphthalate
20 (5.06±1.88%), linoleic acid (10.26±0.07%), campesterol (0.87±0.04%) and stigmasterol
21 (1.13±0.55%). Significant variation ($P = .05$) in most of the metabolites content in leaf and
22 stem was observed during water stress. The above major metabolites played an important role
23 during water stress and can be consider as metabolites responsible for water stress tolerance
24 in *G. hirsutum* under water stress condition. Further, this study will be valuable for the better
25 understanding of overall water stress tolerance mechanism in *G. hirsutum*.

26 **Keywords:** *Gossypium hirsutum*, water stress, metabolites, gas chromatography-mass
27 spectrometry.

28 **Introduction**

29 Water stress is one of the most important environmental factor which affects crop
30 productivity and adversely affects fruit production, square and boll shedding and fiber quality
31 in cotton [1]. Moreover, water stress is considered as the single most devastating
32 environmental factor [2]. It severely affects plant development with substantial reductions in

33 crop growth rate and biomass accumulation by reduction in the cell division, root
34 proliferation, plant water and nutrient relations [3, 4].

35 Previous studies revealed that 2 to 4 °C increase in temperature and the expected 30%
36 decrease in precipitation may adversely affect crop productivity and water availability by the
37 year 2050 [5]. Thus, screening cotton varieties for resistance to water stress conditions and
38 improving cotton tolerance to this stress conditions will mitigate negative consequences of
39 this adversity. Cotton is normally not classified under water stress tolerant crop as some other
40 plants species like sorghum [6]. Nevertheless, cotton has mechanisms that make it well
41 adapted to semi-arid regions [7]. An understanding of the response of cultivars to water
42 deficits is also important to model cotton growth and estimate irrigation needs [8]. The
43 alteration of metabolites due to water stress was previously reported for plant species and
44 considered to be responsible for water stress tolerance [9, 10].

45 Lv et al. evaluated five homozygous transgenic *Gossypium hirsutum* L. plants under
46 water stress condition and the result suggested that glycine betaine may be involved in
47 osmotic adjustment in the plant [11]. Rodriguez-Uribe et al. used microarray analysis to
48 identify water deficit-responsive genes in the *G. hirsutum* under water stress conditions [12].
49 Yoo and Wendel, conducted comparative transcriptome profiling of developing *G. hirsutum*
50 fibres using RNA-Seq by Illumina sequencing [13]. Although some other aspect of the
51 changes in *G. hirsutum* under water stresses conditions have been reported. But still there was
52 need to study the non polar metabolites changes in *G. hirsutum* under water stress condition,
53 so that the metabolites responsible for water stress tolerance can be investigated. Therefore, it
54 was imperative to study the variation of non-polar metabolites in *G. hirsutum* L. plants under
55 water stressed condition. Further, the finding of this study may helpful for agriculture
56 researchers in better understanding of metabolic pathways during water stress. To the best of
57 our knowledge, this was the first study which deals with the variations of non-polar
58 metabolites content in *G. hirsutum* L. plants under water stressed condition by gas
59 chromatography-mass spectrometry (GC-MS) method.

60 **Material and Methods**

61 Cotton seeds were purchased from Central Institute for Cotton research, Regional
62 station, Coimbatore, Tamil Nadu, India. These seeds were sown in trays (52 cm x 27 cm)
63 placed in a cultivation chamber. The seedlings were transplanted into pots. After four months,
64 the best plants of approximately the same height and with the same number of leaves were

65 selected for the study. Further, these selected plants were divided into two groups. First group
66 of plants were irrigated in every 12 hour interval at room temperature and considered as
67 control plant. While second group plants were maintained in the same environment as the
68 control plants but without addition of water to the container for 4 days. This will allow the
69 pots to dry out and plants were considered as water stressed. Finally leaf and stem samples
70 were collected from each group of plants for further study.

71 Dried samples of 3g each leaves and stems were taken for extraction with hexane
72 (1:10 w/v). The solvent portion was collected by filtration and repeated five times until the
73 hexane layer became almost colourless. The separated solvent layer was concentrated under
74 reduced pressure by using rota vapour. The resulting sticky mass was stored at -5 °C. Volatile
75 trimethylsilyl (TMS) derivatives of the samples were prepared by using 3.6 mg of the sample,
76 40 µl of methoxylamine hydrochloride in GC grade pyridine (20 mg/ml). The mixture was
77 shaken for 2 h at 37 °C in a temperature controlled vortex, followed by the addition of 70 µl
78 of the N-methyl-N-(trimethylsilyl) trifluoroacetamide (MSTFA). Thereafter, the mixture was
79 further continuously shaken for 30 min at the same condition. After completion of TMS
80 derivatization 1µl of derivatized mixture was taken for GCMS analysis. The GC-MS
81 analysis was performed using a GCs-Agilent 7890 A coupled with a 5975 C MS: MS detector
82 and Electron Impact Ionization to generate mass spectra. The scan mass range was 30m/z-
83 600m/z and the total run time in minutes was 54 min.

84 The resulting GC-MS profile was analyzed using the NIST mass spectral library and
85 by matching the chromatogram with appropriate standards. The estimation of the metabolites
86 was done using the percentage peak area that appeared at the total ion chromatogram in the
87 GC-MS analysis. The molecular weights and fragmentation patterns were ascertained by use
88 of the NIST library and the Duke phytochemical data base.

89 The Mann-Whitney U test, a nonparametric test of the null hypothesis was used to
90 compare differences in metabolites content between two independent groups i.e., control and
91 water stressed leaf or stem. Statistical analysis of GC-MS data was carried out by Mann-
92 Whitney U test without normal distribution using statistical software SYSTAT version 12.0
93 (Microsoft Corp. SYSTAT Software, Inc., USA).

94 **Results and Discussion**

95 Different class of non-polar metabolites were identified from non-polar extracts of leaf and
96 stem of *G. hirsutum* (Table 1).

97 **Table1:** Mass data of GC-MS identified metabolites from control and water-stressed *G.*
 98 *hirsutum* leaf and stem.

Serial Number	tR (min)	Compound	Molecular Formula	Molecular Weight	Mass Data (m/z)
1.	11.66	Dodecene	C ₁₂ H ₂₄	168	m/z 168 (M ⁺) (6%), 97 (24%), 84 (28%), 83 (30%), 70 (48%), 56 (62%), 55 (72%), 43 (100%)
2.	17.12	Tetradecene	C ₁₄ H ₂₈	196	m/z 196 (M ⁺) (2%), 125 (8%), 111 (34%), 97 (70%), 70 (82%), 69 (100%), 55 (78%),
3.	17.45	Nonanoic acid	C ₁₂ H ₂₆ O ₂ Si	230	m/z 230 (M ⁺) (2%), 215 (70%), 129 (22%), 117 (52%), 97 (62%), 73 (100%), 75 (80%)
4.	19.75	L-Lysine	C ₁₈ H ₄₆ N ₂ O ₃ Si ₄	450	m/z 450 (M ⁺) (2%), 360 (4%), 258 (12%), 232 (34%), 172 (30%), 102 (88%), 77 (48%), 73 (100%)
5.	19.87	Caryophyllene	C ₁₅ H ₂₄	204	m/z 204 (M ⁺) (2%), 189 (24%), 147 (34%), 133 (84%), 105 (58%), 93 (74%), 69 (100%)
6.	22.36	Quinoline derivative	C ₁₈ H ₁₈ N ₂ O	278	m/z 278 (M ⁺) (16%), 264 (20%), 263 (100%), 73 (26%)
7.	24.23	2-Keto-d-gluconic acid	C ₂₁ H ₅₀ O ₇ Si ₅	554	m/z 554 (M ⁺) (2%), 437 (22%), 292 (10%), 217 (30%), 204 (72%), 73 (100%) (Me ₃ Si)
8.	24.56	Cinnamic acid	C ₁₂ H ₆ O ₂ Si	220	m/z 220 (M ⁺), (98%), 215 (72%), 132 (26%), 75 (94%), 73 (100%)
9.	25.86	Maleic acid dibutylester	C ₁₂ H ₂₀ O ₄	228	m/z 228 (M ⁺) (2%), 173 (10%), 155 (16%), 117 (42%), 57 (48%), 41 (38%), 99 (100%)
10.	26.15	Butanal	C ₁₈ H ₄₅ NO ₅ Si ₄	467	m/z 467 (M ⁺) (2%), 307 (28%), 217(20%), 160(10%), 147 (18%), 103 (64%), 73 (100%),
11.	26.39	2- Methylhexadecan-1-ol	C ₁₇ H ₃₆ O	256	m/z 256 (M ⁺) (2%), 125 (10%), 111 (22%), 97 (38%), 71 (52%), 69 (58%), 57 (100%)
12.	26.72	Octadecene	C ₁₈ H ₃₆	252	m/z 252 (M ⁺) (2%), 139 (10%), 111 (44%), 97 (89%), 83 (92%), 69 (76%), 57 (100%),
13.	27.78	Phytol	C ₂₀ H ₄₀ O	296	m/z 296 (M ⁺) (2%), 123 (28%), 95(32%), 82 (38%), 81 (46%), 71 (100%), 57 (64%)
14.	28.53	Myristic acid	C ₁₄ H ₂₈ O ₂	300	m/z 300 (M ⁺) (4%), 285 (86%), 143 (18%), 75 (100%), 73 (80%)
15.	29.61	Tridecanedial	C ₁₃ H ₂₄ O ₂	212	m/z 212 (M ⁺) (2%), 150 (18%), 109 (42%), 95 (96%), 81 (78%), 67 (84%), 55 (100%)
16.	29.94	Hexadecanol	C ₁₉ H ₄₂ OSi	314	m/z 314 (M ⁺) (2%), 300 (22%), 299 (100%), 103 (18%), 75 (50%), 73 (22%)
17.	31.12	Nonadecane	C ₁₈ H ₃₈	266	m/z 266 (M ⁺) (2%), 111 (32%), 97 (62%) 83 (64%), 57 (80%), 55 (92%), 43 (98%), 41 (100%)
18.	32.16	Quinoline Acetamide derivative	C ₂₀ H ₁₈ N ₂ O ₅	366	m/z 366 (M ⁺) (28%), 351 (26%), 235 (68%), 219 (58%), 75 (38%), 73 (100%)
19.	32.22	Palmitic acid	C ₁₉ H ₄₀ O ₂ Si	328	m/z 328 (M ⁺) (4%), 314 (6%),

					313 (34%), 201 (2%), 145 (26%), 132 (38%), 117 (72%), 75 (82%)
20.	35.87	Dibutylphthalate	C ₁₆ H ₂₂ O ₄	278	m/z 278 (M ⁺) (2%), 149 (100%), 150 (10%), 104 (6%), 41 (8%)
21.	36.05	Linoleic acid	C ₂₁ H ₄₀ O ₂ Si	352	m/z 352 (M ⁺) (6%), 337 (70%), 129 (44%), 95 (40%), 73 (100%), 54 (52%)
22.	36.14	Stearic acid	C ₁₈ H ₃₆ O ₄	284	m/z 284 (M ⁺) (4%), 145 (24%), 132 (38%), 129 (64%), 117 (72%), 75 (72%), 73 (100%)
23.	38.32	Docosene	C ₂₂ H ₄₄	308	m/z 308 (M ⁺) (2%), 139 (6%), 125 (12%), 111 (28%), 97 (62%), 69 (68%), 55 (100%)
24.	41.50	n-Eicosanol	C ₂₀ H ₄₂ O	298	m/z 298 (M ⁺) (2%), 153 (4%), 139 (6%), 125 (12%), 111 (30%), 97 (52%), 53 (60%)
25.	44.60	Diethylphthalate	C ₂₄ H ₃₈ O ₄	390	m/z 390 (M ⁺) (2%), 280 (4%), 279 (20%), 167 (40%), 149 (100%), 113 (14%), 71 (26%), 57 (38%)
26.	47.25	Nonacosanol	C ₂₉ H ₆₀ O	424	m/z 424 (M ⁺) (2%), 139 (10%), 125 (22%), 111 (38%), 97 (90%), 69 (68%), 57 (100%)
27.	48.22	Octacosanol	C ₃₁ H ₆₆ O ₂ Si	482	m/z 482 (M ⁺) (2%), 468 (12%), 467 (76%), 111 (18%), 103 (44%), 83 (34%), 75 (100%), 57 (58%)
28.	52.56	Campesterol	C ₃₁ H ₅₆ O ₂ Si	472	m/z 472 (M ⁺) (4%), 343 (28%), 257 (20%), 147 (24%), 137 (44%), 69 (74%), 73 (100%), 57 (72%)
29.	53.77	Stigmasterol	C ₃₂ H ₅₈ O ₂ Si	486	m/z 486 (M ⁺) (38%), 398 (6%), 217 (34%), 147 (36%), 129 (18%), 95 (34%),

99 **Metabolites in leaf**

100 Total 17 non-polar metabolites were detected from leaves of water stressed *G. hirsutum*. The
101 higher amount of quinoline derivative (26.37%), 2- methylhexadecan-1-ol (7.47%), phytol
102 (7.71%), myristic acid (5.94%), hexadecanol (14.30%), nonadecane(1.67%) and palmitic
103 acid (3.20%) were detected in water stressed leaves in compare to control. Moreover two
104 metabolites i.e. caryophyllene and phytol were detected only in stressed leaves.




Figure 1. Variation of major non polar metabolites in control vs water stressed *G. hirsutum* leaf

105

106

107

108

109

110

111

112

113

114

115

The higher amount of metabolites cinnamic acid (23.93%), octadecene (6.74%), quinoline acetamide derivative (1.03%) and stearic acid (2.06%) were present in control leaf in compare to stressed leaf. While the higher amount of quinoline derivative (26.37%), myristic acid (5.94%), hexadecanol (14.30%), nonadecane (1.67%) and palmitic acid (3.20%) were detected in stressed leaf in compare to control leaf. The other non-polar metabolites such as 2-keto-d-gluconic acid (7.13%), maleic acid dibutylester (1.16%), butanal (2.92%) and tridecanedial (1.63%) were detected only in control leaf. The caryophyllene (0.58%) and phytol (7.71%) were present only in stressed leaf (Table 2 and Figure 1).

116

Table 2: Variation of non-polar metabolites in control and water stressed *G. hirsutum* leaf.


Serial Number	Compound Name	Control Leaf (Area %)	Stress Leaf (Area %)
1.	Caryophyllene	ND	0.58 ± 0.02 ^a
2.	Quinoline derivative	7.70 ± 0.11 ^a	26.37 ± 0.29 ^a
3.	2-Keto-d-gluconic acid	7.13 ± 0.17 ^a	ND
4.	Cinnamic acid	23.93 ± 0.49 ^a	9.18 ± 0.11 ^a
5.	Maleic acid dibutylester	1.16 ± 0.07 ^a	ND
6.	Butanal	2.92 ± 0.24 ^a	ND
7.	2- Methylhexadecan-1-ol	1.05 ± 0.01 ^a	7.47 ± 0.07 ^a
8.	Octadecene	6.74 ± 0.38 ^a	1.64 ± 0.17 ^a
9.	Phytol	ND	7.71 ± 0.02 ^a
10.	Myristic acid	0.63 ± 0.01 ^a	5.94 ± 0.04 ^a
11.	Tridecanedial	1.63 ± 0.03 ^a	ND
12.	Hexadecanol	6.14 ± 0.24 ^a	14.30 ± 0.94 ^a
13.	Nonadecane	0.49 ± 0.05 ^a	1.67 ± 0.05 ^a
14.	QuinolineAcetamide derivative	1.03 ± 0.06 ^a	0.79 ± 0.12 ^a

15.	Palmitic acid	0.81± 0.21 ^a	3.20 ± 1.39 ^a
16.	Dibutylphthalate	1.43± 1.05	0.88 ± 0.57
17.	Stearic acid	2.06± 0.03 ^a	0.43 ± 0.21 ^a

117 Mean values ± SD (standard deviation) values of mg/gm of fresh weight. ND = Not Detected;
 118 **a** denotes statistical significance $P = .05$ between groups (control vs stress).

119 Metabolites in stem

120 Total 14 non-polar metabolites were detected from water stressed *G. hirsutum* stem (Table 3).
 121 The higher amount of L-lysine (0.65%), linoleic acid (10.26%) and campesterol (0.87%)
 122 were detected in water stressed stem in compare to control. While the other metabolites were
 123 slightly decreased than control in compare to stress stem.



124
 125

126 **Figure 3.** Variation of major non-polar metabolites in control vs water stressed *G. hirsutum*
 127 stem

128 The higher average amount of maleic acid dibutylester (0.72%) and dioctylphthalate
 129 (4.56%) were detected in control stem compare to stress stem. The higher average amount of
 130 dodecene (1.67%), L-lysine (0.65%), linoleic acid (10.26%) and campesterol (0.87%) were
 131 found in stress stem compare to control. 2- Methylhexadecan-1-ol (0.73%) was present only
 132 in control stem. Statistically significant variation ($P = .05$) in few metabolites content was
 133 found between control and water stressed *G. hirsutum* stem (Table 3, Figure 2).

134 **Table 3:** Variation of non-polar metabolites in control and water stressed *G. hirsutum* stem.

Serial Number	Compound Name	Control Stem (Area %)	Stress Stem (Area %)
1.	Dodecene	1.04 ± 0.04 ^a	1.67 ± 0.11 ^a

2.	Nonanoic acid	5.36 ± 0.24	5.24 ± 0.05
3.	L-Lysine	0.43± 0.11 ^a	0.65 ± 0.06 ^a
4.	Quinoline derivative	28.01 ±0.17	25.87± 1.16
5.	Maleic acid dibutylester	0.72± 0.11 ^a	0.51 ± 0.03 ^{sa}
6.	2- Methylhexadecan-1-ol	0.73± 0.03 ^a	ND
7.	Dibutylphthalate	4.85± 0.21	5.06 ± 1.88
8.	Linoleic acid	3.63± 0.49 ^a	10.26±0.07 ^a
9.	Docosene	3.47± 0.23	3.05 ± 0.28
10.	n-Eicosanol	2.20± 0.08	2.06 ± 0.25
11.	Diocetylphthalate	4.56± 0.07 ^a	3.77 ± 0.09 ^a
12.	Nonacosanol	0.50± 0.06	0.46 ± 0.05
13.	Campesterol	0.31± 0.04 ^a	0.87 ± 0.04 ^a
14.	Stigmasterol	0.44± 0.26	1.13 ± 0.55

135 Mean values ± SD (standard deviation) values of mg/gm of fresh weight. ND= Not Detected;
 136 **a** denotes statistical significance $P = .05$ between groups (control vs stress).

137 Mainly 2- methylhexadecan-1-ol, hexadecanol and palmitic acid in leaf while linoleic
 138 acid in stem was found to be accumulating under water stress condition. The accumulation of
 139 these metabolites was previously reported for other plant species. These metabolites were
 140 observed to be responsible for water stress tolerance [9, 10]. Moreover, plant sterol i.e.
 141 campesterol was found in high amount in stress stem. Plant sterols regulate fluidity and
 142 permeability of phospholipid bilayer [14], cell division and plant growth [15]. Sterols are also
 143 essential for synthesis of prostaglandins and leukotrienes, important component for immune
 144 system [16].

145 **Conclusion**

146 The higher content of the metabolites such as quinoline derivative, 2- methylhexadecan-1-ol,
 147 phytol, myristic acid, hexadecanol and palmitic acid was observed in the stressed leaf
 148 compare to control leaf. The high content of the metabolites such as L-lysine, linoleic acid,
 149 campesterol and stigmasterol was detected in stressed stem compare to control stem.
 150 Similarly, consumption of the metabolites i.e., cinnamic acid, octadecene, stearic acid in leaf
 151 and quinoline derivative, docosene, dioctylphthalate in stem was observed. These
 152 observations indicate that the selective accumulation and consumption of the metabolites
 153 were occurred during the water stress in *G. hirsutum* leaf and stem. It concludes that above
 154 metabolites played a crucial role during the water stress and can be considered as metabolites
 155 responsible for water stress tolerance in *G. hirsutum*.

156 **References**

157 **1.** El-Zik KM and Thaxton PM. Genetic improvement for
 158 resistance to pests and stresses in cotton. In Frisbie, RE, El-Zik, KM and Wilson, LT.

- 159 (Eds) Integrated pest management systems and cotton production. John Wiley and
160 Sons, New York, N.Y.1989; 191-224.
- 161 2. Lambers H, Chapin F, Pons T. Plant physiological ecology.
162 Springer, New York. 2008; 540.
- 163 3. Li LC, Grimshaw JM, Nielsen CP, PCM Judd Coyte, Graham
164 ID. Evolution of Wenger's concept of community of practice. Implementation
165 Science. 2009; 4:11.
- 166 4. Farooq M, Wahid A, Kobayashi N, Fujita D and Basra SMA.
167 Plant drought stress: effects, mechanisms and management. Agronomy for
168 Sustainable Development. 2009; 29: 185-212.
- 169 5. Ben-Asher JI, Tsuyuki BA, Bravdo and Sagih M. Irrigation of
170 grapevines with saline water, I. leaf area index, stomatal conductance, transpiration
171 and photosynthesis. Agriculture Water Management. 2006; 83: 13-21.
- 172 6. Poehlman ET, Tremblay A, Perusse L et al. Heredity and
173 changes in hormones and metabolic rates with short-term training. American Journal
174 of Physiology. 1986; 350: E711-717.
- 175 7. Malik MFA, Qureshi AS, Ashraf M and Ghafoor A. Genetic
176 variability of the main yield related characters in soybean. International Journal of
177 Agriculture and Biology. 2006; 8(6): 815-619
- 178 8. Pace ML, Cole JJ, Carpenter SR, and Kitchell JF. Trophic
179 cascades revealed in diverse ecosystems. Trends in Ecology and Evolution. 1999; 14:
180 483-488.
- 181 9. Witt S, Galicia L, Lisek J, Cairns J, Tiessen A, Araus JL et al.
182 Metabolic and phenotypic responses of greenhouse-grown maize hybrids to
183 experimentally controlled drought stress. Molecular Plant. 2012; 5: 401-417.
- 184 10. Charlton AJ, Donarski JA, Harrison M, Jones SA, Godward J,
185 Oehlschlager S *et al.* Responses of the pea (*Pisum sativum* L.) leaf metabolome to
186 drought stress assessed by nuclear magnetic resonance spectroscopy. Metabolomics.
187 2008; 4: 312-27.
- 188 11. Lv S, Yang A, Zhang K, Wang L and Zhang J. Increase of
189 glycinebetaine synthesis improves drought tolerance in cotton. Molecular Breeding.
190 2007; 20(3): 233-248.
- 191 12. Rodriguez-Uribe L, Abdelraheem A, Tiwari R, Sengupta-
192 Gopalan C, Hughs S.E. and Zhang J. Identification of drought-responsive genes in a
193 drought-tolerant cotton (*Gossypium hirsutum* L.) cultivar under reduced irrigation
194 field conditions and development of candidate gene markers for drought tolerance.
195 Molecular Breeding. 2014; 34(4): 1777-1796.
- 196 13. Yoo M and Wendel J. Comparative Evolutionary and
197 Developmental Dynamics of the Cotton (*Gossypium hirsutum*) Fiber Transcriptome.
198 PLoS Genetics. 2014; 10(1): e1004073.
- 199 14. Hartmann MA. Plant sterols and the membrane environment.
200 Trends Plant Science. 1998; 3(5): 170-5
- 201 15. Clouse SD, Sasse JM. Brassinosteroids: essential regulators of
202 plant growth and development. Annual Review Plant Physiology. Plant Molecular
203 Biology. 1998; 49: 427- 51.
- 204 16. Horrocks AL, Farooqui AA. Docosahexaenoic acid in the diet:
205 its importance in maintenance and restoration of neural membrane function.
206 Prostaglandins, Leukotrienes and Essential Fatty Acids. 2004; 70(4): 361-372.